



# Full wwPDB X-ray Structure Validation Report i

Jan 31, 2016 – 08:12 PM GMT

PDB ID : 1J96  
Title : Human 3alpha-HSD type 3 in Ternary Complex with NADP and Testosterone  
Authors : Nahoum, V.; Labrie, F.; Lin, S.-X.  
Deposited on : 2001-05-23  
Resolution : 1.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	91344	1442 (1.30-1.22)
Clashscore	102246	1530 (1.30-1.22)
Ramachandran outliers	100387	1467 (1.30-1.22)
Sidechain outliers	100360	1465 (1.30-1.22)
RSRZ outliers	91569	1442 (1.30-1.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	B	906	-	-	-	X
4	TES	A	903	-	-	-	X
4	TES	B	904	-	-	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6146 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3alpha-hydroxysteroid dehydrogenase type 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	2590	1657	445	477	11	0	1	0
1	B	318	2552	1636	438	467	11	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ASP	-	CLONING ARTIFACT	UNP P52895
B	1	ASP	-	CLONING ARTIFACT	UNP P52895

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub><sup>-</sup>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O P 48 21 7 17 3	0	0
3	B	1	Total C N O P 48 21 7 17 3	0	0

- Molecule 4 is TESTOSTERONE (three-letter code: TES) (formula:  $C_{19}H_{28}O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 21 19 2	0	0
4	B	1	Total C O 21 19 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	428	Total O 428 428	0	0
5	B	430	Total O 430 430	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3alpha-hydroxysteroid dehydrogenase type 3



- Molecule 1: 3alpha-hydroxysteroid dehydrogenase type 3



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.07Å    87.15Å    76.88Å 90.00°    107.36°    90.00°	Depositor
Resolution (Å)	10.00 – 1.25 14.94 – 1.01	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.25) 95.0 (14.94-1.01)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.28 (at 1.00Å)	Xtriage
Refinement program	REFMAC	Depositor
$R$ , $R_{free}$	0.181 , 0.199 0.177 , 0.196	Depositor DCC
$R_{free}$ test set	9212 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	9.0	Xtriage
Anisotropy	0.420	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 51.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.33$	Xtriage
Outliers	0 of 348000 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6146	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, TES, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.44	0/2663	1.35	36/3604 (1.0%)
1	B	0.45	0/2624	1.39	34/3551 (1.0%)
All	All	0.45	0/5287	1.37	70/7155 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

There are no bond length outliers.

All (70) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	200	ARG	NE-CZ-NH2	28.55	134.57	120.30
1	A	200	ARG	NE-CZ-NH2	23.25	131.92	120.30
1	A	66	ARG	CD-NE-CZ	19.56	150.99	123.60
1	B	127	GLU	OE1-CD-OE2	-18.99	100.51	123.30
1	B	66	ARG	CD-NE-CZ	15.80	145.72	123.60
1	A	200	ARG	NH1-CZ-NH2	-15.27	102.60	119.40
1	B	93	GLU	OE1-CD-OE2	-14.82	105.52	123.30
1	A	127	GLU	OE1-CD-OE2	-14.69	105.67	123.30
1	B	127	GLU	CG-CD-OE2	12.16	142.62	118.30
1	A	66	ARG	NE-CZ-NH2	12.04	126.32	120.30
1	B	28	GLU	OE1-CD-OE2	-11.98	108.93	123.30
1	B	28	GLU	CG-CD-OE2	11.98	142.25	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	93	GLU	OE1-CD-OE2	-11.88	109.04	123.30
1	B	291	GLU	OE1-CD-OE2	-11.75	109.20	123.30
1	B	200	ARG	NH1-CZ-NH2	-11.58	106.67	119.40
1	A	127	GLU	CG-CD-OE2	11.41	141.12	118.30
1	B	77	GLU	OE1-CD-OE2	-10.81	110.33	123.30
1	A	77	GLU	OE1-CD-OE2	-10.77	110.37	123.30
1	A	43	GLU	CG-CD-OE2	10.60	139.49	118.30
1	A	43	GLU	OE1-CD-OE2	-10.59	110.59	123.30
1	A	200	ARG	NE-CZ-NH1	10.36	125.48	120.30
1	B	36	GLU	OE1-CD-OE2	-10.22	111.03	123.30
1	A	36	GLU	OE1-CD-OE2	10.10	135.42	123.30
1	A	36	GLU	CG-CD-OE2	-9.98	98.33	118.30
1	A	275	GLN	CG-CD-OE1	9.65	140.91	121.60
1	A	66	ARG	NE-CZ-NH1	-9.19	115.71	120.30
1	B	77	GLU	CG-CD-OE2	9.04	136.39	118.30
1	B	225	GLU	CA-C-O	-8.95	101.30	120.10
1	B	36	GLU	CG-CD-OE2	8.85	135.99	118.30
1	A	225	GLU	CA-C-O	-8.76	101.70	120.10
1	B	66	ARG	NE-CZ-NH1	8.67	124.63	120.30
1	B	229	ASP	CB-CG-OD1	8.67	126.10	118.30
1	B	28	GLU	CB-CA-C	-8.52	93.35	110.40
1	A	77	GLU	CG-CD-OE2	8.05	134.40	118.30
1	B	250	ARG	NE-CZ-NH2	-8.02	116.29	120.30
1	A	226	PRO	CA-N-CD	-7.79	100.59	111.50
1	A	28	GLU	CG-CD-OE2	-7.77	102.76	118.30
1	A	28	GLU	CB-CA-C	-7.76	94.87	110.40
1	B	226	PRO	CA-N-CD	-7.76	100.64	111.50
1	B	66	ARG	NE-CZ-NH2	-7.69	116.45	120.30
1	A	275	GLN	N-CA-CB	7.38	123.88	110.60
1	A	275	GLN	CG-CD-NE2	-7.31	99.15	116.70
1	A	200	ARG	CG-CD-NE	7.23	126.98	111.80
1	B	275	GLN	CB-CG-CD	-7.08	93.19	111.60
1	B	36	GLU	CB-CG-CD	-6.82	95.80	114.20
1	A	200	ARG	CD-NE-CZ	6.72	133.01	123.60
1	A	28	GLU	OE1-CD-OE2	6.62	131.25	123.30
1	B	200	ARG	CG-CD-NE	6.38	125.20	111.80
1	B	275	GLN	CG-CD-NE2	6.28	131.77	116.70
1	A	196	TYR	CB-CG-CD1	6.22	124.73	121.00
1	A	36	GLU	N-CA-CB	-6.20	99.44	110.60
1	B	93	GLU	CG-CD-OE1	5.98	130.26	118.30
1	B	2	ASP	CB-CG-OD2	5.96	123.67	118.30
1	A	93	GLU	CG-CD-OE1	5.86	130.03	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2	ASP	CB-CA-C	-5.67	99.06	110.40
1	B	291	GLU	CG-CD-OE2	5.58	129.45	118.30
1	A	275	GLN	CB-CA-C	-5.57	99.27	110.40
1	A	263	ARG	NE-CZ-NH2	-5.47	117.56	120.30
1	B	250	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	A	231	ASN	CB-CG-OD1	5.38	132.36	121.60
1	A	2	ASP	CB-CG-OD2	-5.37	113.47	118.30
1	A	196	TYR	CB-CG-CD2	-5.27	117.84	121.00
1	A	226	PRO	N-CD-CG	5.22	111.03	103.20
1	B	57	ASN	OD1-CG-ND2	-5.13	110.10	121.90
1	B	226	PRO	N-CD-CG	5.11	110.87	103.20
1	B	216	TYR	CA-CB-CG	5.07	123.02	113.40
1	A	275	GLN	CB-CG-CD	-5.04	98.50	111.60
1	B	36	GLU	N-CA-CB	-5.01	101.59	110.60
1	A	282	GLN	OE1-CD-NE2	-5.00	110.39	121.90
1	B	275	GLN	CB-CA-C	-5.00	100.39	110.40

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	225	GLU	Mainchain,Peptide
1	B	225	GLU	Mainchain,Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2590	0	2585	29	0
1	B	2552	0	2552	24	0
2	A	4	0	3	0	0
2	B	4	0	3	0	0
3	A	48	0	25	4	0
3	B	48	0	25	3	0
4	A	21	0	28	1	0
4	B	21	0	28	4	0
5	A	428	0	0	15	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	430	0	0	7	0
All	All	6146	0	5249	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:CYS:HB3	5:A:1316:HOH:O	1.24	1.34
1:B:222:HIS:HD2	1:B:224:GLU:H	1.24	0.82
1:A:222:HIS:HD2	1:A:224:GLU:H	1.24	0.82
1:A:306:LEU:HD21	5:A:1093:HOH:O	1.82	0.78
1:A:123:LYS:HE3	5:A:1141:HOH:O	1.90	0.71
1:A:280:ASN:HD21	3:A:901:NAP:H61A	1.38	0.68
1:A:123:LYS:HG2	5:A:1141:HOH:O	1.93	0.67
1:B:1:ASP:OD2	5:B:1275:HOH:O	2.13	0.66
1:A:56:ASN:HB2	5:A:1310:HOH:O	2.01	0.60
1:A:36:GLU:HG2	5:A:1273:HOH:O	2.02	0.59
1:B:242:CYS:HB3	5:B:1253:HOH:O	2.02	0.59
1:B:123:LYS:HG3	1:B:124:PRO:HD2	1.86	0.58
1:A:57:ASN:HD22	1:A:57:ASN:C	2.07	0.58
1:A:231:ASN:OD1	5:A:1102:HOH:O	2.17	0.57
1:A:123:LYS:HG3	1:A:124:PRO:HD2	1.86	0.56
1:B:145:CYS:HB3	1:B:179:LYS:HD2	1.87	0.56
1:B:279:GLN:NE2	3:B:902:NAP:H62A	2.03	0.56
1:A:279:GLN:NE2	3:A:901:NAP:H62A	2.04	0.56
1:B:66:ARG:NH1	5:B:1005:HOH:O	2.39	0.55
1:A:222:HIS:HE1	5:A:1093:HOH:O	1.90	0.55
1:A:136:LYS:HG2	5:A:1272:HOH:O	2.08	0.53
1:B:270:LYS:O	3:B:902:NAP:H8A	2.10	0.52
1:B:149:GLU:HG3	1:B:179:LYS:HE2	1.91	0.51
1:B:123:LYS:CG	1:B:124:PRO:HD2	2.40	0.51
1:B:123:LYS:HD3	1:B:138:LEU:CD2	2.40	0.51
1:B:24:TYR:OH	4:B:904:TES:H11	2.11	0.51
1:A:270:LYS:O	3:A:901:NAP:H8A	2.11	0.50
1:A:63:LEU:HD12	5:A:1324:HOH:O	2.11	0.50
1:A:123:LYS:CG	1:A:124:PRO:HD2	2.42	0.49
1:B:24:TYR:CE1	4:B:904:TES:H193	2.47	0.49
1:B:128:VAL:HG23	4:B:904:TES:H162	1.94	0.48
1:B:123:LYS:HD2	5:B:1279:HOH:O	2.13	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:LYS:HE2	5:A:1272:HOH:O	2.14	0.47
1:A:3:SER:HB2	5:A:1123:HOH:O	2.14	0.47
1:B:250:ARG:NH1	1:B:282:GLN:HE21	2.12	0.47
1:B:123:LYS:HD3	1:B:138:LEU:HD22	1.96	0.47
1:B:66:ARG:NH2	5:B:1024:HOH:O	2.47	0.46
1:B:36:GLU:HB2	5:B:1270:HOH:O	2.16	0.46
1:A:274:GLU:OE2	1:A:278:ARG:NH2	2.49	0.45
1:A:66:ARG:NH1	5:A:1048:HOH:O	2.51	0.44
1:A:280:ASN:ND2	3:A:901:NAP:H61A	2.09	0.43
1:A:200:ARG:NH2	5:A:1265:HOH:O	2.51	0.43
1:A:246:LYS:NZ	1:B:126:GLU:OE2	2.45	0.43
1:B:129:ILE:HG13	4:B:904:TES:H151	1.99	0.42
3:B:902:NAP:O2N	3:B:902:NAP:H3D	2.20	0.42
1:A:301:ARG:HG3	1:A:303:VAL:HG23	2.02	0.42
1:A:275:GLN:HG3	5:A:1326:HOH:O	2.19	0.41
1:A:278:ARG:O	1:A:281:VAL:HG23	2.20	0.41
1:B:285:GLU:HG3	5:B:1334:HOH:O	2.19	0.41
1:A:129:ILE:CD1	4:A:903:TES:H151	2.50	0.41
1:B:274:GLU:OE2	1:B:278:ARG:NH2	2.54	0.40
1:A:57:ASN:C	1:A:57:ASN:ND2	2.74	0.40
1:B:278:ARG:O	1:B:281:VAL:HG23	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/323 (99%)	313 (98%)	7 (2%)	1 (0%)	46  17 
1	B	314/323 (97%)	307 (98%)	6 (2%)	1 (0%)	46  17 
All	All	635/646 (98%)	620 (98%)	13 (2%)	2 (0%)	46  17 

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	226	PRO
1	A	226	PRO

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	286/285 (100%)	280 (98%)	6 (2%)	61 19
1	B	282/285 (99%)	273 (97%)	9 (3%)	46 8
All	All	568/570 (100%)	553 (97%)	15 (3%)	52 12

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	VAL
1	A	57	ASN
1	A	138	LEU
1	A	281	VAL
1	A	306	LEU
1	A	320	SER
1	B	3	SER
1	B	28	GLU
1	B	36	GLU
1	B	43	GLU
1	B	138	LEU
1	B	226	PRO
1	B	281	VAL
1	B	306	LEU
1	B	320	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	105	ASN
1	A	222	HIS
1	A	231	ASN
1	A	275	GLN
1	A	279	GLN
1	A	280	ASN
1	B	105	ASN
1	B	222	HIS
1	B	279	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAP	A	901	-	42,52,52	1.50	7 (16%)	54,80,80	2.45	9 (16%)
4	TES	A	903	-	24,24,24	0.69	0	39,39,39	1.31	5 (12%)
2	ACT	A	905	-	1,3,3	6.14	1 (100%)	0,3,3	0.00	-
3	NAP	B	902	-	42,52,52	1.50	8 (19%)	54,80,80	2.58	13 (24%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	TES	B	904	-	24,24,24	2.16	7 (29%)	39,39,39	3.00	17 (43%)
2	ACT	B	906	-	1,3,3	3.67	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	901	-	-	0/27/67/67	0/5/5/5
4	TES	A	903	-	-	0/0/58/58	0/4/4/4
2	ACT	A	905	-	-	0/0/0/0	0/0/0/0
3	NAP	B	902	-	-	0/27/67/67	0/5/5/5
4	TES	B	904	-	-	0/0/58/58	0/4/4/4
2	ACT	B	906	-	-	0/0/0/0	0/0/0/0

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	906	ACT	CH3-C	-3.67	1.43	1.48
3	A	901	NAP	O5B-C5B	-3.16	1.31	1.44
3	B	902	NAP	O5B-C5B	-3.05	1.32	1.44
3	B	902	NAP	PN-O1N	-2.79	1.41	1.51
3	A	901	NAP	PN-O1N	-2.78	1.41	1.51
3	A	901	NAP	O4B-C4B	-2.68	1.38	1.45
3	B	902	NAP	O4B-C4B	-2.37	1.39	1.45
3	B	902	NAP	C3D-C4D	2.02	1.58	1.53
4	B	904	TES	O17-C17	2.16	1.47	1.43
3	B	902	NAP	C4N-C3N	2.23	1.43	1.39
4	B	904	TES	C13-C17	2.28	1.58	1.54
3	A	901	NAP	C4N-C3N	2.34	1.43	1.39
4	B	904	TES	C12-C11	2.66	1.59	1.53
3	A	901	NAP	O4D-C1D	2.75	1.44	1.41
4	B	904	TES	C8-C14	2.82	1.59	1.53
3	B	902	NAP	O4D-C1D	3.09	1.45	1.41
3	A	901	NAP	C4A-N3A	3.26	1.40	1.35
4	B	904	TES	C8-C9	3.30	1.60	1.53
3	B	902	NAP	C4A-N3A	3.40	1.40	1.35
3	A	901	NAP	C6N-N1N	3.59	1.45	1.35
3	B	902	NAP	C6N-N1N	3.83	1.45	1.35
4	B	904	TES	C11-C9	4.64	1.62	1.53
4	B	904	TES	C7-C8	6.01	1.64	1.53

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	905	ACT	CH3-C	6.14	1.57	1.48

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	902	NAP	N3A-C2A-N1A	-9.48	121.63	128.89
3	A	901	NAP	N3A-C2A-N1A	-7.46	123.18	128.89
3	B	902	NAP	O3-PN-O5D	-7.29	83.59	102.94
3	A	901	NAP	O3-PN-O5D	-7.19	83.87	102.94
4	B	904	TES	C14-C8-C9	-7.17	99.64	109.06
4	B	904	TES	C11-C9-C10	-6.63	104.32	113.11
4	B	904	TES	C6-C7-C8	-6.55	100.70	111.67
4	B	904	TES	C12-C11-C9	-5.76	103.40	113.10
4	B	904	TES	C7-C8-C9	-4.96	104.52	110.46
4	B	904	TES	C16-C15-C14	-4.53	96.01	105.12
4	B	904	TES	C15-C14-C8	-4.08	112.59	119.03
4	B	904	TES	C11-C12-C13	-3.94	105.79	112.84
4	B	904	TES	C7-C8-C14	-3.75	105.80	112.02
4	A	903	TES	C11-C9-C10	-3.38	108.64	113.11
4	B	904	TES	C13-C14-C8	-3.14	109.45	114.37
4	B	904	TES	C10-C9-C8	-3.10	108.09	112.67
4	B	904	TES	C14-C13-C17	-2.96	96.12	99.28
4	A	903	TES	C14-C13-C17	-2.77	96.32	99.28
3	B	902	NAP	C3N-C7N-N7N	-2.66	114.91	117.82
4	B	904	TES	C7-C6-C5	-2.47	107.34	111.90
3	B	902	NAP	C3N-C2N-N1N	-2.28	117.74	120.36
4	A	903	TES	C14-C8-C9	-2.23	106.12	109.06
4	B	904	TES	C12-C13-C17	-2.20	112.02	115.31
3	B	902	NAP	O4B-C4B-C5B	-2.06	101.97	109.32
3	B	902	NAP	O3-PA-O5B	2.04	108.35	102.94
4	B	904	TES	C18-C13-C14	2.07	115.80	111.75
3	B	902	NAP	O2N-PN-O1N	2.07	123.77	112.53
3	A	901	NAP	O2N-PN-O1N	2.09	123.85	112.53
3	B	902	NAP	C5N-C4N-C3N	2.14	123.03	120.33
3	B	902	NAP	C5B-C4B-C3B	2.16	123.79	115.21
4	B	904	TES	C19-C10-C1	2.19	112.72	109.43
3	A	901	NAP	C5B-C4B-C3B	2.20	123.92	115.21
4	A	903	TES	C11-C9-C8	2.21	114.96	111.74
3	A	901	NAP	O2N-PN-O3	2.48	116.36	105.09
3	A	901	NAP	O3-PA-O5B	2.62	109.88	102.94
4	B	904	TES	C16-C17-C13	2.65	106.64	104.58
3	B	902	NAP	O2N-PN-O3	2.76	117.61	105.09

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	903	TES	C7-C8-C9	2.93	113.97	110.46
3	A	901	NAP	O5B-C5B-C4B	3.32	121.36	109.12
3	B	902	NAP	O5B-C5B-C4B	3.56	122.23	109.12
3	A	901	NAP	O5D-C5D-C4D	4.08	124.17	109.12
3	B	902	NAP	O5D-C5D-C4D	4.20	124.60	109.12
3	B	902	NAP	PN-O3-PA	10.14	161.20	132.73
3	A	901	NAP	PN-O3-PA	11.04	163.74	132.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	901	NAP	4	0
4	A	903	TES	1	0
3	B	902	NAP	3	0
4	B	904	TES	4	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	323/323 (100%)	0.59	26 (8%)	15	11	7, 11, 21, 37
1	B	318/323 (98%)	0.64	26 (8%)	14	10	7, 11, 21, 32
All	All	641/646 (99%)	0.62	52 (8%)	15	11	7, 11, 21, 37

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ASP	7.7
1	B	3	SER	6.0
1	B	28	GLU	5.6
1	B	2	ASP	5.6
1	B	137	ILE	5.6
1	B	138	LEU	5.6
1	A	133	GLU	4.7
1	A	28	GLU	4.4
1	B	27	ALA	4.3
1	B	249	LYS	4.1
1	B	123	LYS	4.0
1	B	278	ARG	3.9
1	B	129	ILE	3.9
1	A	138	LEU	3.8
1	B	131	LYS	3.6
1	A	134	ASN	3.5
1	B	246	LYS	3.5
1	B	247	LYS	3.0
1	A	3	SER	3.0
1	B	275	GLN	2.9
1	A	36	GLU	2.9
1	B	128	VAL	2.9
1	A	27	ALA	2.8
1	A	248	HIS	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	123	LYS	2.8
1	A	43	GLU	2.8
1	A	278	ARG	2.7
1	B	127	GLU	2.7
1	A	1	ASP	2.7
1	A	249	LYS	2.6
1	A	2	ASP	2.5
1	A	136	LYS	2.5
1	A	29	VAL	2.5
1	A	306	LEU	2.5
1	B	248	HIS	2.5
1	A	30	PRO	2.4
1	B	320	SER	2.4
1	A	275	GLN	2.3
1	B	5	TYR	2.3
1	B	308	LEU	2.3
1	B	110	TYR	2.3
1	A	7	CYS	2.3
1	A	231	ASN	2.3
1	B	36	GLU	2.2
1	A	33	LYS	2.2
1	A	63	LEU	2.2
1	A	308	LEU	2.2
1	B	130	PRO	2.1
1	B	226	PRO	2.1
1	B	43	GLU	2.1
1	A	246	LYS	2.0
1	A	137	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	TES	B	904	21/21	0.37	0.34	5.98	34,34,35,37	0
4	TES	A	903	21/21	0.56	0.23	5.49	23,24,26,27	0
2	ACT	B	906	4/4	0.67	0.24	4.66	21,25,29,30	0
3	NAP	A	901	48/48	0.96	0.09	-0.03	8,10,14,20	0
3	NAP	B	902	48/48	0.96	0.08	-0.11	8,11,15,20	0
2	ACT	A	905	4/4	0.60	0.26	-	27,28,28,32	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.