



## wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:49 PM BST

PDB ID : 2J9I  
EMDB ID: : EMD-1290  
Title : Lengsin is a survivor of an ancient family of class I glutamine synthetases in eukaryotes that has undergone evolutionary re- engineering for a tissue-specific role in the vertebrate eye lens.  
Authors : Wyatt, K.; White, H.E.; Wang, L.; Bateman, O.A.; Slingsby, C.; Orlova, E.V.; Wistow, G.  
Deposited on : 2006-11-09  
Resolution : 17.00 Å(reported)  
Based on PDB ID : 1F52

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.

For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>

---

MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk27241


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 17.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
Ramachandran outliers	111179	726
Sidechain outliers	111093	686

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	421	81% 18% .
1	B	421	81% 18% .
1	C	421	82% 17% .
1	D	421	82% 17% .
1	E	421	82% 17% .
1	F	421	82% 17% .
1	G	421	81% 18% .
1	H	421	82% 17% .
1	I	421	81% 18% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	J	421	 81% 18% •
1	K	421	 82% 17% •
1	L	421	 81% 18% •

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 33420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	B	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	C	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	D	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	E	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	F	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	G	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	H	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	I	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	J	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	K	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		
1	L	421	Total	C	N	O	S	0	0
			2785	1731	492	554	8		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




#### • Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




#### • Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




#### • Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




#### • Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




- Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




- Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




- Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




- Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1




- Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1






- Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1

Chain K: 82% 17%



- Molecule 1: GLUTAMATE-AMMONIA LIGASE DOMAIN-CONTAINING PROTEIN 1

Chain L: 81% 18%



## 4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of images	Not provided	Depositor
Resolution determination method	Not provided	Depositor
CTF correction method	PHASE FLIPPING	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	Not provided	Depositor
Minimum defocus (nm)	170	Depositor
Maximum defocus (nm)	320	Depositor
Magnification	50000	Depositor
Image detector	Not provided	Depositor



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >2	RMSZ	# Z  >2
1	A	0.69	0/2824	1.03	9/3853 (0.2%)
1	B	0.69	0/2824	1.03	9/3853 (0.2%)
1	C	0.69	0/2824	1.03	9/3853 (0.2%)
1	D	0.69	0/2824	1.03	9/3853 (0.2%)
1	E	0.69	0/2824	1.03	9/3853 (0.2%)
1	F	0.69	0/2824	1.03	9/3853 (0.2%)
1	G	0.69	0/2824	1.03	8/3853 (0.2%)
1	H	0.69	0/2824	1.03	8/3853 (0.2%)
1	I	0.69	0/2824	1.03	9/3853 (0.2%)
1	J	0.69	0/2824	1.03	8/3853 (0.2%)
1	K	0.69	0/2824	1.03	8/3853 (0.2%)
1	L	0.69	0/2824	1.03	9/3853 (0.2%)
All	All	0.69	0/33888	1.03	104/46236 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	3
1	C	0	3
1	D	0	3
1	E	0	3
1	F	0	3
1	G	0	3
1	H	0	3
1	I	0	3
1	J	0	3
1	K	0	3
1	L	0	3
All	All	0	36

There are no bond length outliers.

The worst 5 of 104 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	201	ARG	NE-CZ-NH1	7.22	123.91	120.30
1	F	201	ARG	NE-CZ-NH1	7.22	123.91	120.30
1	A	201	ARG	NE-CZ-NH1	7.19	123.89	120.30
1	D	201	ARG	NE-CZ-NH1	7.19	123.89	120.30
1	B	201	ARG	NE-CZ-NH1	7.15	123.88	120.30

There are no chirality outliers.

5 of 36 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	362	ALA	Peptide
1	A	364	SER	Peptide
1	A	388	LEU	Peptide
1	B	362	ALA	Peptide
1	B	364	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2785	0	2714	426	0
1	B	2785	0	2714	427	0
1	C	2785	0	2714	424	0
1	D	2785	0	2714	427	0
1	E	2785	0	2714	425	0
1	F	2785	0	2714	424	0
1	G	2785	0	2708	389	0
1	H	2785	0	2708	386	0
1	I	2785	0	2708	399	0
1	J	2785	0	2708	388	0
1	K	2785	0	2708	386	0
1	L	2785	0	2708	397	0
All	All	33420	0	32532	2467	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 37.

The worst 5 of 2467 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:133:PRO:HD2	1:F:162:GLY:CA	1.22	1.66
1:E:37:PHE:HE1	1:F:185:LEU:CD2	1.02	1.64
1:A:162:GLY:CA	1:F:133:PRO:HD2	1.22	1.63
1:I:162:GLY:CA	1:J:133:PRO:HD2	1.17	1.62
1:D:133:PRO:HD2	1:E:162:GLY:CA	1.22	1.62

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/421 (100%)	373 (89%)	36 (9%)	10 (2%)	7	47
1	B	419/421 (100%)	373 (89%)	36 (9%)	10 (2%)	7	47
1	C	419/421 (100%)	373 (89%)	36 (9%)	10 (2%)	7	47
1	D	419/421 (100%)	373 (89%)	36 (9%)	10 (2%)	7	47
1	E	419/421 (100%)	373 (89%)	36 (9%)	10 (2%)	7	47
1	F	419/421 (100%)	373 (89%)	36 (9%)	10 (2%)	7	47
1	G	419/421 (100%)	372 (89%)	37 (9%)	10 (2%)	7	47
1	H	419/421 (100%)	372 (89%)	37 (9%)	10 (2%)	7	47
1	I	419/421 (100%)	372 (89%)	37 (9%)	10 (2%)	7	47
1	J	419/421 (100%)	372 (89%)	37 (9%)	10 (2%)	7	47
1	K	419/421 (100%)	372 (89%)	37 (9%)	10 (2%)	7	47
1	L	419/421 (100%)	372 (89%)	37 (9%)	10 (2%)	7	47
All	All	5028/5052 (100%)	4470 (89%)	438 (9%)	120 (2%)	12	47

5 of 120 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	360	ALA
1	B	360	ALA
1	C	360	ALA
1	D	360	ALA
1	E	360	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	B	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	C	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	D	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	E	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	F	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	G	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	H	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	I	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	J	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	K	212/212 (100%)	210 (99%)	2 (1%)	84	93
1	L	212/212 (100%)	210 (99%)	2 (1%)	84	93
All	All	2544/2544 (100%)	2520 (99%)	24 (1%)	85	93

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	214	ILE
1	G	333	LEU
1	L	214	ILE
1	F	333	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	G	214	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	54	ASN
1	G	156	GLN
1	L	54	ASN
1	F	156	GLN
1	G	54	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.