



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:14 PM GMT

PDB ID : 1JEV  
Title : OLIGO-PEPTIDE BINDING PROTEIN (OPPA) COMPLEXED WITH KWK  
Authors : Tame, J.; Wilkinson, A.J.  
Deposited on : 1996-07-03  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1031 (1.32-1.28)
Ramachandran outliers	100387	1504 (1.34-1.26)
Sidechain outliers	100360	1503 (1.34-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	517	 88% 11%
2	B	3	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IUM	A	520	-	-	X	-
3	IUM	A	522	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4748 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called OLIGO-PEPTIDE BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	517	Total	C	N	O	S	0	7	0
			4180	2677	700	798	5			

- Molecule 2 is a protein called PEPTIDE LYS TRP LYS.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	1	0
			38	27	7	4			

- Molecule 3 is URANYL (VI) ION (three-letter code: IUM) (formula: O<sub>2</sub>U).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	U	0	0
			1	1		
3	A	1	Total	U	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total U 1 1	0	0
3	A	1	Total U 1 1	0	0
3	A	1	Total U 1 1	0	0
3	A	1	Total U 1 1	0	0
3	A	1	Total U 1 1	0	0
3	A	1	Total U 1 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	518	Total O 519 519	0	1
4	B	3	Total O 3 3	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: OLIGO-PEPTIDE BINDING PROTEIN



- Molecule 2: PEPTIDE LYS TRP LYS



There are no outlier residues recorded for this chain.

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.82Å 75.33Å 70.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.30	Depositor
% Data completeness (in resolution range)	98.7 (10.00-1.30)	Depositor
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.203 , 0.226	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4748	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IUM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/4322	0.86	3/5894 (0.1%)
2	B	0.58	0/43	0.66	0/50
All	All	0.46	0/4365	0.86	3/5944 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	485	TYR	CA-CB-CG	5.91	124.62	113.40
1	A	413	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	413	ARG	NE-CZ-NH1	5.12	122.86	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4180	0	4097	35	0
2	B	38	0	47	0	0
3	A	8	0	0	6	0
4	A	519	0	0	12	1
4	B	3	0	0	0	0
All	All	4748	0	4144	41	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:519:IUM:U	4:A:528:HOH:O	0.99	1.34
3:A:520:IUM:U	4:A:531:HOH:O	1.11	1.27
3:A:522:IUM:U	4:A:532:HOH:O	1.22	1.20
3:A:520:IUM:U	4:A:530:HOH:O	1.19	1.19
3:A:522:IUM:U	4:A:533:HOH:O	1.26	1.14
1:A:160[A]:VAL:CG2	4:A:1040:HOH:O	2.12	0.96
1:A:160[A]:VAL:HG21	4:A:1040:HOH:O	1.84	0.68
1:A:288:ARG:HH21	1:A:351:ALA:HA	1.59	0.68
1:A:160[A]:VAL:HG23	4:A:1040:HOH:O	1.88	0.66
1:A:273:TYR:CZ	1:A:413:ARG:HD3	2.36	0.60
3:A:518:IUM:U	4:A:527:HOH:O	1.82	0.59
1:A:229:GLU:HB3	1:A:249:PRO:HD3	1.86	0.55
1:A:43:LEU:O	1:A:187:THR:HB	2.09	0.53
1:A:428:ASN:HA	1:A:431:LEU:HD22	1.91	0.52
1:A:301:ILE:HA	1:A:305:LYS:CG	2.40	0.51
1:A:299:ARG:HB3	1:A:313:PRO:HB3	1.94	0.50
1:A:323:ASP:O	1:A:423:PRO:HD3	2.12	0.50
1:A:237:ARG:NH1	1:A:255:LYS:HE2	2.27	0.50
1:A:115:TYR:CE1	1:A:428:ASN:HB3	2.47	0.50
1:A:254:GLN:O	1:A:257:LYS:HG3	2.14	0.48
1:A:342:GLU:HA	1:A:345:LYS:HE3	1.96	0.48
1:A:160[B]:VAL:HG12	4:A:1024:HOH:O	2.16	0.46
1:A:265:ARG:O	1:A:488:ALA:HA	2.17	0.45
1:A:302:ILE:HA	1:A:306:VAL:HB	1.97	0.45
1:A:460:THR:O	1:A:464:GLU:HG3	2.18	0.44
1:A:366:ASN:HA	1:A:395:GLN:O	2.18	0.44
1:A:260:ILE:O	1:A:264:VAL:HG23	2.18	0.43
1:A:506:ASN:HB3	4:A:1024:HOH:O	2.18	0.43
1:A:252:LEU:O	1:A:256:LEU:HG	2.18	0.43
1:A:34:VAL:HB	1:A:35:PRO:HD3	2.00	0.43
1:A:57[B]:SER:HB2	1:A:58:PRO:HD2	2.00	0.43
1:A:301:ILE:HA	1:A:305:LYS:HG2	2.00	0.43
1:A:338:GLN:HG3	1:A:339:LYS:N	2.33	0.43
1:A:501:LYS:HD2	4:A:996:HOH:O	2.19	0.42
1:A:57[A]:SER:HB3	1:A:58:PRO:HD2	2.00	0.42
1:A:12:LYS:HE3	1:A:14:THR:OG1	2.19	0.42
1:A:398:LYS:HB3	1:A:398:LYS:HE2	1.91	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:ILE:HA	1:A:305:LYS:HG3	2.01	0.42
1:A:302:ILE:HG23	1:A:374:LEU:HD22	2.01	0.41
1:A:230:VAL:HG13	1:A:252:LEU:HD11	2.02	0.40
1:A:357:LYS:HD3	1:A:357:LYS:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:563:HOH:O	4:A:597:HOH:O[3_555]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	522/517 (101%)	504 (97%)	17 (3%)	1 (0%)	52	20
2	B	1/3 (33%)	1 (100%)	0	0	100	100
All	All	523/520 (101%)	505 (97%)	17 (3%)	1 (0%)	52	20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/455 (102%)	451 (98%)	11 (2%)	57	14
2	B	4/3 (133%)	4 (100%)	0	100	100
All	All	466/458 (102%)	455 (98%)	11 (2%)	57	14

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	78	GLU
1	A	269	TYR
1	A	300	ASP
1	A	338	GLN
1	A	342	GLU
1	A	348	LEU
1	A	384	LYS
1	A	390	VAL
1	A	431	LEU
1	A	516	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	ASN
1	A	199	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 8 are modelled with single atom - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.