



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:57 AM GMT

PDB ID : 2JG9  
Title : CRYSTALLOGRAPHIC STRUCTURE OF HUMAN C1Q GLOBULAR HEADS (P1)  
Authors : Paidassi, H.; Tacnet-Delorme, P.; Garlatti, V.; Darnault, C.; Ghebrehiwet, B.; Gaboriaud, C.; Arlaud, G.J.; Frachet, P.  
Deposited on : 2007-02-09  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	134	<div> <div>2%</div> <div>83%</div> <div>16%</div> <div>..</div> </div>
1	D	134	<div> <div>3%</div> <div>78%</div> <div>20%</div> <div>.</div> </div>
2	B	136	<div> <div>5%</div> <div>76%</div> <div>19%</div> <div>..</div> </div>
2	E	136	<div> <div>5%</div> <div>70%</div> <div>26%</div> <div>..</div> </div>
3	C	131	<div> <div>%</div> <div>81%</div> <div>15%</div> <div>...</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	131	 A horizontal bar chart showing the quality of chain F. The bar is divided into segments: a small red segment at the beginning labeled '2%', a large green segment labeled '84%', a yellow segment labeled '12%', and a small grey segment at the end. Three dots '...' follow the grey segment.

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called COMPLEMENT C1Q SUBCOMPONENT SUBUNIT A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	134	Total	C	N	O	S	0	0	1
			1042	666	178	193	5			
1	D	134	Total	C	N	O	S	0	0	1
			1042	666	178	193	5			

- Molecule 2 is a protein called COMPLEMENT C1Q SUBCOMPONENT SUBUNIT B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	133	Total	C	N	O	S	0	0	1
			1028	653	179	189	7			
2	E	133	Total	C	N	O	S	0	0	1
			1028	653	179	189	7			

- Molecule 3 is a protein called COMPLEMENT C1Q SUBCOMPONENT SUBUNIT C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	129	Total	C	N	O	S	0	0	0
			1006	646	168	188	4			
3	F	129	Total	C	N	O	S	0	0	0
			1006	646	168	188	4			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	E	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	24	Total 24	O 24	0	0
5	B	10	Total 10	O 10	0	0
5	C	26	Total 26	O 26	0	0
5	D	25	Total 25	O 25	0	0
5	E	10	Total 10	O 10	0	0
5	F	14	Total 14	O 14	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT A




- Molecule 1: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT A




- Molecule 2: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT B



- Molecule 2: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT B




- Molecule 3: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT C





● Molecule 3: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT C



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.15Å 48.24Å 87.79Å 92.45° 92.65° 113.53°	Depositor
Resolution (Å)	15.00 – 1.90 44.13 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (15.00-1.90) 90.7 (44.13-1.90)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.11 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.214 , 0.260 0.217 , 0.262	Depositor DCC
$R_{free}$ test set	5284 reflections (11.11%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.0	Xtriage
Anisotropy	0.487	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 19.5	EDS
Estimated twinning fraction	0.225 for -k,-h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.38$ , $\langle L^2 \rangle = 0.20$	Xtriage
Outliers	0 of 52928 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6263	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.74	0/1070	0.81	1/1453 (0.1%)
1	D	0.73	0/1070	0.77	0/1453
2	B	0.69	0/1049	0.76	0/1419
2	E	0.70	1/1049 (0.1%)	0.77	0/1419
3	C	0.75	0/1033	0.82	2/1408 (0.1%)
3	F	0.74	0/1033	0.83	1/1408 (0.1%)
All	All	0.73	1/6304 (0.0%)	0.80	4/8560 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	135	CYS	CB-SG	-5.09	1.73	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	109	LEU	CA-CB-CG	8.73	135.39	115.30
3	C	109	LEU	CA-CB-CG	8.35	134.51	115.30
1	A	99	ARG	NE-CZ-NH2	-6.50	117.05	120.30
3	C	137	LEU	CA-CB-CG	-5.10	103.57	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1042	0	998	18	0
1	D	1042	0	998	21	0
2	B	1028	0	993	18	1
2	E	1028	0	993	27	1
3	C	1006	0	968	25	0
3	F	1006	0	968	23	0
4	B	1	0	0	0	0
4	E	1	0	0	0	0
5	A	24	0	0	4	0
5	B	10	0	0	0	0
5	C	26	0	0	1	0
5	D	25	0	0	2	0
5	E	10	0	0	1	0
5	F	14	0	0	1	0
All	All	6263	0	5918	121	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 121 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:217:ASP:OD1	3:F:217:ASP:OXT	1.53	1.25
3:C:99:GLN:OE1	3:F:97:THR:HG21	1.53	1.08
2:B:125:ASN:OD1	2:B:136:LYS:HE3	1.53	1.06
3:C:217:ASP:OXT	3:C:217:ASP:OD1	1.79	1.00
2:E:159:ARG:HG3	2:E:193:GLU:OE1	1.66	0.96

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:163:ARG:O	2:E:190:GLU:OE2[1_556]	2.12	0.08

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	132/134 (98%)	124 (94%)	7 (5%)	1 (1%)	24	11
1	D	132/134 (98%)	124 (94%)	7 (5%)	1 (1%)	24	11
2	B	131/136 (96%)	122 (93%)	8 (6%)	1 (1%)	24	11
2	E	131/136 (96%)	122 (93%)	8 (6%)	1 (1%)	24	11
3	C	127/131 (97%)	123 (97%)	4 (3%)	0	100	100
3	F	127/131 (97%)	123 (97%)	4 (3%)	0	100	100
All	All	780/802 (97%)	738 (95%)	38 (5%)	4 (0%)	34	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	223	ASP
1	A	222	SER
1	D	222	SER
2	E	223	ASP

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/117 (98%)	114 (99%)	1 (1%)	84	83
1	D	115/117 (98%)	112 (97%)	3 (3%)	54	45
2	B	109/117 (93%)	106 (97%)	3 (3%)	51	41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	109/117 (93%)	105 (96%)	4 (4%)	41	29
3	C	112/115 (97%)	109 (97%)	3 (3%)	52	43
3	F	112/115 (97%)	109 (97%)	3 (3%)	52	43
All	All	672/698 (96%)	655 (98%)	17 (2%)	55	47

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	119	GLU
1	D	157	SER
2	E	202	LYS
3	C	217	ASP
3	F	90	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	186	GLN
1	D	188	GLN
3	F	91	GLN
3	C	146	HIS
2	E	191	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	134/134 (100%)	-0.11	3 (2%) 65 68	4, 11, 27, 35	0
1	D	134/134 (100%)	-0.10	4 (2%) 54 57	4, 11, 27, 35	0
2	B	133/136 (97%)	0.20	7 (5%) 30 33	5, 16, 29, 35	0
2	E	133/136 (97%)	0.33	7 (5%) 30 33	5, 16, 29, 35	0
3	C	129/131 (98%)	-0.25	1 (0%) 87 88	3, 9, 19, 28	0
3	F	129/131 (98%)	-0.24	2 (1%) 74 78	3, 10, 19, 28	0
All	All	792/802 (98%)	-0.02	24 (3%) 54 57	3, 11, 27, 35	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	162	GLU	7.6
2	E	161	ARG	5.4
2	E	92	THR	5.4
2	B	104	ASN	5.1
2	B	92	THR	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CA	E	1224	1/1	0.99	0.04	-2.35	9,9,9,9	0
4	CA	B	1224	1/1	0.99	0.04	-3.44	9,9,9,9	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.