



wwPDB X-ray Structure Validation Summary Report

Sep 29, 2016 – 04:50 PM EDT

PDB ID : 5JQU
Title : Crystal structure of Cytochrome P450 BM3 heme domain G265F/T269V/L272
W/L322I/F405M/A406S (WIVS-FM) variant with iron(III) deuteroporphyrin
IX bound
Authors : Reynolds, E.W.; McHenry, M.W.; Cannac, F.; Gober, J.G.; Snow, C.D.; Brus-
tad, E.M.
Deposited on : 2016-05-05
Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	 64% 30% . .
1	B	471	 65% 30% . .
1	C	471	 64% 29% . .
1	D	471	 63% 32% . .
1	E	471	 69% 25% . .
1	F	471	 59% 35% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	471	 <p>%</p> <p>59% 34%</p>
1	H	471	 <p>62% 32%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 29916 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional cytochrome P450/NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	452	3554	2284	606	646	18	0	0	0
1	B	453	3578	2293	607	660	18	0	0	0
1	C	453	3552	2281	604	650	17	0	0	0
1	D	454	3485	2243	591	633	18	0	0	0
1	E	451	3579	2299	606	656	18	0	1	0
1	F	455	3543	2275	602	648	18	0	0	0
1	G	451	3546	2274	600	655	17	0	2	0
1	H	450	3571	2294	605	654	18	0	0	0

There are 112 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	PHE	GLY	engineered mutation	UNP P14779
A	269	VAL	THR	engineered mutation	UNP P14779
A	272	TRP	LEU	engineered mutation	UNP P14779
A	322	ILE	LEU	engineered mutation	UNP P14779
A	405	MET	PHE	engineered mutation	UNP P14779
A	406	SER	ALA	engineered mutation	UNP P14779
A	464	LEU	-	expression tag	UNP P14779
A	465	GLU	-	expression tag	UNP P14779
A	466	HIS	-	expression tag	UNP P14779
A	467	HIS	-	expression tag	UNP P14779
A	468	HIS	-	expression tag	UNP P14779
A	469	HIS	-	expression tag	UNP P14779
A	470	HIS	-	expression tag	UNP P14779

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	471	HIS	-	expression tag	UNP P14779
B	265	PHE	GLY	engineered mutation	UNP P14779
B	269	VAL	THR	engineered mutation	UNP P14779
B	272	TRP	LEU	engineered mutation	UNP P14779
B	322	ILE	LEU	engineered mutation	UNP P14779
B	405	MET	PHE	engineered mutation	UNP P14779
B	406	SER	ALA	engineered mutation	UNP P14779
B	464	LEU	-	expression tag	UNP P14779
B	465	GLU	-	expression tag	UNP P14779
B	466	HIS	-	expression tag	UNP P14779
B	467	HIS	-	expression tag	UNP P14779
B	468	HIS	-	expression tag	UNP P14779
B	469	HIS	-	expression tag	UNP P14779
B	470	HIS	-	expression tag	UNP P14779
B	471	HIS	-	expression tag	UNP P14779
C	265	PHE	GLY	engineered mutation	UNP P14779
C	269	VAL	THR	engineered mutation	UNP P14779
C	272	TRP	LEU	engineered mutation	UNP P14779
C	322	ILE	LEU	engineered mutation	UNP P14779
C	405	MET	PHE	engineered mutation	UNP P14779
C	406	SER	ALA	engineered mutation	UNP P14779
C	464	LEU	-	expression tag	UNP P14779
C	465	GLU	-	expression tag	UNP P14779
C	466	HIS	-	expression tag	UNP P14779
C	467	HIS	-	expression tag	UNP P14779
C	468	HIS	-	expression tag	UNP P14779
C	469	HIS	-	expression tag	UNP P14779
C	470	HIS	-	expression tag	UNP P14779
C	471	HIS	-	expression tag	UNP P14779
D	265	PHE	GLY	engineered mutation	UNP P14779
D	269	VAL	THR	engineered mutation	UNP P14779
D	272	TRP	LEU	engineered mutation	UNP P14779
D	322	ILE	LEU	engineered mutation	UNP P14779
D	405	MET	PHE	engineered mutation	UNP P14779
D	406	SER	ALA	engineered mutation	UNP P14779
D	464	LEU	-	expression tag	UNP P14779
D	465	GLU	-	expression tag	UNP P14779
D	466	HIS	-	expression tag	UNP P14779
D	467	HIS	-	expression tag	UNP P14779
D	468	HIS	-	expression tag	UNP P14779
D	469	HIS	-	expression tag	UNP P14779
D	470	HIS	-	expression tag	UNP P14779

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	471	HIS	-	expression tag	UNP P14779
E	265	PHE	GLY	engineered mutation	UNP P14779
E	269	VAL	THR	engineered mutation	UNP P14779
E	272	TRP	LEU	engineered mutation	UNP P14779
E	322	ILE	LEU	engineered mutation	UNP P14779
E	405	MET	PHE	engineered mutation	UNP P14779
E	406	SER	ALA	engineered mutation	UNP P14779
E	464	LEU	-	expression tag	UNP P14779
E	465	GLU	-	expression tag	UNP P14779
E	466	HIS	-	expression tag	UNP P14779
E	467	HIS	-	expression tag	UNP P14779
E	468	HIS	-	expression tag	UNP P14779
E	469	HIS	-	expression tag	UNP P14779
E	470	HIS	-	expression tag	UNP P14779
E	471	HIS	-	expression tag	UNP P14779
F	265	PHE	GLY	engineered mutation	UNP P14779
F	269	VAL	THR	engineered mutation	UNP P14779
F	272	TRP	LEU	engineered mutation	UNP P14779
F	322	ILE	LEU	engineered mutation	UNP P14779
F	405	MET	PHE	engineered mutation	UNP P14779
F	406	SER	ALA	engineered mutation	UNP P14779
F	464	LEU	-	expression tag	UNP P14779
F	465	GLU	-	expression tag	UNP P14779
F	466	HIS	-	expression tag	UNP P14779
F	467	HIS	-	expression tag	UNP P14779
F	468	HIS	-	expression tag	UNP P14779
F	469	HIS	-	expression tag	UNP P14779
F	470	HIS	-	expression tag	UNP P14779
F	471	HIS	-	expression tag	UNP P14779
G	265	PHE	GLY	engineered mutation	UNP P14779
G	269	VAL	THR	engineered mutation	UNP P14779
G	272	TRP	LEU	engineered mutation	UNP P14779
G	322	ILE	LEU	engineered mutation	UNP P14779
G	405	MET	PHE	engineered mutation	UNP P14779
G	406	SER	ALA	engineered mutation	UNP P14779
G	464	LEU	-	expression tag	UNP P14779
G	465	GLU	-	expression tag	UNP P14779
G	466	HIS	-	expression tag	UNP P14779
G	467	HIS	-	expression tag	UNP P14779
G	468	HIS	-	expression tag	UNP P14779
G	469	HIS	-	expression tag	UNP P14779
G	470	HIS	-	expression tag	UNP P14779

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	E	1	Total	C	Fe	N	O	0	0
			39	30	1	4	4		
2	F	1	Total	C	Fe	N	O	0	0
			39	30	1	4	4		
2	G	1	Total	C	Fe	N	O	0	0
			39	30	1	4	4		
2	H	1	Total	C	Fe	N	O	0	0
			39	30	1	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	149	Total	O	0	0
			149	149		
3	B	189	Total	O	0	0
			189	189		
3	C	165	Total	O	0	0
			165	165		
3	D	112	Total	O	0	0
			112	112		
3	E	155	Total	O	0	0
			155	155		
3	F	117	Total	O	0	0
			117	117		
3	G	138	Total	O	0	0
			138	138		
3	H	171	Total	O	0	0
			171	171		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase




- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase




- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase




- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase




- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase





• Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase

Chain H: 62% 32%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.06Å 166.30Å 229.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.85 – 2.16 28.85 – 2.16	Depositor EDS
% Data completeness (in resolution range)	98.8 (28.85-2.16) 98.9 (28.85-2.16)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.16Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.258 , 0.334 0.257 , 0.332	Depositor DCC
R_{free} test set	2006 reflections (0.94%)	DCC
Wilson B-factor (Å ²)	16.0	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.21 , 26.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	29916	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: FDE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/3640	0.70	3/4939 (0.1%)
1	B	0.56	0/3665	0.72	1/4972 (0.0%)
1	C	0.53	0/3637	0.72	3/4936 (0.1%)
1	D	0.51	0/3571	0.69	2/4855 (0.0%)
1	E	0.54	1/3668 (0.0%)	0.70	1/4974 (0.0%)
1	F	0.54	2/3629 (0.1%)	0.68	1/4930 (0.0%)
1	G	0.53	0/3638	0.70	1/4939 (0.0%)
1	H	0.51	0/3657	0.70	0/4955
All	All	0.53	3/29105 (0.0%)	0.70	12/39500 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	156	CYS	CB-SG	-6.66	1.71	1.82
1	F	400	CYS	CB-SG	-6.36	1.71	1.82
1	F	156	CYS	CB-SG	-5.72	1.72	1.81

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	20	LEU	CB-CG-CD2	-6.87	99.32	111.00
1	C	168	ASP	C-N-CA	6.84	138.80	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	232	ASP	CB-CG-OD1	6.46	124.11	118.30
1	A	356	LEU	CA-CB-CG	6.11	129.36	115.30
1	C	71	LEU	CA-CB-CG	5.79	128.62	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	2	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3554	0	3443	121	0
1	B	3578	0	3463	114	0
1	C	3552	0	3444	109	0
1	D	3485	0	3286	116	0
1	E	3579	0	3482	90	0
1	F	3543	0	3387	138	0
1	G	3546	0	3392	143	0
1	H	3571	0	3480	121	0
2	A	39	0	25	2	0
2	B	39	0	25	2	0
2	C	39	0	25	2	0
2	D	39	0	25	2	0
2	E	39	0	25	3	0
2	F	39	0	25	1	0
2	G	39	0	25	4	0
2	H	39	0	25	3	0
3	A	149	0	0	27	0
3	B	189	0	0	21	0
3	C	165	0	0	19	1
3	D	112	0	0	10	0
3	E	155	0	0	10	2
3	F	117	0	0	12	0
3	G	138	0	0	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	171	0	0	23	1
All	All	29916	0	27577	941	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 941 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:272:TRP:CD1	1:G:322:ILE:CD1	2.39	1.05
1:G:272:TRP:NE1	1:G:322:ILE:HD13	1.75	1.02
1:G:391:LYS:NZ	3:G:601:HOH:O	1.92	1.02
1:A:68:ASP:HB2	1:A:336:LYS:NZ	1.75	1.01
1:G:272:TRP:CD1	1:G:322:ILE:HD13	1.99	0.96

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:715:HOH:O	3:H:734:HOH:O 2_455	2.02	0.18
3:C:741:HOH:O	3:E:719:HOH:O 2_354	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	448/471 (95%)	429 (96%)	18 (4%)	1 (0%)	52	51
1	B	451/471 (96%)	437 (97%)	13 (3%)	1 (0%)	52	51
1	C	449/471 (95%)	426 (95%)	21 (5%)	2 (0%)	39	34
1	D	450/471 (96%)	427 (95%)	18 (4%)	5 (1%)	17	10

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	448/471 (95%)	426 (95%)	19 (4%)	3 (1%)	26	18
1	F	451/471 (96%)	427 (95%)	19 (4%)	5 (1%)	17	10
1	G	449/471 (95%)	431 (96%)	17 (4%)	1 (0%)	52	51
1	H	446/471 (95%)	428 (96%)	18 (4%)	0	100	100
All	All	3592/3768 (95%)	3431 (96%)	143 (4%)	18 (0%)	34	26

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	247	GLU
1	E	225	ALA
1	E	311	LEU
1	F	224	LYS
1	D	245	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/415 (88%)	348 (95%)	19 (5%)	29	23
1	B	374/415 (90%)	361 (96%)	13 (4%)	43	41
1	C	368/415 (89%)	351 (95%)	17 (5%)	33	29
1	D	343/415 (83%)	330 (96%)	13 (4%)	40	37
1	E	375/415 (90%)	358 (96%)	17 (4%)	34	30
1	F	360/415 (87%)	337 (94%)	23 (6%)	22	15
1	G	364/415 (88%)	344 (94%)	20 (6%)	27	21
1	H	375/415 (90%)	356 (95%)	19 (5%)	29	24
All	All	2926/3320 (88%)	2785 (95%)	141 (5%)	31	27

5 of 141 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	22	THR
1	E	445	VAL
1	H	212	MET
1	E	102	ILE
1	E	266	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	110	GLN
1	E	266	HIS
1	F	387	GLN
1	D	403	GLN
1	F	359	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FDE	A	501	1,3	29,46,46	6.12	15 (51%)	20,76,76	4.21	12 (60%)
2	FDE	B	501	1	29,46,46	6.43	15 (51%)	20,76,76	3.51	14 (70%)
2	FDE	C	501	1	29,46,46	6.23	16 (55%)	20,76,76	3.85	13 (65%)
2	FDE	D	501	1	29,46,46	6.14	15 (51%)	20,76,76	3.77	10 (50%)
2	FDE	E	501	1,3	29,46,46	6.25	15 (51%)	20,76,76	3.85	11 (55%)
2	FDE	F	501	1	29,46,46	6.08	14 (48%)	20,76,76	4.25	12 (60%)
2	FDE	G	501	1	29,46,46	6.23	16 (55%)	20,76,76	3.88	13 (65%)
2	FDE	H	501	1	29,46,46	6.56	15 (51%)	20,76,76	3.88	13 (65%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FDE	A	501	1,3	-	0/6/86/86	0/0/8/8
2	FDE	B	501	1	-	0/6/86/86	0/0/8/8
2	FDE	C	501	1	-	0/6/86/86	0/0/8/8
2	FDE	D	501	1	-	0/6/86/86	0/0/8/8
2	FDE	E	501	1,3	-	0/6/86/86	0/0/8/8
2	FDE	F	501	1	-	0/6/86/86	0/0/8/8
2	FDE	G	501	1	-	0/6/86/86	0/0/8/8
2	FDE	H	501	1	-	0/6/86/86	0/0/8/8

The worst 5 of 121 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	501	FDE	C1C-NC	-13.94	1.30	1.49
2	E	501	FDE	C1C-NC	-12.88	1.31	1.49
2	C	501	FDE	C1C-NC	-12.35	1.32	1.49
2	B	501	FDE	C1C-NC	-12.32	1.32	1.49
2	G	501	FDE	C1C-NC	-12.28	1.32	1.49

The worst 5 of 98 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	501	FDE	CBD-CAD-C3D	-10.44	94.14	112.49
2	C	501	FDE	CMF-C3C-C2C	-7.50	116.67	128.11
2	A	501	FDE	CMF-C3C-C2C	-6.72	117.84	128.11
2	E	501	FDE	CMF-C3C-C2C	-6.72	117.85	128.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	501	FDE	CMF-C3C-C2C	-6.66	117.94	128.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FDE	2	0
2	B	501	FDE	2	0
2	C	501	FDE	2	0
2	D	501	FDE	2	0
2	E	501	FDE	3	0
2	F	501	FDE	1	0
2	G	501	FDE	4	0
2	H	501	FDE	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/471 (95%)	0.14	5 (1%) 82 86	4, 15, 28, 35	0
1	B	453/471 (96%)	0.01	0 100 100	2, 10, 25, 36	0
1	C	453/471 (96%)	0.02	5 (1%) 82 86	2, 11, 25, 37	0
1	D	454/471 (96%)	0.30	12 (2%) 59 68	10, 21, 33, 42	0
1	E	451/471 (95%)	0.06	5 (1%) 82 86	4, 12, 26, 37	0
1	F	455/471 (96%)	0.20	10 (2%) 65 73	7, 19, 32, 38	0
1	G	451/471 (95%)	0.10	5 (1%) 82 86	4, 16, 32, 42	0
1	H	450/471 (95%)	0.10	2 (0%) 93 94	3, 13, 26, 36	0
All	All	3619/3768 (96%)	0.12	44 (1%) 81 85	2, 15, 29, 42	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	THR	6.5
1	F	191	ALA	4.7
1	G	379	PHE	3.8
1	G	11	PHE	3.8
1	D	173	PHE	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FDE	H	501	39/39	0.94	0.18	1.33	2,2,7,7	0
2	FDE	C	501	39/39	0.93	0.19	1.22	2,2,7,10	0
2	FDE	A	501	39/39	0.82	0.19	1.08	5,10,16,127	0
2	FDE	E	501	39/39	0.95	0.17	0.98	2,4,18,24	0
2	FDE	B	501	39/39	0.95	0.18	0.94	2,3,11,16	0
2	FDE	G	501	39/39	0.94	0.17	0.87	3,6,11,16	0
2	FDE	D	501	39/39	0.94	0.18	0.66	5,10,15,17	0
2	FDE	F	501	39/39	0.94	0.15	0.04	4,8,18,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.