



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:57 PM GMT

PDB ID : 4JTC
Title : Crystal structure of Kv1.2-2.1 paddle chimera channel in complex with Charyb-
dotoxin in Cs+
Authors : Banerjee, A.; Lee, A.; Campbell, E.; MacKinnon, R.
Deposited on : 2013-03-23
Resolution : 2.56 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3324 (2.60-2.52)
Clashscore	102246	3729 (2.60-2.52)
Ramachandran outliers	100387	3673 (2.60-2.52)
Sidechain outliers	100360	3673 (2.60-2.52)
RSRZ outliers	91569	3333 (2.60-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	<div> <div>2%</div> <div>79%</div> <div>17%</div> <div>..</div> </div>
1	G	333	<div> <div>2%</div> <div>78%</div> <div>19%</div> <div>..</div> </div>
2	B	514	<div> <div>13%</div> <div>48%</div> <div>25%</div> <div>•</div> <div>25%</div> </div>
2	H	514	<div> <div>32%</div> <div>37%</div> <div>31%</div> <div>•</div> <div>29%</div> </div>
3	Y	37	<div> <div>97%</div> <div>59%</div> <div>38%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAP	G	1001	-	-	-	X
6	PGW	B	504	-	-	-	X
6	PGW	B	505	-	-	-	X
6	PGW	B	509	-	-	-	X
6	PGW	B	510	-	-	-	X
6	PGW	B	513	-	-	-	X
6	PGW	B	515	-	-	-	X
6	PGW	B	516	-	-	-	X
6	PGW	B	517	-	-	-	X
6	PGW	B	518	-	-	-	X
6	PGW	H	504	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11770 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Voltage-gated potassium channel subunit beta-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	0	0	0
			2556	1627	443	470	16			
1	G	326	Total	C	N	O	S	0	0	0
			2556	1627	443	470	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	MET	-	EXPRESSION TAG	UNP P62483
G	35	MET	-	EXPRESSION TAG	UNP P62483

- Molecule 2 is a protein called Potassium voltage-gated channel subfamily A member 2, Potassium voltage-gated channel subfamily B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	386	Total	C	N	O	S	0	0	0
			3088	2022	504	548	14			
2	H	363	Total	C	N	O	S	0	0	0
			2959	1950	478	518	13			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	MET	-	EXPRESSION TAG	UNP P63142
B	-17	ALA	-	EXPRESSION TAG	UNP P63142
B	-16	HIS	-	EXPRESSION TAG	UNP P63142
B	-15	HIS	-	EXPRESSION TAG	UNP P63142
B	-14	HIS	-	EXPRESSION TAG	UNP P63142
B	-13	HIS	-	EXPRESSION TAG	UNP P63142
B	-12	HIS	-	EXPRESSION TAG	UNP P63142
B	-11	HIS	-	EXPRESSION TAG	UNP P63142
B	-10	HIS	-	EXPRESSION TAG	UNP P63142

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	HIS	-	EXPRESSION TAG	UNP P63142
B	-8	HIS	-	EXPRESSION TAG	UNP P63142
B	-7	HIS	-	EXPRESSION TAG	UNP P63142
B	-6	GLY	-	EXPRESSION TAG	UNP P63142
B	-5	LEU	-	EXPRESSION TAG	UNP P63142
B	-4	VAL	-	EXPRESSION TAG	UNP P63142
B	-3	PRO	-	EXPRESSION TAG	UNP P63142
B	-2	ARG	-	EXPRESSION TAG	UNP P63142
B	-1	GLY	-	EXPRESSION TAG	UNP P63142
B	0	SER	-	EXPRESSION TAG	UNP P63142
B	31	SER	CYS	ENGINEERED MUTATION	UNP P63142
B	32	SER	CYS	ENGINEERED MUTATION	UNP P63142
B	207	GLN	ASN	ENGINEERED MUTATION	UNP P63142
B	431	SER	CYS	ENGINEERED MUTATION	UNP P63142
B	478	SER	CYS	ENGINEERED MUTATION	UNP P63142
H	-18	MET	-	EXPRESSION TAG	UNP P63142
H	-17	ALA	-	EXPRESSION TAG	UNP P63142
H	-16	HIS	-	EXPRESSION TAG	UNP P63142
H	-15	HIS	-	EXPRESSION TAG	UNP P63142
H	-14	HIS	-	EXPRESSION TAG	UNP P63142
H	-13	HIS	-	EXPRESSION TAG	UNP P63142
H	-12	HIS	-	EXPRESSION TAG	UNP P63142
H	-11	HIS	-	EXPRESSION TAG	UNP P63142
H	-10	HIS	-	EXPRESSION TAG	UNP P63142
H	-9	HIS	-	EXPRESSION TAG	UNP P63142
H	-8	HIS	-	EXPRESSION TAG	UNP P63142
H	-7	HIS	-	EXPRESSION TAG	UNP P63142
H	-6	GLY	-	EXPRESSION TAG	UNP P63142
H	-5	LEU	-	EXPRESSION TAG	UNP P63142
H	-4	VAL	-	EXPRESSION TAG	UNP P63142
H	-3	PRO	-	EXPRESSION TAG	UNP P63142
H	-2	ARG	-	EXPRESSION TAG	UNP P63142
H	-1	GLY	-	EXPRESSION TAG	UNP P63142
H	0	SER	-	EXPRESSION TAG	UNP P63142
H	31	SER	CYS	ENGINEERED MUTATION	UNP P63142
H	32	SER	CYS	ENGINEERED MUTATION	UNP P63142
H	207	GLN	ASN	ENGINEERED MUTATION	UNP P63142
H	431	SER	CYS	ENGINEERED MUTATION	UNP P63142
H	478	SER	CYS	ENGINEERED MUTATION	UNP P63142


- Molecule 3 is a protein called Potassium channel toxin alpha-KTx 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Y	37	Total	C	N	O	S	0	0	0
			295	176	57	55	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	1	PCA	GLN	MODIFIED RESIDUE	UNP P13487

- Molecule 4 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	G	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 5 is CESIUM ION (three-letter code: CS) (formula: Cs).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	4	Total Cs 4 4	0	0
5	B	4	Total Cs 4 4	0	0

- Molecule 6 is (1R)-2-{[(S)-{[(2S)-2,3-DIHYDROXYPROPYL]OXY}(HYDROXY)PHOSPHORYL]OXY}-1-[(HEXADECANOYLOXY)METHYL]ETHYL (9Z)-OCTADEC-9-ENOIC ACID

TE (three-letter code: PGW) (formula: C₄₀H₇₇O₁₀P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C O 22 17 5	0	0
6	B	1	Total C 9 9	0	0
6	B	1	Total C 9 9	0	0
6	B	1	Total C 9 9	0	0
6	B	1	Total C 9 9	0	0
6	B	1	Total C 9 9	0	0
6	B	1	Total C 9 9	0	0
6	B	1	Total C 7 7	0	0
6	B	1	Total C 9 9	0	0
6	B	1	Total C 8 8	0	0
6	B	1	Total C O P 23 14 8 1	0	0
6	B	1	Total C 8 8	0	0
6	B	1	Total C O P 36 25 10 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C 7 7	0	0
6	B	1	Total C 8 8	0	0
6	B	1	Total C 8 8	0	0
6	H	1	Total C O 22 17 5	0	0

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
-

- Chain G:
-
- 2% 78% 19%
- MET
 L36
 Q37
 F38
 Y39
 R40
 L52
 G55
 T56
 W57
 F60
 E67
 M73
 I74
 L75
 I102
 K106
 F120
 R129
 L144
 V152
 F156
 R159
 E167
 M173
 T174
 N178
 A182
 H185
 G186
 R189
 V201
 A202
 L207
 C212
 E213
 Q214
 M215

- [illegible]



E1	F2	T3	M4	V5	S6	C7	T8	T9	S10	F11	E12	C13	W14	S15	V16	C17	Q18	F19	L20	H21	N22	T23	S24	R25	G26	F27	C28	N29	H30	K31	K32	C33	R34	C35	Y36	S37
----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	145.43 Å 145.43 Å 285.59 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.56 49.93 – 2.56	Depositor EDS
% Data completeness (in resolution range)	91.9 (50.00-2.56) 92.1 (49.93-2.56)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 2.54 Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.237 , 0.262 0.243 , 0.269	Depositor DCC
R_{free} test set	4652 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	42.3	Xtriage
Anisotropy	0.231	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 99051 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	11770	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CS, PGW, NAP, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2608	0.61	0/3524
1	G	0.39	0/2608	0.60	0/3524
2	B	0.37	0/3169	0.55	0/4292
2	H	0.33	0/3036	0.50	0/4114
3	Y	0.26	0/292	0.46	0/389
All	All	0.38	0/11713	0.56	0/15843

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2556	0	2582	44	0
1	G	2556	0	2582	46	0
2	B	3088	0	3034	118	0
2	H	2959	0	2956	161	0
3	Y	295	0	282	10	0
4	A	48	0	25	11	0
4	G	48	0	25	12	0
5	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	H	4	0	0	0	0
6	B	190	0	251	18	0
6	H	22	0	25	7	0
All	All	11770	0	11762	391	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 391 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:400:LEU:HB2	2:B:401:PRO:HD3	1.43	1.00
1:G:55:GLY:HA3	4:G:1001:NAP:O3D	1.68	0.93
2:H:213:GLN:HE21	2:H:215:SER:HB3	1.31	0.91
2:H:146:GLN:HB3	2:H:243:ALA:HA	1.51	0.90
1:A:55:GLY:HA3	4:A:1001:NAP:O3D	1.73	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/333 (97%)	308 (95%)	14 (4%)	2 (1%)	30	52
1	G	324/333 (97%)	313 (97%)	10 (3%)	1 (0%)	46	68
2	B	384/514 (75%)	344 (90%)	27 (7%)	13 (3%)	5	6
2	H	357/514 (70%)	280 (78%)	60 (17%)	17 (5%)	3	3
3	Y	35/37 (95%)	17 (49%)	17 (49%)	1 (3%)	6	8
All	All	1424/1731 (82%)	1262 (89%)	128 (9%)	34 (2%)	7	12

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	132	TYR
2	B	133	ILE
2	B	134	LYS
2	B	135	GLU
2	B	137	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/280 (98%)	267 (98%)	6 (2%)	60	82
1	G	273/280 (98%)	266 (97%)	7 (3%)	54	77
2	B	332/459 (72%)	325 (98%)	7 (2%)	61	83
2	H	324/459 (71%)	319 (98%)	5 (2%)	72	88
3	Y	35/35 (100%)	33 (94%)	2 (6%)	25	46
All	All	1237/1513 (82%)	1210 (98%)	27 (2%)	60	82

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	302	LYS
1	G	129	ARG
2	H	350	ARG
1	G	73	MET
1	A	326	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	207	GLN
2	B	253	ASN
2	H	284	ASN
2	B	213	GLN
2	B	256	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PCA	Y	1	3	7,8,9	0.57	0	9,10,12	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PCA	Y	1	3	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 27 ligands modelled in this entry, 8 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAP	A	1001	-	42,52,52	1.72	3 (7%)	54,80,80	2.57	14 (25%)
6	PGW	B	504	-	21,21,50	0.60	0	23,23,56	1.25	4 (17%)
6	PGW	B	505	-	8,8,50	0.34	0	7,7,56	0.52	0
6	PGW	B	506	-	8,8,50	0.34	0	7,7,56	0.50	0
6	PGW	B	507	-	8,8,50	0.34	0	7,7,56	0.54	0
6	PGW	B	508	-	8,8,50	0.34	0	7,7,56	0.53	0
6	PGW	B	509	-	8,8,50	0.34	0	7,7,56	0.52	0
6	PGW	B	510	-	8,8,50	0.34	0	7,7,56	0.52	0
6	PGW	B	511	-	6,6,50	0.35	0	5,5,56	0.45	0
6	PGW	B	512	-	8,8,50	0.34	0	7,7,56	0.51	0
6	PGW	B	513	-	7,7,50	0.35	0	6,6,56	0.50	0
6	PGW	B	514	-	22,22,50	0.82	0	25,27,56	1.29	5 (20%)
6	PGW	B	515	-	7,7,50	0.34	0	6,6,56	0.49	0
6	PGW	B	516	-	35,35,50	0.66	0	36,41,56	0.94	3 (8%)
6	PGW	B	517	-	6,6,50	0.35	0	5,5,56	0.42	0
6	PGW	B	518	-	7,7,50	0.34	0	6,6,56	0.49	0
6	PGW	B	519	-	7,7,50	0.35	0	6,6,56	0.49	0
4	NAP	G	1001	-	42,52,52	1.67	7 (16%)	54,80,80	2.60	16 (29%)
6	PGW	H	504	-	21,21,50	0.60	0	23,23,56	1.31	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAP	A	1001	-	-	0/27/67/67	0/5/5/5
6	PGW	B	504	-	-	0/23/23/55	0/0/0/0
6	PGW	B	505	-	-	0/6/6/55	0/0/0/0
6	PGW	B	506	-	-	0/6/6/55	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGW	B	507	-	-	0/6/6/55	0/0/0/0
6	PGW	B	508	-	-	0/6/6/55	0/0/0/0
6	PGW	B	509	-	-	0/6/6/55	0/0/0/0
6	PGW	B	510	-	-	0/6/6/55	0/0/0/0
6	PGW	B	511	-	-	0/4/4/55	0/0/0/0
6	PGW	B	512	-	-	0/6/6/55	0/0/0/0
6	PGW	B	513	-	-	0/5/5/55	0/0/0/0
6	PGW	B	514	-	-	0/24/24/55	0/0/0/0
6	PGW	B	515	-	-	0/5/5/55	0/0/0/0
6	PGW	B	516	-	-	0/40/40/55	0/0/0/0
6	PGW	B	517	-	-	0/4/4/55	0/0/0/0
6	PGW	B	518	-	-	0/5/5/55	0/0/0/0
6	PGW	B	519	-	-	0/5/5/55	0/0/0/0
4	NAP	G	1001	-	-	0/27/67/67	0/5/5/5
6	PGW	H	504	-	-	0/23/23/55	0/0/0/0

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	1001	NAP	O3D-C3D	-2.29	1.37	1.43
4	G	1001	NAP	C8A-N7A	-2.28	1.30	1.34
4	G	1001	NAP	P2B-O3X	2.06	1.62	1.54
4	G	1001	NAP	C4N-C3N	2.33	1.43	1.39
4	G	1001	NAP	C2N-C3N	2.78	1.43	1.39

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	1001	NAP	N3A-C2A-N1A	-9.23	121.83	128.89
4	A	1001	NAP	N3A-C2A-N1A	-8.88	122.10	128.89
4	G	1001	NAP	PN-O3-PA	-8.23	109.62	132.73
4	A	1001	NAP	PN-O3-PA	-8.00	110.26	132.73
4	A	1001	NAP	O3-PA-O5B	-6.89	84.66	102.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

10 monomers are involved in 48 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1001	NAP	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	504	PGW	3	0
6	B	510	PGW	3	0
6	B	512	PGW	2	0
6	B	513	PGW	1	0
6	B	514	PGW	1	0
6	B	515	PGW	3	0
6	B	516	PGW	8	0
4	G	1001	NAP	12	0
6	H	504	PGW	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/333 (97%)	0.28	8 (2%) 61 59	17, 33, 60, 83	0
1	G	326/333 (97%)	0.20	8 (2%) 61 59	21, 39, 72, 96	0
2	B	386/514 (75%)	0.96	68 (17%) 2 1	25, 66, 110, 125	0
2	H	363/514 (70%)	3.17	163 (44%) 0 0	38, 115, 185, 204	0
3	Y	36/37 (97%)	6.99	36 (100%) 0 0	102, 107, 122, 126	36 (100%)
All	All	1437/1731 (83%)	1.34	283 (19%) 1 1	17, 56, 176, 204	36 (2%)

The worst 5 of 283 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	272	LEU	22.1
2	H	218	PHE	19.6
2	H	280	LEU	17.6
2	H	248	ALA	17.4
2	H	214	GLN	16.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PCA	Y	1	8/9	0.71	0.42	-	125,126,126,126	8

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	PGW	B	504	22/51	0.48	0.68	15.87	85,102,113,114	0
6	PGW	B	518	8/51	0.44	0.86	12.43	102,105,107,107	0
6	PGW	B	509	9/51	0.81	0.52	8.23	82,86,89,89	0
6	PGW	H	504	22/51	0.42	0.84	7.35	142,147,149,149	0
6	PGW	B	515	8/51	0.53	0.53	5.97	81,87,91,91	0
6	PGW	B	513	8/51	0.65	0.46	4.62	74,78,79,79	0
6	PGW	B	510	9/51	0.79	0.44	4.07	106,107,109,109	0
6	PGW	B	517	7/51	0.56	0.48	3.69	63,66,68,69	0
4	NAP	G	1001	48/48	0.95	0.20	2.64	25,39,51,51	0
6	PGW	B	505	9/51	0.64	0.36	2.25	79,82,84,85	0
6	PGW	B	516	36/51	0.58	0.39	2.22	107,125,141,141	0
6	PGW	B	514	23/51	0.70	0.31	1.48	109,119,123,124	0
4	NAP	A	1001	48/48	0.96	0.18	0.41	23,33,43,47	0
6	PGW	B	512	9/51	0.22	1.17	-	126,128,128,128	0
6	PGW	B	508	9/51	0.56	0.47	-	100,102,102,102	0
6	PGW	B	519	8/51	0.64	0.39	-	97,99,100,100	0
5	CS	H	505	1/1	0.92	0.39	-	126,126,126,126	1
5	CS	H	501	1/1	0.97	0.22	-	85,85,85,85	1
6	PGW	B	511	7/51	0.85	0.20	-	70,73,73,73	0
5	CS	B	503	1/1	0.93	0.14	-	101,101,101,101	1
5	CS	B	501	1/1	0.95	0.19	-	45,45,45,45	1
5	CS	B	520	1/1	1.00	0.21	-	61,61,61,61	1
5	CS	B	502	1/1	0.88	0.12	-	43,43,43,43	1
6	PGW	B	507	9/51	0.62	0.58	-	94,96,98,98	0
6	PGW	B	506	9/51	0.62	0.41	-	87,90,91,91	0
5	CS	H	503	1/1	0.94	0.24	-	75,75,75,75	1
5	CS	H	502	1/1	0.98	0.22	-	74,74,74,74	1

6.5 Other polymers ⓘ

There are no such residues in this entry.