



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:20 PM GMT

PDB ID : 1JUS  
Title : Crystal structure of the multidrug binding transcriptional repressor QacR bound to rhodamine 6G  
Authors : Schumacher, M.A.; Miller, M.C.; Grkovic, S.; Brown, M.H.; Skurray, R.A.; Brennan, R.G.  
Deposited on : 2001-08-27  
Resolution : 2.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry. We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org). A user guide is available at <http://wwpdb.org/validation/2016/XrayValidationReportHelp> with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3170 (2.88-2.80)
Clashscore	102246	3658 (2.88-2.80)
Ramachandran outliers	100387	3591 (2.88-2.80)
Sidechain outliers	100360	3594 (2.88-2.80)
RSRZ outliers	91569	3184 (2.88-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	 2% 58% 32% 6% •
1	B	194	 2% 45% 41% 8% ••
1	D	194	 4% 53% 39% 5% •
1	E	194	 6% 57% 31% 7% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	599	-	-	X	-
2	SO4	A	699	-	-	-	X
2	SO4	A	799	-	-	-	X
2	SO4	A	800	-	-	X	-
2	SO4	A	802	-	-	-	X
2	SO4	B	199	-	-	-	X
2	SO4	B	600	-	-	-	X
2	SO4	D	1000	-	-	-	X
2	SO4	D	899	-	-	-	X
2	SO4	D	900	-	-	X	-
2	SO4	D	901	-	-	-	X
2	SO4	E	399	-	-	-	X
2	SO4	E	901	-	-	-	X
3	RHQ	A	201	-	X	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6398 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HYPOTHETICAL TRANSCRIPTIONAL REGULATOR IN QACA 5'REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	B	186	1547	998	252	295	2	0	0	0
1	D	186	1547	998	252	295	2	0	0	0
1	A	186	1547	998	252	295	2	0	0	0
1	E	186	1547	998	252	295	2	0	0	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	72	ALA	CYS	ENGINEERED	UNP P0A0N4
B	141	SER	CYS	ENGINEERED	UNP P0A0N4
B	189	HIS	-	EXPRESSION TAG	UNP P0A0N4
B	190	HIS	-	EXPRESSION TAG	UNP P0A0N4
B	191	HIS	-	EXPRESSION TAG	UNP P0A0N4
B	192	HIS	-	EXPRESSION TAG	UNP P0A0N4
B	193	HIS	-	EXPRESSION TAG	UNP P0A0N4
B	194	HIS	-	EXPRESSION TAG	UNP P0A0N4
D	72	ALA	CYS	ENGINEERED	UNP P0A0N4
D	141	SER	CYS	ENGINEERED	UNP P0A0N4
D	189	HIS	-	EXPRESSION TAG	UNP P0A0N4
D	190	HIS	-	EXPRESSION TAG	UNP P0A0N4
D	191	HIS	-	EXPRESSION TAG	UNP P0A0N4
D	192	HIS	-	EXPRESSION TAG	UNP P0A0N4
D	193	HIS	-	EXPRESSION TAG	UNP P0A0N4
D	194	HIS	-	EXPRESSION TAG	UNP P0A0N4
A	72	ALA	CYS	ENGINEERED	UNP P0A0N4
A	141	SER	CYS	ENGINEERED	UNP P0A0N4
A	189	HIS	-	EXPRESSION TAG	UNP P0A0N4
A	190	HIS	-	EXPRESSION TAG	UNP P0A0N4

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	191	HIS	-	EXPRESSION TAG	UNP P0A0N4
A	192	HIS	-	EXPRESSION TAG	UNP P0A0N4
A	193	HIS	-	EXPRESSION TAG	UNP P0A0N4
A	194	HIS	-	EXPRESSION TAG	UNP P0A0N4
E	72	ALA	CYS	ENGINEERED	UNP P0A0N4
E	141	SER	CYS	ENGINEERED	UNP P0A0N4
E	189	HIS	-	EXPRESSION TAG	UNP P0A0N4
E	190	HIS	-	EXPRESSION TAG	UNP P0A0N4
E	191	HIS	-	EXPRESSION TAG	UNP P0A0N4
E	192	HIS	-	EXPRESSION TAG	UNP P0A0N4
E	193	HIS	-	EXPRESSION TAG	UNP P0A0N4
E	194	HIS	-	EXPRESSION TAG	UNP P0A0N4

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is RHODAMINE 6G (three-letter code: RHQ) (formula:  $C_{28}H_{31}N_2O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	33	28	2	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	25	Total	O	0	0
			25	25		
4	B	11	Total	O	0	0
			11	11		
4	D	10	Total	O	0	0
			10	10		
4	E	16	Total	O	0	0
			16	16		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HYPOTHETICAL TRANSCRIPTIONAL REGULATOR IN QACA 5'REGION



- Molecule 1: HYPOTHETICAL TRANSCRIPTIONAL REGULATOR IN QACA 5'REGION



- Molecule 1: HYPOTHETICAL TRANSCRIPTIONAL REGULATOR IN QACA 5'REGION



HIS  
HIS  
HIS  
HIS

- Molecule 1: HYPOTHETICAL TRANSCRIPTIONAL REGULATOR IN QACA 5'REGION

Chain E:  6% 57% 31% 7%

MSE  
R2  
L3  
T25  
V29  
K30  
S34  
S35  
K36  
G37  
N38  
L39  
Y40  
Y41  
H42  
F43  
K44  
T45  
K46  
E47  
N48  
L49  
F50  
L51  
L54  
N55  
E58  
Q62  
E63  
Q64  
K65  
K66  
R67  
E68  
A72  
K73  
T74  
M75  
R76  
E77  
L85  
Y92  
Q96  
N97  
I100  
E101  
F102  
Y103  
T104

E105  
Y106  
K107  
K108  
M109  
S110  
S111  
I112  
M113  
E114  
K115  
M116  
M117  
K118  
L119  
E120  
M121  
K122  
Y123  
I124  
K132  
E133  
G134  
M135  
L136  
M137  
M140  
S141  
I142  
M143  
D144  
Y145  
M146  
A147  
V148  
S149  
K150  
I151  
M157  
T161  
F162  
T163  
H164  
E165  
I168  
M169  
E170  
R171  
L183  
M184  
G185  
L186  
S187  
LYS

HIS  
HIS  
HIS  
HIS  
HIS  
HIS

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	172.30Å 172.30Å 95.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	77.06 – 2.84 77.05 – 2.84	Depositor EDS
% Data completeness (in resolution range)	99.7 (77.06-2.84) 99.8 (77.05-2.84)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.65 (at 2.82Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.230 , 0.272 0.226 , 0.268	Depositor DCC
$R_{free}$ test set	3421 reflections (9.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	70.8	Xtrriage
Anisotropy	0.011	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 67.3	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	1 of 34321 reflections (0.003%)	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6398	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RHQ, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1576	0.60	0/2120
1	B	0.45	0/1576	0.69	0/2120
1	D	0.41	0/1576	0.62	0/2120
1	E	0.43	0/1576	0.61	0/2120
All	All	0.44	0/6304	0.63	0/8480

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	107	TYR	Sidechain
1	A	20	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1547	0	1536	80	0
1	B	1547	0	1536	107	0
1	D	1547	0	1536	81	0
1	E	1547	0	1536	93	0
2	A	35	0	0	6	0
2	B	30	0	0	1	0
2	D	25	0	0	4	0
2	E	25	0	0	0	0
3	A	33	0	31	10	0
4	A	25	0	0	0	0
4	B	11	0	0	0	0
4	D	10	0	0	0	0
4	E	16	0	0	1	1
All	All	6398	0	6175	339	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 339 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:97:ASN:HB2	1:E:100:ILE:HD11	1.23	1.16
1:D:165:GLU:HB3	1:E:103:TYR:HE1	1.06	1.07
1:D:165:GLU:HB3	1:E:103:TYR:CE1	1.91	1.06
1:E:168:ILE:H	1:E:168:ILE:HD12	1.24	0.98
1:E:44:LYS:HG3	1:E:45:THR:HG23	1.45	0.98

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:907:HOH:O	4:E:907:HOH:O[7_556]	1.50	0.70

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/194 (95%)	174 (95%)	9 (5%)	1 (0%)	34	67
1	B	184/194 (95%)	156 (85%)	22 (12%)	6 (3%)	5	15
1	D	184/194 (95%)	163 (89%)	17 (9%)	4 (2%)	8	26
1	E	184/194 (95%)	163 (89%)	16 (9%)	5 (3%)	6	20
All	All	736/776 (95%)	656 (89%)	64 (9%)	16 (2%)	8	26

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	44	LYS
1	D	186	LEU
1	E	110	ASN
1	B	41	TYR
1	B	89	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/174 (97%)	154 (91%)	15 (9%)	12	32
1	B	169/174 (97%)	148 (88%)	21 (12%)	6	16
1	D	169/174 (97%)	155 (92%)	14 (8%)	14	36
1	E	169/174 (97%)	155 (92%)	14 (8%)	14	36
All	All	676/696 (97%)	612 (90%)	64 (10%)	11	28

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	71	LYS
1	A	14	LEU
1	E	107	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	73	LYS
1	D	164	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	184	ASN
1	A	135	ASN
1	E	143	ASN
1	A	38	ASN
1	A	143	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	RHQ	A	201	-	33,36,36	7.15	26 (78%)	40,51,51	4.72	23 (57%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	299	-	4,4,4	0.24	0	6,6,6	0.07	0
2	SO4	A	599	-	4,4,4	0.10	0	6,6,6	0.08	0
2	SO4	A	699	-	4,4,4	0.22	0	6,6,6	0.12	0
2	SO4	A	799	-	4,4,4	0.17	0	6,6,6	0.09	0
2	SO4	A	800	-	4,4,4	0.31	0	6,6,6	0.07	0
2	SO4	A	801	-	4,4,4	0.27	0	6,6,6	0.07	0
2	SO4	A	802	-	4,4,4	0.27	0	6,6,6	0.09	0
2	SO4	B	1000	-	4,4,4	0.17	0	6,6,6	0.08	0
2	SO4	B	199	-	4,4,4	0.14	0	6,6,6	0.16	0
2	SO4	B	499	-	4,4,4	0.18	0	6,6,6	0.10	0
2	SO4	B	600	-	4,4,4	0.20	0	6,6,6	0.10	0
2	SO4	B	601	-	4,4,4	0.27	0	6,6,6	0.12	0
2	SO4	B	700	-	4,4,4	0.21	0	6,6,6	0.09	0
2	SO4	D	1000	-	4,4,4	0.19	0	6,6,6	0.12	0
2	SO4	D	899	-	4,4,4	0.36	0	6,6,6	0.11	0
2	SO4	D	900	-	4,4,4	0.24	0	6,6,6	0.09	0
2	SO4	D	901	-	4,4,4	0.27	0	6,6,6	0.12	0
2	SO4	D	999	-	4,4,4	0.23	0	6,6,6	0.17	0
2	SO4	E	399	-	4,4,4	0.19	0	6,6,6	0.13	0
2	SO4	E	900	-	4,4,4	0.23	0	6,6,6	0.10	0
2	SO4	E	901	-	4,4,4	0.19	0	6,6,6	0.16	0
2	SO4	E	902	-	4,4,4	0.21	0	6,6,6	0.09	0
2	SO4	E	903	-	4,4,4	0.18	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RHQ	A	201	-	-	0/15/21/21	0/4/4/4
2	SO4	A	299	-	-	0/0/0/0	0/0/0/0
2	SO4	A	599	-	-	0/0/0/0	0/0/0/0
2	SO4	A	699	-	-	0/0/0/0	0/0/0/0
2	SO4	A	799	-	-	0/0/0/0	0/0/0/0
2	SO4	A	800	-	-	0/0/0/0	0/0/0/0
2	SO4	A	801	-	-	0/0/0/0	0/0/0/0
2	SO4	A	802	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1000	-	-	0/0/0/0	0/0/0/0
2	SO4	B	199	-	-	0/0/0/0	0/0/0/0
2	SO4	B	499	-	-	0/0/0/0	0/0/0/0
2	SO4	B	600	-	-	0/0/0/0	0/0/0/0
2	SO4	B	601	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	B	700	-	-	0/0/0/0	0/0/0/0
2	SO4	D	1000	-	-	0/0/0/0	0/0/0/0
2	SO4	D	899	-	-	0/0/0/0	0/0/0/0
2	SO4	D	900	-	-	0/0/0/0	0/0/0/0
2	SO4	D	901	-	-	0/0/0/0	0/0/0/0
2	SO4	D	999	-	-	0/0/0/0	0/0/0/0
2	SO4	E	399	-	-	0/0/0/0	0/0/0/0
2	SO4	E	900	-	-	0/0/0/0	0/0/0/0
2	SO4	E	901	-	-	0/0/0/0	0/0/0/0
2	SO4	E	902	-	-	0/0/0/0	0/0/0/0
2	SO4	E	903	-	-	0/0/0/0	0/0/0/0

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	201	RHQ	C19-C26	-3.37	1.42	1.50
3	A	201	RHQ	C10-C7	2.29	1.42	1.37
3	A	201	RHQ	C13-C12	2.68	1.45	1.37
3	A	201	RHQ	C22-N1	3.17	1.56	1.46
3	A	201	RHQ	C20-C12	3.39	1.57	1.51

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	201	RHQ	C2-C9-C8	-17.78	106.39	119.57
3	A	201	RHQ	C1-O1-C7	-7.17	113.11	122.37
3	A	201	RHQ	C15-C14-C19	-6.59	110.49	118.03
3	A	201	RHQ	C3-C2-C9	-6.35	116.97	124.27
3	A	201	RHQ	C10-C11-N1	-5.22	112.26	121.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	201	RHQ	10	0
2	A	599	SO4	2	0
2	A	699	SO4	1	0
2	A	800	SO4	3	0
2	B	600	SO4	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	899	SO4	1	0
2	D	900	SO4	2	0
2	D	999	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	184/194 (94%)	0.09	3 (1%) 74 67	34, 53, 89, 118	0
1	B	184/194 (94%)	0.27	3 (1%) 74 67	36, 70, 121, 145	0
1	D	184/194 (94%)	0.44	7 (3%) 44 33	42, 70, 105, 119	0
1	E	184/194 (94%)	0.44	12 (6%) 22 14	39, 60, 97, 111	0
All	All	736/776 (94%)	0.31	25 (3%) 49 38	34, 62, 108, 145	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	187	SER	4.7
1	E	187	SER	3.5
1	D	186	LEU	3.3
1	A	187	SER	2.9
1	D	131	PHE	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	802	5/5	0.67	0.53	13.44	145,145,146,146	0
2	SO4	E	399	5/5	0.76	0.63	12.91	175,175,176,176	0
2	SO4	A	699	5/5	0.77	0.37	6.53	142,143,143,144	0
3	RHQ	A	201	33/33	0.82	0.47	5.60	30,38,44,48	0
2	SO4	D	901	5/5	0.87	0.40	4.78	131,131,132,132	0
2	SO4	B	199	5/5	0.89	0.33	4.60	123,124,124,124	0
2	SO4	A	799	5/5	0.98	0.31	4.31	83,83,85,85	0
2	SO4	B	600	5/5	0.80	0.37	3.65	143,143,143,144	0
2	SO4	E	901	5/5	0.73	0.29	3.32	137,137,137,138	0
2	SO4	D	1000	5/5	0.58	0.48	3.32	144,144,145,145	0
2	SO4	D	899	5/5	0.75	0.37	2.68	118,119,121,122	0
2	SO4	B	700	5/5	0.81	0.21	-0.03	174,174,174,174	0
2	SO4	D	900	5/5	0.92	0.18	-0.60	127,127,128,128	0
2	SO4	E	903	5/5	0.80	0.32	-	174,174,174,174	0
2	SO4	B	601	5/5	0.83	0.23	-	133,133,133,134	0
2	SO4	E	900	5/5	0.89	0.18	-	116,116,116,116	0
2	SO4	B	499	5/5	0.79	0.22	-	137,137,138,138	0
2	SO4	E	902	5/5	0.83	0.21	-	138,138,138,138	0
2	SO4	A	800	5/5	0.91	0.26	-	98,98,99,100	0
2	SO4	A	299	5/5	0.78	0.24	-	130,130,130,130	0
2	SO4	A	599	5/5	0.89	0.28	-	121,122,123,124	0
2	SO4	D	999	5/5	0.91	0.19	-	122,122,123,123	0
2	SO4	A	801	5/5	0.83	0.19	-	129,129,129,129	0
2	SO4	B	1000	5/5	0.72	0.37	-	199,199,199,199	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.