



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 05:53 PM GMT

PDB ID : 4JU1
Title : Crystal structure of hcv ns5b polymerase in complex with compound 6
Authors : Coulombe, R.
Deposited on : 2013-03-24
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	576	73%	22%	..
1	B	576	%	70%	26% ..


The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	3001	-	-	-	X
2	GOL	B	601	-	-	-	X
4	MG	A	3004	-	-	-	X
4	MG	B	603	-	-	-	X



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is 6-[3-([1,3]OXAZOLO[4,5-B]PYRIDIN-2-YL)-2-(TRIFLUOROMETHYL)PHENOXY]-1-(2,4,6-TRIFLUOROBENZYL)QUINAZOLIN-4(1H)-ONE (three-letter code: 1NZ) (formula: C₂₈H₁₄F₆N₄O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C F N O 41 28 6 4 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C F N O 41 28 6 4 3	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total Mg 2 2	0	0
4	A	2	Total Mg 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	178	Total O 178 178	0	0
5	B	146	Total O 146 146	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Genome polyprotein



- Molecule 1: Genome polyprotein



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.82Å 107.76Å 135.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.90 39.26 – 2.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-2.90) 94.1 (39.26-2.90)	Depositor EDS
R _{merge}	0.09	Depositor
R _{sym}	(Not available)	Depositor
<I/σ(I)> ¹	2.75 (at 2.90Å)	Xtriage
Refinement program	CNX 2002	Depositor
R, R _{free}	0.195 , 0.252 0.192 , 0.245	Depositor DCC
R _{free} test set	3334 reflections (11.09%)	DCC
Wilson B-factor (Å ²)	54.3	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent k _{sol} (e/Å ³), B _{sol} (Å ²)	0.30 , 45.9	EDS
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
L-test for twinning ²	< L > = 0.48, <L ² > = 0.31	Xtriage
Outliers	0 of 33463 reflections	Xtriage
F _o ,F _c correlation	0.94	EDS
Total number of atoms	9146	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of <|L|>, <L²> for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	147	VAL
1	B	541	ALA
1	A	96	SER
1	B	400	ALA
1	B	540	PRO
1	A	310	ASP
1	A	347	SER
1	B	404	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	477/491 (97%)	466 (98%)	11 (2%)	58 87
1	B	477/491 (97%)	466 (98%)	11 (2%)	58 87
All	All	954/982 (97%)	932 (98%)	22 (2%)	58 87

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LEU
1	A	57	LEU
1	A	83	LEU
1	A	220	ASP
1	A	273	ASN
1	A	313	MET
1	A	336	LEU
1	A	402	HIS
1	A	434	LEU
1	A	461	GLN
1	A	547	LEU
1	B	34	HIS
1	B	106	LYS
1	B	198	LYS
1	B	248	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	273	ASN
1	B	311	CYS
1	B	331	GLU
1	B	366	CYS
1	B	465	ARG
1	B	469	LEU
1	B	492	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	251	GLN
1	A	273	ASN
1	A	438	GLN
1	A	514	GLN
1	A	544	GLN
1	B	24	ASN
1	B	34	HIS
1	B	110	ASN
1	B	148	GLN
1	B	206	ASN
1	B	251	GLN
1	B	273	ASN
1	B	406	ASN
1	B	467	HIS
1	B	514	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	1NZ	F32-C5-C6	2.47	120.49	117.45
3	A	3002	1NZ	C7-N10-C11	2.49	123.49	120.38
3	A	3002	1NZ	C1-C2-C3	2.54	119.28	116.63
3	B	602	1NZ	F31-C1-C6	2.55	120.58	117.45
3	B	602	1NZ	C13-C12-C11	2.58	122.57	118.49
3	B	602	1NZ	C1-C2-C3	2.65	119.39	116.63
3	B	602	1NZ	C5-C4-C3	2.69	119.44	116.63
3	B	602	1NZ	C26-C25-C33	2.82	126.05	120.09
3	A	3002	1NZ	C25-C33-N35	3.09	128.04	123.68
3	A	3002	1NZ	C26-C25-C33	3.17	126.80	120.09
3	B	602	1NZ	C7-N10-C11	3.22	124.41	120.38
3	A	3002	1NZ	C16-C11-N10	3.45	125.44	121.47
3	B	602	1NZ	C25-C33-N35	3.50	128.60	123.68
3	B	602	1NZ	C16-C11-N10	3.94	126.00	121.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3001	GOL	1	0
3	B	602	1NZ	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	560/576 (97%)	-0.45	0 (100%) 100 (100%)	27, 45, 68, 90	0
1	B	560/576 (97%)	-0.21	4 (0%) 89 88	31, 56, 92, 105	0
All	All	1120/1152 (97%)	-0.33	4 (0%) 93 92	27, 49, 84, 105	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	507	VAL	2.7
1	B	23	ILE	2.5
1	B	24	ASN	2.5
1	B	511	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	MG	B	603	1/1	0.97	0.27	7.96	57,57,57,57	0
4	MG	A	3004	1/1	0.72	0.43	6.24	64,64,64,64	0
2	GOL	B	601	6/6	0.93	0.26	2.82	76,77,77,77	0
2	GOL	A	3001	6/6	0.92	0.23	2.77	61,62,63,63	0
3	1NZ	A	3002	41/41	0.95	0.18	1.35	45,51,59,60	0
3	1NZ	B	602	41/41	0.95	0.18	0.33	56,61,71,71	0
4	MG	A	3003	1/1	0.90	0.08	-5.87	36,36,36,36	0
4	MG	B	604	1/1	0.78	0.22	-	77,77,77,77	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.