



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:21 PM GMT

PDB ID : 1JXP
Title : BK STRAIN HEPATITIS C VIRUS (HCV) NS3-NS4A
Authors : Yan, Y.; Munshi, S.; Chen, Z.
Deposited on : 1997-08-21
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.





Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	186	
1	B	186	
2	C	16	
2	D	16	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2987 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NS3 SERINE PROTEASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	177	Total	C	N	O	S	0	0	0
			1301	809	234	248	10			
1	B	177	Total	C	N	O	S	0	0	0
			1301	809	234	248	10			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	SER	ALA	CONFLICT	UNP P26663
A	18	ILE	VAL	CONFLICT	UNP P26663
A	56	TYR	PHE	CONFLICT	UNP P26663
A	86	GLN	HIS	CONFLICT	UNP P26663
A	122	SER	GLY	CONFLICT	UNP P26663
A	170	VAL	ILE	CONFLICT	UNP P26663
B	7	SER	ALA	CONFLICT	UNP P26663
B	18	ILE	VAL	CONFLICT	UNP P26663
B	56	TYR	PHE	CONFLICT	UNP P26663
B	86	GLN	HIS	CONFLICT	UNP P26663
B	122	SER	GLY	CONFLICT	UNP P26663
B	170	VAL	ILE	CONFLICT	UNP P26663

- Molecule 2 is a protein called NS4A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	12	Total	C	N	O	0	0	0
			84	55	15	14			
2	D	12	Total	C	N	O	0	0	0
			84	55	15	14			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Zn 1	0	0
3	A	1	Total 1	Zn 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	103	Total 103	O 103	0	0
4	B	101	Total 101	O 101	0	0
4	C	8	Total 8	O 8	0	0
4	D	3	Total 3	O 3	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: NS3 SERINE PROTEASE



• Molecule 1: NS3 SERINE PROTEASE



• Molecule 2: NS4A



• Molecule 2: NS4A



LYS	G221	S222	Y223	Y224	I225	I229	I230	L231	S232	GLY	ARG	LYS
-----	------	------	------	------	------	------	------	------	------	-----	-----	-----

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	96.96 Å 96.96 Å 167.10 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.20)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.245 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2987	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	2/1326 (0.2%)	0.90	1/1806 (0.1%)
1	B	0.64	1/1326 (0.1%)	0.89	1/1806 (0.1%)
2	C	0.60	0/83	0.94	0/111
2	D	0.65	0/83	0.89	0/111
All	All	0.67	3/2818 (0.1%)	0.89	2/3834 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	16	CYS	CB-SG	8.53	1.96	1.82
1	A	97	CYS	CB-SG	7.57	1.95	1.82
1	A	16	CYS	CB-SG	6.02	1.92	1.82

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	97	CYS	N-CA-C	5.67	126.31	111.00
1	A	143	LEU	N-CA-C	-5.47	96.22	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1301	0	1306	67	0
1	B	1301	0	1305	60	0
2	C	84	0	99	7	0
2	D	84	0	99	8	0
3	A	1	0	0	1	0
3	B	1	0	0	0	0
4	A	103	0	0	2	0
4	B	101	0	0	1	0
4	C	8	0	0	1	0
4	D	3	0	0	0	0
All	All	2987	0	2809	128	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 128 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:890:HOH:O	1:B:16:CYS:SG	2.15	1.03
1:A:16:CYS:SG	4:A:727:HOH:O	2.19	0.91
1:A:145:CYS:HG	3:A:190:ZN:ZN	0.85	0.87
1:B:33:VAL:HG13	2:D:229:ILE:HB	1.61	0.82
1:A:103:ASP:O	1:A:146:PRO:HD3	1.79	0.82

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	175/186 (94%)	149 (85%)	23 (13%)	3 (2%)	11 7
1	B	175/186 (94%)	155 (89%)	14 (8%)	6 (3%)	5 2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	10/16 (62%)	9 (90%)	1 (10%)	0	100	100
2	D	10/16 (62%)	10 (100%)	0	0	100	100
All	All	370/404 (92%)	323 (87%)	38 (10%)	9 (2%)	7	4

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	91	ALA
1	B	99	CYS
1	A	129	PRO
1	A	146	PRO
1	B	97	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/151 (95%)	126 (88%)	18 (12%)	6	4
1	B	144/151 (95%)	124 (86%)	20 (14%)	4	3
2	C	10/13 (77%)	10 (100%)	0	100	100
2	D	10/13 (77%)	8 (80%)	2 (20%)	1	1
All	All	308/328 (94%)	268 (87%)	40 (13%)	5	4

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	4	THR
1	B	43	PHE
1	B	161	ARG
1	B	16	CYS
1	B	51	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	B	80	GLN
1	A	86	GLN
1	A	27	ASN
1	B	41	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.