



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:56 PM GMT

PDB ID : 4K03
Title : Crystal structure of Drosophila Cryochrome
Authors : Berndt, A.; Wolf, E.
Deposited on : 2013-04-03
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	561	 36% 46% 12% • 6%
1	B	561	 37% 48% 11% ••

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8746 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cryptochrome-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	529	4238	2707	749	758	24	0	0	0
1	B	543	4350	2775	769	781	25	0	0	0

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	EXPRESSION TAG	UNP O77059
A	-17	ALA	-	EXPRESSION TAG	UNP O77059
A	-16	MET	-	EXPRESSION TAG	UNP O77059
A	-15	GLY	-	EXPRESSION TAG	UNP O77059
A	-14	SER	-	EXPRESSION TAG	UNP O77059
A	-13	GLY	-	EXPRESSION TAG	UNP O77059
A	-12	ILE	-	EXPRESSION TAG	UNP O77059
A	-11	GLN	-	EXPRESSION TAG	UNP O77059
A	-10	ARG	-	EXPRESSION TAG	UNP O77059
A	-9	PRO	-	EXPRESSION TAG	UNP O77059
A	-8	THR	-	EXPRESSION TAG	UNP O77059
A	-7	SER	-	EXPRESSION TAG	UNP O77059
A	-6	THR	-	EXPRESSION TAG	UNP O77059
A	-5	SER	-	EXPRESSION TAG	UNP O77059
A	-4	SER	-	EXPRESSION TAG	UNP O77059
A	-3	LEU	-	EXPRESSION TAG	UNP O77059
A	-2	VAL	-	EXPRESSION TAG	UNP O77059
A	-1	ALA	-	EXPRESSION TAG	UNP O77059
A	0	ALA	-	EXPRESSION TAG	UNP O77059
B	-18	GLY	-	EXPRESSION TAG	UNP O77059
B	-17	ALA	-	EXPRESSION TAG	UNP O77059
B	-16	MET	-	EXPRESSION TAG	UNP O77059
B	-15	GLY	-	EXPRESSION TAG	UNP O77059
B	-14	SER	-	EXPRESSION TAG	UNP O77059
B	-13	GLY	-	EXPRESSION TAG	UNP O77059

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	ILE	-	EXPRESSION TAG	UNP O77059
B	-11	GLN	-	EXPRESSION TAG	UNP O77059
B	-10	ARG	-	EXPRESSION TAG	UNP O77059
B	-9	PRO	-	EXPRESSION TAG	UNP O77059
B	-8	THR	-	EXPRESSION TAG	UNP O77059
B	-7	SER	-	EXPRESSION TAG	UNP O77059
B	-6	THR	-	EXPRESSION TAG	UNP O77059
B	-5	SER	-	EXPRESSION TAG	UNP O77059
B	-4	SER	-	EXPRESSION TAG	UNP O77059
B	-3	LEU	-	EXPRESSION TAG	UNP O77059
B	-2	VAL	-	EXPRESSION TAG	UNP O77059
B	-1	ALA	-	EXPRESSION TAG	UNP O77059
B	0	ALA	-	EXPRESSION TAG	UNP O77059

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total 27	O 27	0	0
3	B	25	Total 25	O 25	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cryptochrome-1



• Molecule 1: Cryptochrome-1



C485	L486	L487	G488	V489	H490	L491	P492	E493	R494	L495	L496	D497	L498	S499	M500	K503	R504	N505	M506	L507	A508	S511	L512	R513	N514	S515	L516	L517	T518	P519	P520	H521	C522	R523	R524	P525	S526	N527	E528	E529	E530	V531	R532	G533	F534	F535	N536	L537	ALA	ASP	VAL	VAL	VAL			
V415	C416	A417	G418	N419	V420	M421	A427	F428	R429	R430	L431	L432	D433	S434	S435	L436	V437	T438	C439	P440	V441	A442	L443	A444	R445	R446	L447	D448	P449	D450	G451	Y453	T452	I454	K455	Q456	T457	V458	P459	E460	L461	V464	P465	K466	E467	F468	E471	P472	W473	R474	N475	S476	A477	E478	Q479	Q480
L349	L350	Q351	S352	W353	R354	L355	G356	Q357	T358	G359	F360	P361	L362	I363	D364	G365	A366	M367	R368	L371	L376	R377	R378	T379	L380	T383	V384	A385	T386	F387	L388	I454	T389	R390	G391	G392	L393	W394	E460	Q395	S396	N397	E398	L401	F402	L405	K406	Y407	L408	L409	D410	A411	D412	W413	S414	
S276	V277	R278	R279	F280	Y281	H282	S283	D286	N287	F288	K289	H290	V291	Q292	L293	C296	V297	R298	G299	V300	Q301	N302	T303	H307	I308	T309	G310	G311	L312	I313	Y319	T320	R321	S322	W323	N324	W325	P326	N327	V328	D329	R330	N331	N334	D335	L336	C337	L338	K342	A343	R344	N346				
P203	T204	P205	E206	Y211	G212	D213	G216	N219	L218	A219	K220	I221	I222	N223	R224	T228	Q229	A230	G231	L232	L233	L234	D235	E236	R237	L238	K239	V240	E241	H242	A244	F245	E246	R247	G248	F249	Y250	L251	P252	L256	P257	N258	I259	H260	D261	S262	P263	L270	R271	F272	G273	C274	L275			
S120	I121	R122	S123	L124	C125	R126	E127	L128	I130	E134	K135	V136	S137	H138	T149	M150	G151	G152	I153	P154	P155	L156	L157	Y158	Q159	L162	H163	T164	V165	Q166	L170	P171	P172	T175	A176	D177	A178	A179	L180	E181	T184	F185	V186	E191	F192	L196	F199	L202								

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.12Å 121.81Å 79.72Å 90.00° 114.78° 90.00°	Depositor
Resolution (Å)	40.49 – 3.20 46.60 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.49-3.20) 99.7 (46.60-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 3.19Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.188 , 0.269 0.188 , 0.269	Depositor DCC
R_{free} test set	1041 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	58.7	Xtrriage
Anisotropy	0.178	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 85.4	EDS
Estimated twinning fraction	0.035 for h,-k,-h-l	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Outliers	0 of 20371 reflections	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8746	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/4355	0.71	3/5929 (0.1%)
1	B	0.50	0/4469	0.72	1/6082 (0.0%)
All	All	0.51	0/8824	0.72	4/12011 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	251	LEU	CA-CB-CG	8.68	135.26	115.30
1	A	393	LEU	N-CA-C	6.16	127.62	111.00
1	A	291	VAL	N-CA-C	-5.91	95.03	111.00
1	B	232	LEU	CA-CB-CG	5.01	126.83	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4238	0	4076	343	0
1	B	4350	0	4190	342	0
2	A	53	0	31	3	0
2	B	53	0	31	1	0
3	A	27	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	25	0	0	1	0
All	All	8746	0	8328	670	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 670 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:VAL:HG13	1:A:298:ARG:H	0.97	1.09
1:A:297:VAL:HG13	1:A:298:ARG:N	1.79	0.98
1:A:192:PHE:CZ	1:A:196:LEU:HD22	2.04	0.93
1:A:297:VAL:CG1	1:A:298:ARG:H	1.81	0.92
1:A:17:LEU:HD12	1:A:70:GLN:OE1	1.70	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/561 (94%)	409 (78%)	90 (17%)	26 (5%)	3	21
1	B	541/561 (96%)	444 (82%)	70 (13%)	27 (5%)	3	21
All	All	1066/1122 (95%)	853 (80%)	160 (15%)	53 (5%)	3	21

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	221	ILE
1	A	252	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	254	GLN
1	A	291	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	447/488 (92%)	377 (84%)	70 (16%)	3 15
1	B	458/488 (94%)	384 (84%)	74 (16%)	3 14
All	All	905/976 (93%)	761 (84%)	144 (16%)	3 14

5 of 144 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	512	LEU
1	B	83	VAL
1	B	460	GLU
1	A	523	CYS
1	B	46	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	505	ASN
1	A	533	GLN
1	B	301	GLN
1	A	403	HIS
1	B	96	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	601	-	48,58,58	1.08	5 (10%)	54,89,89	2.63	10 (18%)
2	FAD	B	601	-	48,58,58	1.06	5 (10%)	54,89,89	2.62	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	0/30/50/50	0/6/6/6
2	FAD	B	601	-	-	0/30/50/50	0/6/6/6

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C4X-N5	-2.42	1.29	1.33
2	A	601	FAD	C10-N10	-2.24	1.36	1.39
2	A	601	FAD	C10-N1	-2.17	1.32	1.35
2	B	601	FAD	C10-N10	-2.11	1.36	1.39
2	B	601	FAD	C10-N1	-2.02	1.32	1.35

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	FAD	C4X-C4-N3	-6.85	114.22	123.59
2	A	601	FAD	C4X-C4-N3	-6.57	114.60	123.59
2	B	601	FAD	C4X-C10-N10	-6.36	116.77	120.52
2	A	601	FAD	C4X-C10-N10	-6.25	116.84	120.52
2	B	601	FAD	C4-C4X-C10	-3.88	117.46	119.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	FAD	3	0
2	B	601	FAD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	529/561 (94%)	-0.29	3 (0%) 90 84	23, 52, 87, 135	0
1	B	543/561 (96%)	-0.19	6 (1%) 82 72	17, 54, 92, 161	0
All	All	1072/1122 (95%)	-0.24	9 (0%) 87 80	17, 54, 90, 161	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	485	CYS	3.0
1	B	0	ALA	2.9
1	B	490	HIS	2.7
1	B	461	LEU	2.7
1	B	473	TRP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FAD	A	601	53/53	0.96	0.19	0.09	37,37,60,60	0
2	FAD	B	601	53/53	0.96	0.17	-0.28	39,39,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.