



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:52 AM GMT

PDB ID : 3K2L
Title : Crystal Structure of dual-specificity tyrosine phosphorylation regulated kinase 2 (DYRK2)
Authors : Filippakopoulos, P.; Myrianthopoulos, V.; Soundararajan, M.; Krojer, T.; Hapka, E.; Fedorov, O.; Berridge, G.; Wang, J.; Shrestha, L.; Pike, A.C.W.; Ugochukwu, E.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.; Weigelt, J.; Bountra, C.; Mikros, E.; Knapp, S.; Structural Genomics Consortium (SGC)
Deposited on : 2009-09-30
Resolution : 2.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity tyrosine-phosphorylation-regulated kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	P	S	0	2	0
			3157	2004	546	579	5	23			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	MET	-	INITIATING METHIONINE	UNP Q92630
A	51	HIS	-	EXPRESSION TAG	UNP Q92630
A	52	HIS	-	EXPRESSION TAG	UNP Q92630
A	53	HIS	-	EXPRESSION TAG	UNP Q92630
A	54	HIS	-	EXPRESSION TAG	UNP Q92630
A	55	HIS	-	EXPRESSION TAG	UNP Q92630
A	56	HIS	-	EXPRESSION TAG	UNP Q92630
A	57	SER	-	EXPRESSION TAG	UNP Q92630
A	58	SER	-	EXPRESSION TAG	UNP Q92630
A	59	GLY	-	EXPRESSION TAG	UNP Q92630
A	60	VAL	-	EXPRESSION TAG	UNP Q92630
A	61	ASP	-	EXPRESSION TAG	UNP Q92630
A	62	LEU	-	EXPRESSION TAG	UNP Q92630
A	63	GLY	-	EXPRESSION TAG	UNP Q92630
A	64	THR	-	EXPRESSION TAG	UNP Q92630
A	65	GLU	-	EXPRESSION TAG	UNP Q92630
A	66	ASN	-	EXPRESSION TAG	UNP Q92630
A	67	LEU	-	EXPRESSION TAG	UNP Q92630
A	68	TYR	-	EXPRESSION TAG	UNP Q92630
A	69	PHE	-	EXPRESSION TAG	UNP Q92630
A	71	GLN	-	EXPRESSION TAG	UNP Q92630
A	72	SER	-	EXPRESSION TAG	UNP Q92630

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cl	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	58	Total	O	0	0
			58	58		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₂ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	84.28 Å 84.28 Å 148.50 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.54 – 2.36 40.54 – 2.36	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.54-2.36) 99.7 (40.54-2.36)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.37 Å)	Xtriage
Refinement program	REFMAC 5.5.0089	Depositor
R, R_{free}	0.225 , 0.288 0.222 , 0.280	Depositor DCC
R_{free} test set	1160 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	56.4	Xtriage
Anisotropy	0.397	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 40.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 22694 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3228	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, TPO, NA, SEP, SO4, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	1/3179 (0.0%)	0.76	0/4304

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	392	CYS	CB-SG	-5.13	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3157	0	2961	20	0
2	A	10	0	0	0	0
3	A	1	0	0	0	0
4	A	2	0	0	0	0
5	A	58	0	0	0	0
All	All	3228	0	2961	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:446:ASP:OD2	1:A:449:VAL:HG23	1.84	0.78
1:A:272:ILE:HG22	1:A:274:CYS:SG	2.35	0.67
1:A:154:VAL:HA	1:A:164:VAL:HG12	1.76	0.66
1:A:417:PRO:O	1:A:420:SER:HB3	2.01	0.60
1:A:434:LEU:HD22	1:A:460:PRO:HB2	1.84	0.60
1:A:171:VAL:O	1:A:171:VAL:HG12	2.07	0.55
1:A:378:ARG:O	1:A:381:ASN:HB2	2.08	0.54
1:A:369:SEP:HB2	1:A:372:LEU:HB2	1.91	0.53
1:A:133:ASP:OD1	1:A:133:ASP:N	2.41	0.53
1:A:272:ILE:CG2	1:A:274:CYS:SG	3.00	0.49
1:A:87:MET:HE1	1:A:99:HIS:HA	1.95	0.47
1:A:267:HIS:HA	1:A:328:MET:SD	2.56	0.46
1:A:171:VAL:HG12	1:A:173:GLN:HG2	1.98	0.45
1:A:394:VAL:HG12	1:A:396:THR:HG23	2.00	0.43
1:A:373:LEU:HB3	1:A:379:ALA:HB1	2.01	0.43
1:A:328:MET:N	1:A:329:PRO:CD	2.83	0.41
1:A:437:ASP:OD2	1:A:441:GLN:NE2	2.44	0.41
1:A:79:THR:HB	1:A:80:PRO:CD	2.51	0.41
1:A:113:LEU:O	1:A:114:ASN:CG	2.59	0.41
1:A:462:LEU:HA	1:A:462:LEU:HD23	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/429 (93%)	378 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	310/370 (84%)	301 (97%)	9 (3%)	50 64

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	ASN
1	A	133	ASP
1	A	253[A]	ARG
1	A	253[B]	ARG
1	A	332	MET
1	A	371	LYS
1	A	395	THR
1	A	402	VAL
1	A	411	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	HIS
1	A	199	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	A	159	1	8,9,10	1.72	3 (37%)	8,12,14	2.59	2 (25%)
1	TPO	A	308	1	8,10,11	0.99	1 (12%)	7,14,16	1.45	1 (14%)
1	PTR	A	309	1	14,16,17	1.93	1 (7%)	18,22,24	0.84	1 (5%)
1	SEP	A	369	1	8,9,10	1.65	2 (25%)	8,12,14	2.43	1 (12%)
1	SEP	A	385	1	8,9,10	1.63	1 (12%)	8,12,14	1.37	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	159	1	-	0/6/8/10	0/0/0/0
1	TPO	A	308	1	-	0/8/11/13	0/0/0/0
1	PTR	A	309	1	-	0/9/11/13	0/1/1/1
1	SEP	A	369	1	-	0/6/8/10	0/0/0/0
1	SEP	A	385	1	-	0/6/8/10	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	309	PTR	OH-CZ	-6.90	1.24	1.40
1	A	308	TPO	P-OG1	2.07	1.66	1.60
1	A	369	SEP	P-O2P	2.12	1.62	1.54
1	A	159	SEP	P-O3P	2.12	1.62	1.54
1	A	159	SEP	P-O2P	2.18	1.62	1.54
1	A	385	SEP	P-O1P	3.01	1.61	1.51
1	A	369	SEP	P-O1P	3.13	1.61	1.51
1	A	159	SEP	P-O1P	3.29	1.62	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	309	PTR	O-C-CA	-2.89	117.95	125.49
1	A	159	SEP	O-C-CA	-2.14	119.91	125.49
1	A	385	SEP	O-C-CA	-2.02	120.22	125.49
1	A	385	SEP	O2P-P-OG	2.21	112.91	106.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	308	TPO	C-CA-N	2.25	114.53	109.83
1	A	369	SEP	OG-CB-CA	5.87	113.28	108.27
1	A	159	SEP	OG-CB-CA	6.53	113.85	108.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	369	SEP	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1	-	4,4,4	0.48	0	6,6,6	0.42	0
2	SO4	A	2	-	4,4,4	0.81	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	2	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	402/429 (93%)	0.17	24 (5%)	25 38	14, 32, 54, 84	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	124	GLY	6.8
1	A	135	GLY	6.4
1	A	470	PRO	6.2
1	A	125	PRO	5.6
1	A	306	VAL	5.2
1	A	402	VAL	4.4
1	A	399	ASP	4.1
1	A	396	THR	3.9
1	A	136	SER	3.8
1	A	126	ASN	3.8
1	A	127	ASN	3.7
1	A	431	ASP	3.4
1	A	401	SER	3.0
1	A	394	VAL	2.7
1	A	137	TYR	2.6
1	A	400	GLY	2.4
1	A	466	LEU	2.4
1	A	374	ASP	2.3
1	A	133	ASP	2.3
1	A	395	THR	2.2
1	A	138	VAL	2.2
1	A	436	LEU	2.2
1	A	74	GLY	2.1
1	A	187	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	A	308	11/12	0.82	0.14	-	46,52,89,95	0
1	SEP	A	369	10/11	0.67	0.17	-	36,47,80,80	0
1	SEP	A	385	10/11	0.83	0.14	-	37,47,53,64	0
1	PTR	A	309	16/17	0.96	0.11	-	31,40,62,65	0
1	SEP	A	159	10/11	0.77	0.17	-	59,64,80,86	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NA	A	480	1/1	0.95	0.11	-0.56	41,41,41,41	0
2	SO4	A	1	5/5	0.94	0.15	-	60,68,78,97	0
2	SO4	A	2	5/5	0.93	0.22	-	46,55,72,87	0
4	CL	A	482	1/1	0.97	0.14	-	79,79,79,79	0
4	CL	A	481	1/1	0.99	0.07	-	61,61,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.