wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 11:27 PM BST

PDB ID	:	2KI8
Title	:	Solution NMR structure of tungsten formylmethanofuran dehydrogenase subunit D from Archaeoglobus fulgidus, Northeast Structural Genomics Consortium target AtT7
Authors	:	Eletsky, A.; Wu, Y.; Yee, A.; Fares, C.; Lee, H.W.; Arrowsmith, C.H.; Prestegard, J.H.; Szyperski, T.; Montelione, G.T.; Northeast Structural Genomics Consortium (NESG)
Deposited on	:	2009-04-28

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 96%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	146	54% 9% 37%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:23-A:31, A:47-A:109, A:122-A:141 (92)	0.24	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 5 single-model clusters were found.

Cluster number	Models
1	4, 10, 14, 17, 20
2	5, 8, 16
3	3, 13, 18
4	11, 12
5	1, 15
Single-model clusters	2; 6; 7; 9; 19

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2242 atoms, of which 1108 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Tungsten formylmethanofuran dehydrogenase, subunit D (FwdD-2).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	146	2242	713	1108	192	223	6	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP O28351
A	2	GLY	-	expression tag	UNP O28351
A	3	SER	-	expression tag	UNP O28351
A	4	SER	-	expression tag	UNP O28351
A	5	HIS	-	expression tag	UNP O28351
A	6	HIS	-	expression tag	UNP O28351
A	7	HIS	-	expression tag	UNP O28351
A	8	HIS	-	expression tag	UNP O28351
A	9	HIS	-	expression tag	UNP O28351
A	10	HIS	-	expression tag	UNP O28351
A	11	SER	-	expression tag	UNP O28351
A	12	SER	-	expression tag	UNP O28351
A	13	GLY	-	expression tag	UNP O28351
A	14	ARG	-	expression tag	UNP O28351
A	15	GLU	-	expression tag	UNP O28351
A	16	ASN	-	expression tag	UNP O28351
A	17	LEU	-	expression tag	UNP O28351
A	18	TYR	-	expression tag	UNP O28351
A	19	PHE	-	expression tag	UNP O28351
A	20	GLN	-	expression tag	UNP O28351
A	21	GLY	-	expression tag	UNP O28351
A	22	HIS	-	expression tag	UNP O28351
A	145	GLY	-	expression tag	UNP O28351
A	146	SER	-	expression tag	UNP O28351

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Tungsten formylmethanofuran dehydrogenase, subunit D (FwdD-2)

Chain A: 54% 9% 37%

M1 G2 S3 S4 H5 H6 H7 H8 H9 H10 S11 S12 G13 R14 E15 N16 L17 Y18 F19 Q20 G21 H22 E27 S30 G31 R32 T33 L34 N35 Q36 G37 A38 T39 V40 E41 E42 K43 L44 T45 E46 E47 Y48 N53 L67 T77 I98 P99 M100 M106 V107 I108 D109 P110

S111 T112 D113 G114 T115 G116 M117 P118 Q119 F120 K121 G125 K133 L140 L141 E142 A143 I144 G145 S146

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 3. Colouring as in section 4.1 above.

- Molecule 1: Tungsten formylmethanofuran dehydrogenase, subunit D (FwdD-2)

Chain A: 49% 13% • 37%

M1 G2 S3 S4 H5 H6 H7 H8 H9 H10 S11 S12 G13 R14 E15 N16 L17 Y18 F19 Q20 G21 H22 S30 G31 R32 T33 L34 N35 Q36 G37 A38 T39 V40 E41 E42 K43 L44 T45 E46 A51 V52 N53 I57 L65 G66 L67 Q68 E69 K74 V75 K76 T77 E81

I98 A104 N105 M106 V107 I108 D109 P110 S111 T112 D113 G114 T115 G116 M117 P118 Q119 F120 K121 G125 T126 V127 E128 K129 V137 L140 L141 E142 A143 I144 G145 S146

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	3.0
CNS	refinement	1.2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 16267
Number of chemical shift lists	1
Total number of shifts	1711
Number of shifts mapped to atoms	1711
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	96%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	716	720	720	9±2
All	All	14320	14400	14400	187

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

5 of 66 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:138:LYS:HD3	1:A:141:LEU:HD12	0.66	1.67	9	1
1:A:30:SER:HA	1:A:98:ILE:O	0.61	1.95	17	19
1:A:77:THR:HG22	1:A:125:GLY:HA3	0.59	1.74	15	14
1:A:106:MET:SD	1:A:140:LEU:HD21	0.58	2.38	15	17
1:A:74:LYS:HD3	1:A:76:LYS:HE3	0.58	1.76	3	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	92/146 (63%)	87±1 (94±2%)	5±1 (5±2%)	1±1 (1±1%)	34 78
All	All	1840/2920 (63%)	1731 (94%)	98 (5%)	11 (1%)	34 78

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	69	GLU	9
1	A	135	LEU	1
1	A	100	MET	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	78/123 (63%)	75±1 (96±2%)	3±1 (4±2%)	40 83
All	All	1560/2460 (63%)	1493 (96%)	67 (4%)	40 83

5 of 17 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	53	ASN	13
1	A	100	MET	13
1	A	133	LYS	9
1	A	48	TYR	6
1	A	81	GLU	5

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 96% for the well-defined parts and 87% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 16267

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1711
Number of shifts mapped to atoms	1711
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	135	-0.09 ± 0.09	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	120	-0.22 ± 0.10	None needed (< 0.5 ppm)
$^{13}C'$	135	0.13 ± 0.10	None needed (< 0.5 ppm)
^{15}N	129	0.37 ± 0.53	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 96%, i.e. 1059 atoms were assigned a chemical shift out of a possible 1104. 20 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	454/454 (100%)	181/181 (100%)	184/184 (100%)	89/89 (100%)
Sidechain	533/578 (92%)	326/333 (98%)	200/227 (88%)	7/18 (39%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	72/72 (100%)	38/38 (100%)	33/33 (100%)	1/1 (100%)
Overall	1059/1104 (96%)	545/552 (99%)	417/444 (94%)	97/108 (90%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	67	LEU	HB2	-0.37	3.32 – -0.08	-5.9
1	A	88	LYS	HB2	0.37	3.03 – 0.53	-5.6
1	A	67	LEU	HB3	-0.47	3.34 – -0.26	-5.6
1	A	88	LYS	HE3	1.91	3.86 – 1.96	-5.3

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: