wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 11:24 PM BST

PDB ID	:	2KIV
Title	:	AIDA-1 SAM domain tandem
Authors	:	Donaldson, L.W.; Kurabi, A.
Deposited on	:	2009-05-12

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 80%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	148	59% 17% 16% 9%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:2-A:31, A:35-A:46, A:51-A:62, A:76-A:133 (112)	0.47	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 3 single-model clusters were found.

Cluster number	Models
1	3, 4, 5, 8, 9, 11, 12, 14, 15, 17, 18, 19, 20
2	1, 2, 6, 13
Single-model clusters	7; 10; 16

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2134 atoms, of which 1075 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Ankyrin repeat and sterile alpha motif domain-containing protein 1B.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	135	2134	670	1075	187	197	5	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	EXPRESSION TAG	UNP Q7Z6G8
A	-11	HIS	-	EXPRESSION TAG	UNP Q7Z6G8
A	-10	HIS	-	EXPRESSION TAG	UNP Q7Z6G8
A	-9	HIS	-	EXPRESSION TAG	UNP Q7Z6G8
A	-8	HIS	-	EXPRESSION TAG	UNP Q7Z6G8
A	-7	HIS	-	EXPRESSION TAG	UNP Q7Z6G8
A	-6	HIS	-	EXPRESSION TAG	UNP Q7Z6G8
A	-5	LEU	-	EXPRESSION TAG	UNP Q7Z6G8
A	-4	VAL	-	EXPRESSION TAG	UNP Q7Z6G8
A	-3	PRO	-	EXPRESSION TAG	UNP Q7Z6G8
A	-2	ARG	-	EXPRESSION TAG	UNP Q7Z6G8
A	-1	GLY	-	EXPRESSION TAG	UNP Q7Z6G8
A	0	SER	-	EXPRESSION TAG	UNP Q7Z6G8
A	30	ALA	PHE	ENGINEERED	UNP Q7Z6G8
A	73	ALA	TYR	ENGINEERED	UNP Q7Z6G8
A	109	ALA	TRP	ENGINEERED	UNP Q7Z6G8

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Ankyrin repeat and sterile alpha motif domain-containing protein 1B

Chain A: 59% 17% 16% 9%

MET HIS HIS HIS HIS HIS HIS LEU VAL PRO ARG GLY SER V1 L8 H19 L20 M31 G32 S33 N34 V35 M36 L41 L42 E43 I44 G45 I46 L47 N48 S49 G50 H51 R52 I59 L62 P63 K64 M65 R66 P67 I68 G69 H70 D71 G72 A73 H74 P75 V78 W81

L87 Y90 F94 Y99 T100 L104 L105 L113 I119 R125 I128 D134 R135

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 9. Colouring as in section 4.1 above.

- Molecule 1: Ankyrin repeat and sterile alpha motif domain-containing protein 1B

Chain A: 57% 17% • 16% 9%

MET HIS HIS HIS HIS HIS HIS LEU VAL PRO ARG GLY SER V1 Q2 T3 G12 L13 H19 L20 N23 G24 F25 G32 S33 N34 D38 L41 I44 G45 I46 L47 N48 S49 G50 I59 D60 I61 L62 P63 K64 M65 R66 P67 I68 G69 H70 D71 G72 A73 H74 P75

V78 L87 F94 Y99 M102 D103 L104 L105 E112 L113 V116 I119 N120 L121 H124 D134 R135

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 500 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	refinement	2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 16297
Number of chemical shift lists	1
Total number of shifts	1413
Number of shifts mapped to atoms	1413
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	80%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	891	914	914	15±5
All	All	17820	18280	18280	301

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

5 of 147 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:113:LEU:HD11	1:A:119:ILE:HD12	0.83	1.51	16	13
1:A:79:ALA:HB2	1:A:91:THR:HG21	0.78	1.55	10	1
1:A:113:LEU:HD12	1:A:117:LEU:HD23	0.77	1.57	5	3
1:A:113:LEU:HD12	1:A:119:ILE:HD12	0.69	1.63	19	2
1:A:59:ILE:HD12	1:A:62:LEU:HD12	0.67	1.67	5	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	112/148 (76%)	97±3 (87±2%)	13±2 (11±2%)	2±1 (2±1%)	13 52
All	All	2240/2960 (76%)	1938 (87%)	253 (11%)	49 (2%)	13 52

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	45	GLY	18
1	A	44	ILE	14
1	A	88	GLY	9
1	A	86	GLU	2
1	A	35	VAL	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	97/127 (76%)	86±3 (88±4%)	11±3 (12±4%)	11 53
All	All	1940/2540 (76%)	1711 (88%)	229 (12%)	11 53

5 of 55 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	19	HIS	17
1	A	125	ARG	14
1	A	100	THR	12
1	A	20	LEU	12
1	A	52	ARG	9

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 80% for the well-defined parts and 74% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 16297

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1413
Number of shifts mapped to atoms	1413
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	122	-0.56 ± 0.14	Should be applied
$^{13}C_{\beta}$	111	0.20 ± 0.08	None needed (< 0.5 ppm)
$^{13}C'$	116	-0.34 ± 0.19	None needed (< 0.5 ppm)
^{15}N	113	1.19 ± 0.40	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 80%, i.e. 1121 atoms were assigned a chemical shift out of a possible 1406. 17 out of 24 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	520/558 (93%)	206/223 (92%)	212/224 (95%)	102/111 (92%)
Sidechain	532/761 (70%)	317/439 (72%)	209/291 (72%)	6/31 (19%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	69/87 (79%)	38/46 (83%)	29/36 (81%)	2/5 (40%)
Overall	1121/1406 (80%)	561/708 (79%)	450/551 (82%)	110/147 (75%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: