



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:06 PM GMT

PDB ID : 4KLS
Title : DNA polymerase beta mismatched reactant complex with Mn²⁺, 10 min
Authors : Freudenthal, B.D.; Beard, W.A.; Shock, D.D.; Wilson, S.H.
Deposited on : 2013-05-07
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	8664 (2.00-1.96)
Clashscore	102246	9905 (2.00-1.96)
Ramachandran outliers	100387	9792 (2.00-1.96)
Sidechain outliers	100360	9791 (2.00-1.96)
RSRZ outliers	91569	8679 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	<div> <div>15%</div> <div>73%</div> <div>21%</div> <div>••</div> </div>
2	D	5	<div> <div>60%</div> <div>40%</div> </div>
3	P	11	<div> <div>55%</div> <div>36%</div> <div>9%</div> </div>
4	T	16	<div> <div>6%</div> <div>56%</div> <div>44%</div> </div>

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 3562 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	6	0
			2632	1661	457	505	9			

- Molecule 2 is a DNA chain called 5'-D(P*GP*TP*CP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	5	Total	C	N	O	P	0	0	0
			106	49	20	32	5			


- Molecule 3 is a DNA chain called 5'-D(*GP*CP*TP*GP*AP*TP*GP*CP*GP*CP*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	11	Total	C	N	O	P	0	2	0
			243	116	46	70	11			

- Molecule 4 is a DNA chain called 5'-D(*CP*CP*GP*AP*CP*GP*GP*CP*GP*CP*AP*TP*CP*AP*GP*C)-3'.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	T	16	Total	C	N	O	P	0	0	0
			323	153	63	92	15			

- Molecule 5 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: C₁₀H₁₆N₅O₁₂P₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	
			30	10	5	12	3	

- Molecule 6 is PYROPHOSPHATE (three-letter code: PPV) (formula: $\text{H}_4\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	O P		
			9	7 2	0	1

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	3	Total 3	Na 3	0	0

- Molecule 8 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	P	1	Total 1	Mn 1	0	0
8	A	2	Total 2	Mn 2	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	144	Total 144	O 144	0	0
9	D	11	Total 11	O 11	0	0
9	P	29	Total 29	O 29	0	1
9	T	29	Total 29	O 29	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase beta



- Molecule 2: 5'-D(P*GP*TP*CP*GP*G)-3'



- Molecule 3: 5'-D(*GP*CP*TP*GP*AP*TP*GP*CP*GP*CP*A)-3'



- Molecule 4: 5'-D(*CP*CP*GP*AP*CP*GP*GP*CP*GP*CP*AP*TP*CP*AP*GP*C)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.92Å 78.61Å 55.03Å 90.00° 112.88° 90.00°	Depositor
Resolution (Å)	24.13 – 1.98 24.13 – 1.98	Depositor EDS
% Data completeness (in resolution range)	89.1 (24.13-1.98) 84.9 (24.13-1.98)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.98Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.226 , 0.285 0.229 , 0.277	Depositor DCC
R_{free} test set	1513 reflections (5.59%)	DCC
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.0	EDS
Estimated twinning fraction	0.036 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 27118 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3562	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, DTP, PPV, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	1/2684 (0.0%)	0.57	2/3608 (0.1%)
2	D	1.18	0/118	1.43	2/179 (1.1%)
3	P	1.36	4/272 (1.5%)	2.95	7/418 (1.7%)
4	T	0.85	1/362 (0.3%)	1.28	3/556 (0.5%)
All	All	0.63	6/3436 (0.2%)	1.13	14/4761 (0.3%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	10[A]	DC	O3'-P	-6.24	1.53	1.61
3	P	10[B]	DC	O3'-P	-6.24	1.53	1.61
1	A	189	GLY	C-N	-5.55	1.21	1.34
3	P	10[A]	DC	P-O5'	-5.51	1.54	1.59
3	P	10[B]	DC	P-O5'	-5.51	1.54	1.59
4	T	12	DT	C4'-O4'	-5.49	1.39	1.45

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	10[A]	DC	P-O3'-C3'	36.40	163.38	119.70
3	P	10[B]	DC	P-O3'-C3'	36.40	163.38	119.70
1	A	189	GLY	O-C-N	-5.99	113.11	122.70
2	D	2	DT	O4'-C1'-N1	-5.92	103.86	108.00
3	P	11[B]	DA	O4'-C1'-N9	5.84	112.09	108.00
1	A	190	ASP	O-C-N	-5.78	113.46	122.70
3	P	11[B]	DA	O4'-C4'-C3'	-5.70	102.22	104.50
3	P	10[A]	DC	C1'-O4'-C4'	-5.67	104.42	110.10
3	P	10[B]	DC	C1'-O4'-C4'	-5.67	104.42	110.10
4	T	13	DC	O4'-C1'-C2'	-5.46	101.53	105.90
4	T	14	DA	O4'-C1'-N9	-5.40	104.22	108.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	T	5	DC	C1'-O4'-C4'	-5.09	105.01	110.10
2	D	1	DG	O4'-C1'-N9	-5.08	104.45	108.00
3	P	4	DG	O4'-C1'-N9	-5.00	104.50	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2632	0	2610	84	0
2	D	106	0	57	0	0
3	P	243	0	136	11	0
4	T	323	0	179	2	0
5	A	30	0	11	5	0
6	A	9	0	0	1	0
7	A	3	0	0	0	0
8	A	2	0	0	0	0
8	P	1	0	0	0	0
9	A	144	0	0	4	0
9	D	11	0	0	0	0
9	P	29	0	0	0	0
9	T	29	0	0	0	0
All	All	3562	0	2993	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:282:MET:HG2	1:A:325:TRP:CZ3	1.82	1.14
1:A:282:MET:HG2	1:A:325:TRP:HZ3	0.94	1.09
1:A:287:LEU:HD21	1:A:292:THR:HG22	1.33	1.07
1:A:287:LEU:CD2	1:A:292:THR:HG22	1.85	1.06

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:GLU:HG2	1:A:204:SER:N	1.68	1.04
1:A:203:GLU:HG2	1:A:204:SER:H	0.90	1.04
1:A:203:GLU:CG	1:A:204:SER:H	1.71	1.01
1:A:282:MET:CE	1:A:320:PHE:CE2	2.49	0.96
1:A:36:TYR:CZ	1:A:40:ARG:HD2	2.04	0.93
1:A:203:GLU:CG	1:A:204:SER:N	2.26	0.91
1:A:282:MET:CE	1:A:320:PHE:HE2	1.90	0.83
6:A:402[B]:PPV:O12	3:P:11[B]:DA:H5''	1.80	0.81
1:A:282:MET:HE2	1:A:320:PHE:HE2	1.45	0.80
1:A:301:LEU:HG	1:A:302:GLY:H	1.45	0.80
1:A:183:ARG:O	1:A:331:LYS:HG2	1.85	0.76
1:A:301:LEU:HG	1:A:302:GLY:N	2.01	0.75
1:A:282:MET:CG	1:A:325:TRP:HZ3	1.88	0.71
1:A:276:ASP:HA	3:P:11[B]:DA:C5	2.26	0.70
1:A:287:LEU:HD23	1:A:292:THR:HG22	1.71	0.70
1:A:123:GLU:HA	1:A:126:ARG:HD2	1.77	0.66
1:A:170:ASP:HB3	1:A:173:TYR:CD2	2.32	0.64
1:A:276:ASP:HA	5:A:401[A]:DTP:C5	2.27	0.63
1:A:301:LEU:CG	1:A:302:GLY:H	2.11	0.63
1:A:276:ASP:HB3	3:P:11[B]:DA:N7	2.16	0.60
1:A:265:TYR:CZ	1:A:269:VAL:HG21	2.36	0.60
1:A:276:ASP:HB3	5:A:401[A]:DTP:N7	2.18	0.59
1:A:234:LYS:HE2	9:A:558:HOH:O	2.03	0.59
4:T:1:DC:H2''	4:T:2:DC:H5'	1.86	0.58
1:A:271:TYR:HH	5:A:401[A]:DTP:C2	2.16	0.57
1:A:311:LEU:HD13	1:A:322:TYR:CE1	2.40	0.57
1:A:276:ASP:HA	3:P:11[B]:DA:C4	2.40	0.56
1:A:282:MET:HE1	1:A:320:PHE:CE2	2.38	0.56
1:A:282:MET:HE2	1:A:320:PHE:CE2	2.27	0.55
1:A:159:GLN:NE2	9:A:507:HOH:O	2.39	0.55
1:A:287:LEU:CD2	1:A:292:THR:CG2	2.75	0.54
1:A:282:MET:HE3	1:A:320:PHE:CE2	2.41	0.54
4:T:2:DC:H2''	4:T:3:DG:C8	2.44	0.53
1:A:134:HIS:O	1:A:138:ILE:HD13	2.09	0.52
1:A:150:ILE:HG12	1:A:253:ARG:HD2	1.91	0.51
1:A:123:GLU:O	1:A:126:ARG:HG2	2.09	0.51
1:A:18:MET:SD	1:A:82:LEU:HD22	2.50	0.51
1:A:271:TYR:HE1	1:A:283:ARG:HH22	1.58	0.51
3:P:11[B]:DA:O4'	3:P:11[B]:DA:OP1	2.29	0.51
1:A:134:HIS:HE1	1:A:227:THR:O	1.94	0.50
1:A:283:ARG:HB2	1:A:283:ARG:CZ	2.42	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:ILE:HD13	1:A:59:ALA:HB2	1.93	0.50
1:A:103:VAL:HB	1:A:106:ILE:HD12	1.94	0.50
3:P:1:DG:H2"	3:P:2:DC:C6	2.46	0.50
1:A:209:LYS:O	1:A:213:GLN:HG3	2.12	0.49
1:A:277:ILE:HG22	1:A:335:GLU:HB3	1.94	0.49
1:A:96:SER:HB3	1:A:120:LYS:HB2	1.94	0.48
1:A:264:GLN:NE2	1:A:296[A]:TYR:CD1	2.82	0.48
1:A:203:GLU:HG3	1:A:204:SER:N	2.25	0.48
1:A:277:ILE:CG2	1:A:335:GLU:HB3	2.44	0.48
3:P:1:DG:H2"	3:P:2:DC:H6	1.79	0.47
1:A:271:TYR:O	5:A:401[A]:DTP:H2'1	2.13	0.47
1:A:264:GLN:NE2	1:A:296[B]:TYR:CD1	2.82	0.47
1:A:36:TYR:CE2	1:A:40:ARG:HD2	2.48	0.47
1:A:327:TYR:HD1	1:A:328:ARG:N	2.13	0.46
1:A:182:ARG:HD3	1:A:273:THR:OG1	2.15	0.46
1:A:317:LYS:HG3	1:A:327:TYR:CD2	2.50	0.46
1:A:198:PRO:HA	1:A:262:LYS:HD3	1.98	0.46
1:A:271:TYR:O	3:P:11[B]:DA:H2"	2.15	0.45
1:A:276:ASP:HA	5:A:401[A]:DTP:C4	2.46	0.45
1:A:36:TYR:CE1	1:A:40:ARG:HD2	2.48	0.45
1:A:287:LEU:HD23	1:A:292:THR:CG2	2.41	0.45
1:A:276:ASP:OD1	1:A:277:ILE:N	2.40	0.45
1:A:279:ASN:HB2	3:P:11[B]:DA:C2	2.52	0.44
1:A:289:LYS:HE2	1:A:324:GLN:CB	2.48	0.44
1:A:276:ASP:CA	3:P:11[B]:DA:C5	2.98	0.44
1:A:135:HIS:CD2	1:A:228:LEU:HD22	2.53	0.44
1:A:264:GLN:NE2	1:A:296[B]:TYR:HD1	2.15	0.44
1:A:264:GLN:NE2	1:A:296[A]:TYR:HD1	2.15	0.44
1:A:240:GLN:NE2	1:A:249:GLU:HG2	2.32	0.43
1:A:274:GLY:HA3	1:A:275:SER:HA	1.80	0.43
1:A:275:SER:N	1:A:278:PHE:HB3	2.35	0.42
1:A:289:LYS:HD2	1:A:323:ILE:O	2.20	0.42
1:A:271:TYR:CD2	1:A:295[A]:GLU:HB3	2.54	0.42
1:A:276:ASP:CB	3:P:11[B]:DA:C5	3.02	0.42
1:A:178:CYS:HA	1:A:182:ARG:HB2	2.01	0.42
1:A:275:SER:H	1:A:278:PHE:HB3	1.85	0.41
1:A:300:PRO:HD3	1:A:311:LEU:HG	2.02	0.41
1:A:183:ARG:O	1:A:331:LYS:HA	2.21	0.41
1:A:45:VAL:HG11	1:A:62:LEU:HB3	2.02	0.41
1:A:73:ILE:O	1:A:77:LEU:HG	2.21	0.41
1:A:208:PRO:HA	9:A:638:HOH:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:TYR:CD1	1:A:279:ASN:ND2	2.89	0.41
1:A:165:GLU:OE1	1:A:168:LYS:HD2	2.20	0.41
1:A:204:SER:HB2	9:A:607:HOH:O	2.20	0.40
1:A:190:ASP:OD1	1:A:190:ASP:C	2.60	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/335 (97%)	310 (95%)	15 (5%)	1 (0%)	46	39

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	204	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/295 (97%)	276 (96%)	10 (4%)	43	38

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	LEU
1	A	126	ARG
1	A	138	ILE
1	A	207	GLN
1	A	282	MET
1	A	283	ARG
1	A	315	SER
1	A	325	TRP
1	A	327	TYR
1	A	334	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	159	GLN
1	A	164	ASN
1	A	207	GLN
1	A	240	GLN
1	A	264	GLN
1	A	285	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	DTP	A	401[A]	8	24,32,32	1.32	5 (20%)	32,50,50	2.57	10 (31%)
6	PPV	A	402[B]	8	6,8,8	0.65	0	11,13,13	1.22	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DTP	A	401[A]	8	-	0/18/34/34	0/3/3/3
6	PPV	A	402[B]	8	-	0/6/6/6	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	401[A]	DTP	PG-O1G	-2.66	1.42	1.51
5	A	401[A]	DTP	C8-N7	-2.39	1.30	1.34
5	A	401[A]	DTP	C4-N3	-2.28	1.32	1.35
5	A	401[A]	DTP	PA-O1A	2.07	1.58	1.51
5	A	401[A]	DTP	PG-O2G	2.81	1.64	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	401[A]	DTP	N3-C2-N1	-8.61	122.30	128.89
5	A	401[A]	DTP	PA-O3A-PB	-4.70	119.54	132.73
5	A	401[A]	DTP	PB-O3B-PG	-3.83	119.81	132.67
5	A	401[A]	DTP	C4-C5-N7	-3.65	106.12	109.48
6	A	402[B]	PPV	P2-OPP-P1	-3.24	121.80	132.67
5	A	401[A]	DTP	C2'-C1'-N9	-2.53	108.01	114.16
5	A	401[A]	DTP	O2G-PG-O3B	2.27	115.39	105.09
5	A	401[A]	DTP	O2B-PB-O3B	2.80	117.79	105.09
5	A	401[A]	DTP	O4'-C1'-N9	3.26	113.37	107.72
5	A	401[A]	DTP	O2A-PA-O3A	3.79	122.30	105.09
5	A	401[A]	DTP	O3A-PA-O5'	4.81	115.69	102.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	401[A]	DTP	5	0
6	A	402[B]	PPV	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/335 (96%)	0.88	49 (15%) 3 4	16, 35, 74, 95	0
2	D	5/5 (100%)	-0.27	0 100 100	22, 25, 44, 52	0
3	P	11/11 (100%)	-0.20	0 100 100	24, 32, 40, 49	1 (9%)
4	T	16/16 (100%)	0.15	1 (6%) 23 27	26, 41, 63, 64	0
All	All	356/367 (97%)	0.80	50 (14%) 4 5	16, 35, 73, 95	1 (0%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	LYS	8.4
1	A	303	VAL	8.0
1	A	327	TYR	7.6
1	A	323	ILE	7.2
1	A	305	GLY	6.8
1	A	301	LEU	6.8
1	A	304	THR	6.8
1	A	306	VAL	6.5
1	A	246	ASP	6.4
1	A	302	GLY	6.4
1	A	284	ALA	5.9
1	A	290	GLY	4.9
1	A	257	ILE	4.6
1	A	293	ILE	4.5
1	A	277	ILE	4.3
1	A	193	VAL	3.9
1	A	325	TRP	3.8
1	A	326	LYS	3.8
1	A	203	GLU	3.8
1	A	247	GLU	3.8
1	A	288	GLU	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	324	GLN	3.4
1	A	332	ASP	3.4
1	A	255	ILE	3.3
1	A	287	LEU	3.2
1	A	206	LYS	3.0
1	A	292	THR	3.0
1	A	317	LYS	3.0
1	A	281	ASN	2.8
1	A	271	TYR	2.8
1	A	46	ILE	2.7
1	A	204	SER	2.7
1	A	162	VAL	2.6
1	A	235	PHE	2.6
1	A	140	LEU	2.6
1	A	320	PHE	2.6
1	A	276	ASP	2.6
1	A	279	ASN	2.6
1	A	282	MET	2.5
1	A	194	LEU	2.5
1	A	209	LYS	2.4
1	A	329	GLU	2.3
1	A	248	LYS	2.3
4	T	1	DC	2.3
1	A	195	LEU	2.2
1	A	272	PHE	2.2
1	A	307	ALA	2.1
1	A	283	ARG	2.1
1	A	335	GLU	2.0
1	A	42	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
8	MN	A	405	1/1	0.97	0.19	1.15	22,22,22,22	1
6	PPV	A	402[B]	9/9	0.88	0.15	0.37	32,40,42,48	9
5	DTP	A	401[A]	30/30	0.84	0.19	0.29	12,44,56,59	30
7	NA	A	403	1/1	0.99	0.10	-1.09	21,21,21,21	0
7	NA	A	407	1/1	0.95	0.08	-1.39	30,30,30,30	0
8	MN	A	406	1/1	0.94	0.07	-1.89	40,40,40,40	0
7	NA	A	404	1/1	0.97	0.21	-	27,27,27,27	0
8	MN	P	101	1/1	0.96	0.08	-	69,69,69,69	0

6.5 Other polymers

There are no such residues in this entry.