



wwPDB X-ray Structure Validation Summary Report

Jan 31, 2016 – 08:31 PM GMT

PDB ID : 1KNX
Title : HPr kinase/phosphatase from Mycoplasma pneumoniae
Authors : Allen, G.S.
Deposited on : 2001-12-19
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	 16% 58% 35% ..
1	B	312	 2% 67% 29% ..
1	C	312	 6% 65% 29% ..
1	D	312	 12% 66% 29% ..
1	E	312	 4% 73% 24% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	312	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into segments: a small red segment at the beginning labeled '5%', a large green segment in the middle labeled '71%', and a yellow segment at the end labeled '25%'. There are two small grey dots at the far right end of the bar.</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 14717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable HPr(Ser) kinase/phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	303	Total 2392	C 1546	N 404	O 436	S 6	0	0	0
1	B	304	Total 2415	C 1559	N 409	O 441	S 6	0	0	0
1	C	300	Total 2375	C 1532	N 403	O 434	S 6	0	0	0
1	D	304	Total 2400	C 1548	N 408	O 438	S 6	0	0	0
1	E	310	Total 2446	C 1576	N 414	O 450	S 6	0	0	0
1	F	306	Total 2407	C 1555	N 404	O 442	S 6	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	34	Total 34	O 34	0	0
2	B	55	Total 55	O 55	0	0
2	C	38	Total 38	O 38	0	0
2	D	39	Total 39	O 39	0	0
2	E	61	Total 61	O 61	0	0
2	F	55	Total 55	O 55	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable HPr(Ser) kinase/phosphatase




- Molecule 1: Probable HPr(Ser) kinase/phosphatase




- Molecule 1: Probable HPr(Ser) kinase/phosphatase







• Molecule 1: Probable HPr(Ser) kinase/phosphatase




• Molecule 1: Probable HPr(Ser) kinase/phosphatase



• Molecule 1: Probable HPr(Ser) kinase/phosphatase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	117.73Å 127.84Å 170.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.22 – 2.50 29.22 – 2.40	Depositor EDS
% Data completeness (in resolution range)	91.3 (29.22-2.50) 89.8 (29.22-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 2.39Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.227 , 0.280 0.229 , 0.280	Depositor DCC
R_{free} test set	4108 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	41.2	Xtrriage
Anisotropy	0.334	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.0	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	2 of 97608 reflections (0.002%)	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14717	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/2427	0.55	0/3270
1	B	0.41	0/2450	0.62	0/3301
1	C	0.39	0/2409	0.59	0/3245
1	D	0.41	0/2435	0.59	0/3283
1	E	0.46	0/2481	0.61	0/3346
1	F	0.48	1/2442 (0.0%)	0.66	1/3295 (0.0%)
All	All	0.42	1/14644 (0.0%)	0.60	1/19740 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	233	LEU	C-N	-6.51	1.19	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	235	LEU	C-N-CA	5.16	134.61	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	236	GLU	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2392	0	2488	106	0
1	B	2415	0	2520	62	0
1	C	2375	0	2478	70	0
1	D	2400	0	2490	70	0
1	E	2446	0	2528	55	0
1	F	2407	0	2491	59	0
2	A	34	0	0	0	0
2	B	55	0	0	3	0
2	C	38	0	0	3	0
2	D	39	0	0	0	0
2	E	61	0	0	2	0
2	F	55	0	0	1	0
All	All	14717	0	14995	402	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 402 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:282:ILE:HG12	1:B:218:GLN:HG2	1.21	1.11
1:E:292:ASN:HD22	1:E:295:LEU:H	1.05	0.98
1:B:90:THR:HG23	1:B:92:SER:H	1.28	0.97
1:E:292:ASN:ND2	1:E:295:LEU:H	1.64	0.95
1:A:218:GLN:NE2	1:A:218:GLN:H	1.62	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/312 (96%)	263 (88%)	31 (10%)	5 (2%)	11	19
1	B	300/312 (96%)	288 (96%)	12 (4%)	0	100	100
1	C	296/312 (95%)	279 (94%)	16 (5%)	1 (0%)	46	68
1	D	300/312 (96%)	265 (88%)	32 (11%)	3 (1%)	19	34
1	E	306/312 (98%)	289 (94%)	16 (5%)	1 (0%)	46	68
1	F	302/312 (97%)	285 (94%)	17 (6%)	0	100	100
All	All	1803/1872 (96%)	1669 (93%)	124 (7%)	10 (1%)	30	50

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	97	THR
1	A	94	THR
1	C	178	ASP
1	A	21	ASP
1	D	249	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/279 (95%)	248 (94%)	17 (6%)	22	39
1	B	270/279 (97%)	259 (96%)	11 (4%)	37	63
1	C	265/279 (95%)	250 (94%)	15 (6%)	25	46
1	D	266/279 (95%)	253 (95%)	13 (5%)	31	55
1	E	271/279 (97%)	253 (93%)	18 (7%)	21	38
1	F	267/279 (96%)	253 (95%)	14 (5%)	29	51
All	All	1604/1674 (96%)	1516 (94%)	88 (6%)	27	48

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	249	GLU
1	D	166	ASP
1	F	210	ASN
1	C	256	LEU
1	D	24	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	169	ASN
1	D	49	GLN
1	F	194	GLN
1	C	172	HIS
1	C	231	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/312 (97%)	0.63	49 (16%) 3 2	25, 65, 131, 145	0
1	B	304/312 (97%)	-0.12	7 (2%) 64 67	24, 44, 66, 100	0
1	C	300/312 (96%)	0.22	19 (6%) 23 26	26, 50, 90, 125	0
1	D	304/312 (97%)	0.58	37 (12%) 5 5	26, 53, 100, 113	0
1	E	310/312 (99%)	-0.08	13 (4%) 40 45	25, 41, 68, 108	0
1	F	306/312 (98%)	0.00	16 (5%) 31 35	24, 41, 90, 127	0
All	All	1827/1872 (97%)	0.20	141 (7%) 16 18	24, 46, 106, 145	0

The worst 5 of 141 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	70	VAL	5.9
1	E	236	GLU	5.8
1	C	311	GLU	5.3
1	D	107	GLN	5.3
1	A	115	PHE	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.