



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:09 PM GMT

PDB ID : 4KRV
Title : Crystal structure of catalytic domain of bovine beta1,4-galactosyltransferase mutant M344H-GalT1 complex with 6-sulfo-GlcNAc
Authors : Ramakrishnan, B.; Qasba, P.K.
Deposited on : 2013-05-17
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	
1	B	286	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	A	507	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4751 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-1,4-galactosyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	1	0
			2226	1429	385	400	12			
1	B	272	Total	C	N	O	S	0	1	0
			2226	1429	385	400	12			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	117	ALA	-	EXPRESSION TAG	UNP P08037
A	118	SER	-	EXPRESSION TAG	UNP P08037
A	119	MET	-	EXPRESSION TAG	UNP P08037
A	120	THR	-	EXPRESSION TAG	UNP P08037
A	121	GLY	-	EXPRESSION TAG	UNP P08037
A	122	GLY	-	EXPRESSION TAG	UNP P08037
A	123	GLN	-	EXPRESSION TAG	UNP P08037
A	124	GLN	-	EXPRESSION TAG	UNP P08037
A	125	MET	-	EXPRESSION TAG	UNP P08037
A	126	GLY	-	EXPRESSION TAG	UNP P08037
A	127	ARG	-	EXPRESSION TAG	UNP P08037
A	128	GLY	-	EXPRESSION TAG	UNP P08037
A	129	SER	-	EXPRESSION TAG	UNP P08037
A	342	THR	CYS	ENGINEERED MUTATION	UNP P08037
A	344	HIS	MET	ENGINEERED MUTATION	UNP P08037
B	117	ALA	-	EXPRESSION TAG	UNP P08037
B	118	SER	-	EXPRESSION TAG	UNP P08037
B	119	MET	-	EXPRESSION TAG	UNP P08037
B	120	THR	-	EXPRESSION TAG	UNP P08037
B	121	GLY	-	EXPRESSION TAG	UNP P08037
B	122	GLY	-	EXPRESSION TAG	UNP P08037
B	123	GLN	-	EXPRESSION TAG	UNP P08037
B	124	GLN	-	EXPRESSION TAG	UNP P08037
B	125	MET	-	EXPRESSION TAG	UNP P08037
B	126	GLY	-	EXPRESSION TAG	UNP P08037

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	127	ARG	-	EXPRESSION TAG	UNP P08037
B	128	GLY	-	EXPRESSION TAG	UNP P08037
B	129	SER	-	EXPRESSION TAG	UNP P08037
B	342	THR	CYS	ENGINEERED MUTATION	UNP P08037
B	344	HIS	MET	ENGINEERED MUTATION	UNP P08037

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is 6-AMINOHEXYL-URIDINE-C1,5'-DIPHOSPHATE (three-letter code: UDH) (formula: C₁₅H₂₇N₃O₁₂P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 32	C 15	N 3	O 12	P 2	0	0
3	B	1	Total 32	C 15	N 3	O 12	P 2	0	0

- Molecule 4 is SUGAR (2-(ACETYLAMINO)-2-DEOXY-6-O-SULFO-BETA-D-GLUCOPYRANOSE) (three-letter code: NGS) (formula: C₈H₁₅NO₉S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			19	8	1	9	1		
4	B	1	Total	C	N	O	S	0	0
			19	8	1	9	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	73	Total	O	0	0
			73	73		
6	B	72	Total	O	0	0
			72	72		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-1,4-galactosyltransferase 1



- Molecule 1: Beta-1,4-galactosyltransferase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.28Å 88.90Å 139.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.64 – 2.40 24.64 – 2.40	Depositor EDS
% Data completeness (in resolution range)	89.9 (24.64-2.40) 89.9 (24.64-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.89 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.188 , 0.249 0.197 , 0.248	Depositor DCC
R_{free} test set	1152 reflections (5.45%)	DCC
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 22371 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4751	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UDH, MN, SO4, NGS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/2290	0.40	0/3103
1	B	0.22	0/2290	0.40	0/3103
All	All	0.22	0/4580	0.40	0/6206

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2226	0	2191	18	0
1	B	2226	0	2191	21	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	32	0	25	0	0
3	B	32	0	25	0	0
4	A	19	0	14	0	0
4	B	19	0	14	0	0
5	A	30	0	0	0	0
5	B	20	0	0	0	0
6	A	73	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	72	0	0	0	0
All	All	4751	0	4460	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:258:MET:HE2	1:B:343:ARG:HG2	1.64	0.80
1:A:241:LYS:HB3	1:B:241:LYS:HE3	1.70	0.73
1:A:153:VAL:HG23	1:A:196:LYS:HE3	1.77	0.65
1:A:154:ASP:HB3	1:A:157:LEU:HB2	1.84	0.59
1:A:242:ASP:OD2	1:A:387:ARG:NH2	2.35	0.59
1:B:220:GLN:HB2	1:B:231:LEU:HD21	1.87	0.56
1:A:180:HIS:CE1	1:A:265:ARG:HD2	2.42	0.55
1:A:387:ARG:HG3	1:B:382:VAL:O	2.08	0.53
1:B:252:ASP:HB2	1:B:255:LEU:HD22	1.91	0.52
1:B:382:VAL:HA	1:B:396:VAL:HG12	1.91	0.51
1:B:146:LEU:HD22	1:B:147:ILE:N	2.26	0.51
1:A:241:LYS:HD3	1:B:241:LYS:HB3	1.93	0.48
1:B:255:LEU:HD23	1:B:293:VAL:HG23	1.95	0.48
1:B:275:VAL:HG22	1:B:340:GLY:HA3	1.94	0.48
1:B:233:ASN:HB3	1:B:378:LEU:HD22	1.96	0.48
1:B:180:HIS:CE1	1:B:265:ARG:HD2	2.49	0.47
1:A:318:ASP:OD1	1:A:318:ASP:N	2.47	0.47
1:B:147:ILE:HB	1:B:343:ARG:HG3	1.96	0.47
1:B:191:ARG:HH21	1:B:194:HIS:HD2	1.62	0.46
1:B:231:LEU:HD13	1:B:398:ILE:HD13	1.97	0.46
1:B:272:HIS:HB3	1:B:334:ARG:HG2	1.98	0.46
1:A:274:SER:HA	1:A:342:THR:HG21	1.98	0.46
1:A:258:MET:HE3	1:A:341:LYS:HB3	1.98	0.45
1:B:189:ARG:NH2	1:B:224:SER:O	2.39	0.44
1:A:351:LYS:HD2	1:A:352:LYS:HG2	2.00	0.44
1:B:146:LEU:HD11	1:B:148:GLU:HG3	1.99	0.44
1:A:277:MET:HE2	1:A:279:LYS:HE2	2.00	0.43
1:A:249:VAL:HG22	1:A:295:ALA:HB2	1.99	0.43
1:A:258:MET:HE3	1:A:282:PHE:HE1	1.83	0.43
1:A:233:ASN:HB3	1:A:378:LEU:HD22	2.01	0.43
1:B:191:ARG:HD2	1:B:194:HIS:CD2	2.55	0.41
1:A:139:PRO:HG2	1:A:140:LEU:HD12	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:255:LEU:HA	1:B:255:LEU:HD12	1.89	0.41
1:A:356:ASN:O	1:A:359:ARG:NH1	2.49	0.41
1:A:155:LEU:HD23	1:A:155:LEU:HA	1.91	0.40
1:B:351:LYS:HE3	1:B:351:LYS:HB3	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/286 (95%)	268 (99%)	3 (1%)	0	100	100
1	B	271/286 (95%)	263 (97%)	8 (3%)	0	100	100
All	All	542/572 (95%)	531 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/254 (97%)	239 (97%)	7 (3%)	51	72
1	B	246/254 (97%)	236 (96%)	10 (4%)	37	57
All	All	492/508 (97%)	475 (96%)	17 (4%)	43	64

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	LEU
1	A	167	LEU
1	A	223	GLU
1	A	231	LEU
1	A	255	LEU
1	A	342	THR
1	A	351	LYS
1	B	134	CYS
1	B	141	LEU
1	B	157	LEU
1	B	160	GLN
1	B	195	LEU
1	B	231	LEU
1	B	242	ASP
1	B	255	LEU
1	B	342	THR
1	B	349	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	HIS
1	B	194	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	UDH	A	502	2	25,33,33	1.92	3 (12%)	33,47,47	1.72	7 (21%)
4	NGS	A	503	-	19,19,19	1.75	6 (31%)	22,28,28	1.19	2 (9%)
5	SO4	A	504	-	4,4,4	0.21	0	6,6,6	0.08	0
5	SO4	A	505	-	4,4,4	0.19	0	6,6,6	0.08	0
5	SO4	A	506	-	4,4,4	0.21	0	6,6,6	0.10	0
5	SO4	A	507	-	4,4,4	0.21	0	6,6,6	0.08	0
5	SO4	A	508	-	4,4,4	0.20	0	6,6,6	0.08	0
5	SO4	A	509	-	4,4,4	0.21	0	6,6,6	0.10	0
3	UDH	B	502	2	25,33,33	1.95	3 (12%)	33,47,47	1.77	7 (21%)
4	NGS	B	503	-	19,19,19	1.71	6 (31%)	22,28,28	1.11	1 (4%)
5	SO4	B	504	-	4,4,4	0.23	0	6,6,6	0.07	0
5	SO4	B	505	-	4,4,4	0.23	0	6,6,6	0.06	0
5	SO4	B	506	-	4,4,4	0.21	0	6,6,6	0.09	0
5	SO4	B	507	-	4,4,4	0.20	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UDH	A	502	2	-	0/21/41/41	0/2/2/2
4	NGS	A	503	-	-	0/10/30/30	0/1/1/1
5	SO4	A	504	-	-	0/0/0/0	0/0/0/0
5	SO4	A	505	-	-	0/0/0/0	0/0/0/0
5	SO4	A	506	-	-	0/0/0/0	0/0/0/0
5	SO4	A	507	-	-	0/0/0/0	0/0/0/0
5	SO4	A	508	-	-	0/0/0/0	0/0/0/0
5	SO4	A	509	-	-	0/0/0/0	0/0/0/0
3	UDH	B	502	2	-	0/21/41/41	0/2/2/2
4	NGS	B	503	-	-	0/10/30/30	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SO4	B	504	-	-	0/0/0/0	0/0/0/0
5	SO4	B	505	-	-	0/0/0/0	0/0/0/0
5	SO4	B	506	-	-	0/0/0/0	0/0/0/0
5	SO4	B	507	-	-	0/0/0/0	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	NGS	C1-C2	-3.22	1.49	1.53
3	B	502	UDH	C3B-C2B	-3.22	1.44	1.53
3	A	502	UDH	C3B-C2B	-3.20	1.44	1.53
4	A	503	NGS	O6-S	-3.17	1.47	1.57
4	B	503	NGS	C1-C2	-3.10	1.49	1.53
4	B	503	NGS	O6-S	-3.09	1.47	1.57
4	A	503	NGS	O5-C1	-2.96	1.37	1.43
3	B	502	UDH	C3B-C4B	-2.82	1.45	1.53
4	B	503	NGS	O5-C1	-2.76	1.37	1.43
3	A	502	UDH	C3B-C4B	-2.72	1.45	1.53
4	A	503	NGS	O5-C5	-2.35	1.38	1.44
4	B	503	NGS	O5-C5	-2.29	1.38	1.44
4	B	503	NGS	O4-C4	-2.16	1.37	1.43
4	A	503	NGS	O4-C4	-2.06	1.38	1.43
4	A	503	NGS	C-N	3.25	1.46	1.34
4	B	503	NGS	C-N	3.26	1.46	1.34
3	A	502	UDH	O4-C4	7.78	1.43	1.24
3	B	502	UDH	O4-C4	7.84	1.43	1.24

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	UDH	PB-O3A-PA	-4.42	120.32	132.73
3	A	502	UDH	PB-O3A-PA	-3.71	122.32	132.73
4	A	503	NGS	O9-S-O7	-2.48	101.66	112.46
4	B	503	NGS	O9-S-O7	-2.43	101.87	112.46
3	B	502	UDH	C4B-O4'-C1B	-2.02	107.50	109.72
4	A	503	NGS	CH3-C-N	2.12	120.17	116.11
3	A	502	UDH	O4'-C1B-N1	2.13	112.58	108.08
3	A	502	UDH	O2B-C1'-C2'	2.18	117.02	108.85
3	B	502	UDH	O2B-C1'-C2'	2.18	117.02	108.85
3	B	502	UDH	O5'-C5B-C4B	2.18	117.16	109.12
3	A	502	UDH	O5'-C5B-C4B	2.22	117.31	109.12
3	A	502	UDH	O3A-PA-O5'	2.32	109.08	102.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	UDH	O3A-PA-O5'	2.59	109.82	102.94
3	B	502	UDH	O3A-PB-O2B	2.71	110.14	102.94
3	A	502	UDH	O3A-PB-O2B	2.76	110.25	102.94
3	B	502	UDH	C4-N3-C2	6.45	120.53	114.14
3	A	502	UDH	C4-N3-C2	6.58	120.66	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/286 (95%)	0.01	10 (3%) 45 46	13, 20, 40, 58	0
1	B	272/286 (95%)	0.14	18 (6%) 22 22	14, 24, 49, 64	0
All	All	544/572 (95%)	0.08	28 (5%) 32 32	13, 22, 43, 64	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	131	LEU	7.3
1	A	402	SER	5.7
1	A	132	THR	4.3
1	B	152	PRO	4.1
1	B	174	MET	3.8
1	B	402	SER	3.8
1	B	150	ASN	3.3
1	B	136	GLU	3.2
1	B	163	PRO	3.1
1	B	151	ILE	3.0
1	B	137	GLU	3.0
1	B	131	LEU	3.0
1	B	132	THR	2.8
1	B	351	LYS	2.8
1	A	177	ILE	2.7
1	A	185	ILE	2.5
1	A	251	SER	2.5
1	B	160	GLN	2.5
1	B	206	LEU	2.5
1	B	153	VAL	2.5
1	B	184	ILE	2.5
1	A	186	ILE	2.4
1	A	184	ILE	2.2
1	B	244	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	249	VAL	2.1
1	A	244	ASP	2.1
1	B	293	VAL	2.1
1	B	133	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	SO4	A	507	5/5	0.95	0.31	2.07	53,54,62,63	0
5	SO4	A	505	5/5	0.92	0.25	1.82	43,48,61,64	0
5	SO4	A	508	5/5	0.81	0.34	1.01	50,55,88,88	0
5	SO4	A	504	5/5	0.91	0.16	0.55	35,42,56,69	0
3	UDH	A	502	32/32	0.94	0.15	-0.36	13,17,34,56	0
3	UDH	B	502	32/32	0.96	0.13	-0.55	13,20,37,46	0
4	NGS	B	503	19/19	0.96	0.10	-0.77	17,24,33,40	0
4	NGS	A	503	19/19	0.96	0.12	-1.10	14,25,34,36	0
2	MN	B	501	1/1	0.99	0.06	-7.18	20,20,20,20	0
5	SO4	A	506	5/5	0.95	0.21	-	39,44,50,69	0
5	SO4	B	507	5/5	0.89	0.37	-	53,61,68,74	0
5	SO4	B	506	5/5	0.90	0.30	-	50,51,72,74	0
5	SO4	A	509	5/5	0.94	0.35	-	49,64,69,73	0
5	SO4	B	504	5/5	0.98	0.14	-	31,37,45,53	0
2	MN	A	501	1/1	0.99	0.06	-	17,17,17,17	0
5	SO4	B	505	5/5	0.95	0.20	-	44,51,66,68	0

6.5 Other polymers

There are no such residues in this entry.