



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 11:55 PM BST

PDB ID : 2KUU
Title : Solution Structure of K10 TLS RNA (GC mutant in upper helix)
Authors : Bullock, S.L.; Ringel, I.; Ish-Horowicz, D.; Lukavsky, P.J.
Deposited on : 2010-03-01

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | NMR archive (#Entries) |
|--------------|-----------------------------|---------------------------|
| Clashscore | 114402 | 11133 |
| RNA backbone | 3027 | 600 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 1 | A | 48 |  |

2 Ensemble composition and analysis ⓘ

This entry contains 12 models. This entry does not contain protein, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1523 atoms, of which 513 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called K10 TLS RNA.


| Mol | Chain | Residues | Atoms | | | | | | Trace |
|-----|-------|----------|-------|-----|-----|-----|-----|----|-------|
| 1 | A | 48 | Total | C | H | N | O | P | 0 |
| | | | 1523 | 455 | 513 | 173 | 335 | 47 | |

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: K10 TLS RNA




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: K10 TLS RNA




4.2.2 Score per residue for model 2

- Molecule 1: K10 TLS RNA




4.2.3 Score per residue for model 3

- Molecule 1: K10 TLS RNA




4.2.4 Score per residue for model 4

- Molecule 1: K10 TLS RNA




4.2.5 Score per residue for model 5

- Molecule 1: K10 TLS RNA




4.2.6 Score per residue for model 6

- Molecule 1: K10 TLS RNA




4.2.7 Score per residue for model 7

- Molecule 1: K10 TLS RNA




4.2.8 Score per residue for model 8

- Molecule 1: K10 TLS RNA




4.2.9 Score per residue for model 9

- Molecule 1: K10 TLS RNA




4.2.10 Score per residue for model 10

- Molecule 1: K10 TLS RNA




4.2.11 Score per residue for model 11

- Molecule 1: K10 TLS RNA



4.2.12 Score per residue for model 12

- Molecule 1: K10 TLS RNA



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *simulated annealing, restrained molecular dynamics with RDCs*.

Of the 250 calculated structures, 12 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version |
|---------------|--------------------|---------|
| X-PLOR NIH | structure solution | |
| X-PLOR NIH | refinement | |

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------------------|-------------|----------------------|
| | | RMSZ | #Z>5 | RMSZ | #Z>5 |
| 1 | A | 1.02±0.00 | 0±0/1128 (0.0±0.0%) | 1.62±0.00 | 11±1/1753 (0.6±0.0%) |
| All | All | 1.02 | 0/13536 (0.0%) | 1.62 | 136/21036 (0.6%) |

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) | Models | |
|-----|-------|-----|------|-------------|-------|-------------|----------|--------|-------|
| | | | | | | | | Worst | Total |
| 1 | A | 14 | G | O4'-C1'-N9 | 6.43 | 113.34 | 108.20 | 12 | 12 |
| 1 | A | 36 | U | C5'-C4'-C3' | -6.00 | 106.41 | 116.00 | 12 | 12 |
| 1 | A | 17 | U | O4'-C1'-N1 | 5.91 | 112.93 | 108.20 | 11 | 4 |
| 1 | A | 26 | U | C5'-C4'-C3' | -5.71 | 106.86 | 116.00 | 3 | 12 |
| 1 | A | 16 | G | O4'-C1'-N9 | 5.46 | 112.57 | 108.20 | 2 | 10 |
| 1 | A | 23 | A | C3'-C2'-C1' | 5.43 | 105.84 | 101.50 | 7 | 9 |
| 1 | A | 22 | U | C3'-C2'-C1' | 5.42 | 105.84 | 101.50 | 2 | 7 |
| 1 | A | 25 | U | C5'-C4'-C3' | -5.37 | 107.40 | 116.00 | 3 | 1 |
| 1 | A | 34 | A | C5'-C4'-C3' | -5.26 | 107.59 | 116.00 | 2 | 12 |
| 1 | A | 40 | A | C5'-C4'-C3' | -5.25 | 107.60 | 116.00 | 3 | 12 |
| 1 | A | 35 | C | C5'-C4'-C3' | -5.22 | 107.65 | 116.00 | 12 | 10 |
| 1 | A | 1 | G | C5'-C4'-C3' | -5.21 | 107.66 | 116.00 | 9 | 2 |
| 1 | A | 42 | U | C5'-C4'-C3' | -5.21 | 107.67 | 116.00 | 3 | 12 |
| 1 | A | 47 | C | C5'-C4'-C3' | -5.21 | 107.67 | 116.00 | 12 | 12 |
| 1 | A | 22 | U | C5'-C4'-C3' | -5.18 | 107.71 | 116.00 | 10 | 7 |
| 1 | A | 4 | U | O4'-C1'-N1 | 5.02 | 112.22 | 108.20 | 7 | 2 |

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|-----|-------|-------|----------|----------|---------|
| 1 | A | 1010 | 513 | 513 | 34±4 |
| All | All | 12120 | 6156 | 6156 | 406 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

All unique clashes are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|---------------|---------------|----------|-------------|--------|-------|
| | | | | Worst | Total |
| 1:A:24:A:O2' | 1:A:25:U:H2' | 0.92 | 1.62 | 11 | 11 |
| 1:A:12:A:O2' | 1:A:13:U:H5' | 0.82 | 1.74 | 12 | 12 |
| 1:A:22:U:O2' | 1:A:23:A:H5'' | 0.81 | 1.76 | 2 | 1 |
| 1:A:1:G:H2' | 1:A:2:G:N7 | 0.80 | 1.91 | 12 | 3 |
| 1:A:22:U:H2' | 1:A:24:A:OP2 | 0.76 | 1.80 | 3 | 2 |
| 1:A:46:G:O2' | 1:A:47:C:H5' | 0.74 | 1.83 | 9 | 12 |
| 1:A:36:U:O2 | 1:A:36:U:H2' | 0.68 | 1.85 | 12 | 5 |
| 1:A:43:U:HO2' | 1:A:44:A:H8 | 0.67 | 1.32 | 8 | 2 |
| 1:A:1:G:H2' | 1:A:2:G:C8 | 0.65 | 2.26 | 9 | 12 |
| 1:A:36:U:H2' | 1:A:36:U:O2 | 0.65 | 1.88 | 1 | 3 |
| 1:A:34:A:H2' | 1:A:35:C:O4' | 0.65 | 1.91 | 10 | 10 |
| 1:A:24:A:HO2' | 1:A:25:U:H6 | 0.65 | 1.29 | 1 | 8 |
| 1:A:13:U:H2' | 1:A:14:G:O4' | 0.64 | 1.93 | 2 | 12 |
| 1:A:24:A:O2' | 1:A:25:U:H5'' | 0.64 | 1.92 | 3 | 1 |
| 1:A:23:A:H2' | 1:A:24:A:C8 | 0.61 | 2.31 | 11 | 9 |
| 1:A:42:U:H2' | 1:A:43:U:O4' | 0.59 | 1.98 | 4 | 12 |
| 1:A:23:A:H8 | 1:A:23:A:O5' | 0.59 | 1.81 | 2 | 2 |
| 1:A:39:A:H2' | 1:A:40:A:C8 | 0.58 | 2.33 | 1 | 12 |
| 1:A:24:A:O2' | 1:A:25:U:C6 | 0.58 | 2.56 | 10 | 11 |
| 1:A:1:G:C2' | 1:A:2:G:C8 | 0.57 | 2.88 | 12 | 3 |
| 1:A:22:U:C2' | 1:A:23:A:H5'' | 0.56 | 2.30 | 2 | 1 |
| 1:A:23:A:C8 | 1:A:23:A:O5' | 0.56 | 2.59 | 10 | 2 |
| 1:A:3:C:O2' | 1:A:4:U:H5' | 0.56 | 2.01 | 9 | 2 |
| 1:A:8:U:H2' | 1:A:9:U:C6 | 0.54 | 2.38 | 2 | 12 |
| 1:A:22:U:H1' | 1:A:26:U:O2 | 0.54 | 2.03 | 11 | 1 |
| 1:A:43:U:O2' | 1:A:44:A:H8 | 0.54 | 1.85 | 2 | 3 |
| 1:A:45:A:H2' | 1:A:46:G:C8 | 0.53 | 2.38 | 1 | 12 |

Continued on next page...

Continued from previous page...

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|--------------|--------------|----------|-------------|--------|-------|
| | | | | Worst | Total |
| 1:A:32:A:H2' | 1:A:33:C:C6 | 0.53 | 2.39 | 6 | 12 |
| 1:A:47:C:O5' | 1:A:47:C:H6 | 0.53 | 1.87 | 5 | 3 |
| 1:A:31:C:H2' | 1:A:32:A:C8 | 0.53 | 2.38 | 4 | 11 |
| 1:A:47:C:H6 | 1:A:47:C:O5' | 0.53 | 1.87 | 4 | 6 |
| 1:A:23:A:O5' | 1:A:23:A:C8 | 0.53 | 2.62 | 3 | 1 |
| 1:A:29:U:H2' | 1:A:30:A:O4' | 0.52 | 2.04 | 11 | 1 |
| 1:A:22:U:O2' | 1:A:24:A:C8 | 0.51 | 2.64 | 4 | 5 |
| 1:A:11:U:H2' | 1:A:12:A:C8 | 0.51 | 2.41 | 6 | 11 |
| 1:A:19:A:H2' | 1:A:20:A:C8 | 0.51 | 2.40 | 1 | 11 |
| 1:A:2:G:C8 | 1:A:2:G:O5' | 0.51 | 2.64 | 9 | 2 |
| 1:A:16:G:H2' | 1:A:17:U:C6 | 0.49 | 2.42 | 5 | 11 |
| 1:A:20:A:H2' | 1:A:21:U:O4' | 0.49 | 2.08 | 10 | 7 |
| 1:A:24:A:O2' | 1:A:25:U:H6 | 0.49 | 1.91 | 3 | 1 |
| 1:A:4:U:H2' | 1:A:5:U:C6 | 0.49 | 2.43 | 9 | 11 |
| 1:A:2:G:H2' | 1:A:3:C:C6 | 0.49 | 2.43 | 2 | 11 |
| 1:A:28:U:H2' | 1:A:29:U:C6 | 0.48 | 2.44 | 2 | 8 |
| 1:A:47:C:O2' | 1:A:48:C:H5' | 0.48 | 2.08 | 10 | 3 |
| 1:A:41:A:H2' | 1:A:42:U:C6 | 0.48 | 2.44 | 12 | 11 |
| 1:A:36:U:O2 | 1:A:36:U:C2' | 0.48 | 2.62 | 1 | 2 |
| 1:A:26:U:H2' | 1:A:27:C:C6 | 0.48 | 2.44 | 1 | 8 |
| 1:A:2:G:O5' | 1:A:2:G:C8 | 0.48 | 2.67 | 12 | 1 |
| 1:A:44:A:H2' | 1:A:45:A:C8 | 0.47 | 2.43 | 2 | 2 |
| 1:A:23:A:C6 | 1:A:24:A:C4 | 0.47 | 3.02 | 2 | 2 |
| 1:A:7:A:O2' | 1:A:8:U:H5' | 0.47 | 2.09 | 10 | 3 |
| 1:A:24:A:C6 | 1:A:26:U:H4' | 0.47 | 2.45 | 2 | 2 |
| 1:A:24:A:C2' | 1:A:25:U:C5' | 0.46 | 2.93 | 3 | 1 |
| 1:A:21:U:O2' | 1:A:22:U:H5' | 0.46 | 2.10 | 7 | 1 |
| 1:A:43:U:O2' | 1:A:44:A:C8 | 0.45 | 2.69 | 4 | 7 |
| 1:A:29:U:H2' | 1:A:30:A:C8 | 0.45 | 2.47 | 5 | 8 |
| 1:A:37:A:H2' | 1:A:38:C:C6 | 0.45 | 2.47 | 7 | 9 |
| 1:A:7:A:H2' | 1:A:8:U:C6 | 0.45 | 2.47 | 2 | 4 |
| 1:A:24:A:O3' | 1:A:25:U:H6 | 0.45 | 1.95 | 5 | 10 |
| 1:A:23:A:C6 | 1:A:24:A:C5 | 0.45 | 3.05 | 3 | 1 |
| 1:A:2:G:O5' | 1:A:2:G:H8 | 0.45 | 1.94 | 9 | 2 |
| 1:A:38:C:H2' | 1:A:39:A:C8 | 0.45 | 2.48 | 10 | 8 |
| 1:A:30:A:H2' | 1:A:31:C:C6 | 0.44 | 2.47 | 11 | 9 |
| 1:A:27:C:H2' | 1:A:28:U:C6 | 0.44 | 2.47 | 2 | 3 |
| 1:A:23:A:N6 | 1:A:24:A:C6 | 0.44 | 2.86 | 3 | 1 |
| 1:A:13:U:C2' | 1:A:14:G:O4' | 0.44 | 2.65 | 11 | 1 |
| 1:A:26:U:C4 | 1:A:27:C:N4 | 0.44 | 2.86 | 6 | 3 |
| 1:A:10:G:H2' | 1:A:11:U:C6 | 0.43 | 2.47 | 2 | 1 |

Continued on next page...

Continued from previous page...

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|--------------|---------------|----------|-------------|--------|-------|
| | | | | Worst | Total |
| 1:A:33:C:H2' | 1:A:34:A:C8 | 0.43 | 2.48 | 4 | 1 |
| 1:A:35:C:H4' | 1:A:36:U:OP1 | 0.43 | 2.13 | 4 | 1 |
| 1:A:2:G:H8 | 1:A:2:G:O5' | 0.43 | 1.96 | 10 | 1 |
| 1:A:22:U:O2' | 1:A:24:A:H8 | 0.43 | 1.97 | 3 | 1 |
| 1:A:14:G:H2' | 1:A:15:U:C6 | 0.43 | 2.48 | 4 | 3 |
| 1:A:17:U:H2' | 1:A:18:A:C8 | 0.42 | 2.49 | 11 | 1 |
| 1:A:23:A:O2' | 1:A:24:A:O4' | 0.41 | 2.36 | 5 | 1 |
| 1:A:25:U:H3' | 1:A:26:U:C5' | 0.41 | 2.45 | 3 | 1 |
| 1:A:22:U:H2' | 1:A:23:A:H5'' | 0.41 | 1.92 | 12 | 1 |
| 1:A:36:U:H2' | 1:A:37:A:C8 | 0.41 | 2.51 | 9 | 2 |
| 1:A:12:A:H2' | 1:A:13:U:C6 | 0.40 | 2.51 | 2 | 1 |
| 1:A:22:U:C2 | 1:A:26:U:C2 | 0.40 | 3.09 | 4 | 1 |

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers | Suiteness |
|-----|-------|---------------|-------------------|-----------------|-----------|
| 1 | A | 47/48 (98%) | 28±2 (60±4%) | 5±1 (11±2%) | 0.07±0.01 |
| All | All | 566/576 (98%) | 338 (60%) | 62 (11%) | 0.07 |

The overall RNA backbone suiteness is 0.07.

All unique RNA backbone outliers are listed below:

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 1 | A | 27 | C | 12 |
| 1 | A | 26 | U | 12 |
| 1 | A | 35 | C | 12 |
| 1 | A | 12 | A | 12 |
| 1 | A | 43 | U | 12 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 1 | A | 6 | G | 12 |
| 1 | A | 46 | G | 12 |
| 1 | A | 17 | U | 12 |
| 1 | A | 45 | A | 12 |
| 1 | A | 24 | A | 12 |
| 1 | A | 2 | G | 12 |
| 1 | A | 14 | G | 12 |
| 1 | A | 39 | A | 12 |
| 1 | A | 36 | U | 12 |
| 1 | A | 41 | A | 12 |
| 1 | A | 37 | A | 12 |
| 1 | A | 25 | U | 12 |
| 1 | A | 23 | A | 12 |
| 1 | A | 33 | C | 12 |
| 1 | A | 30 | A | 12 |
| 1 | A | 28 | U | 11 |
| 1 | A | 3 | C | 10 |
| 1 | A | 31 | C | 10 |
| 1 | A | 16 | G | 10 |
| 1 | A | 11 | U | 10 |
| 1 | A | 18 | A | 9 |
| 1 | A | 21 | U | 9 |
| 1 | A | 44 | A | 9 |
| 1 | A | 4 | U | 6 |
| 1 | A | 34 | A | 4 |
| 1 | A | 22 | U | 4 |
| 1 | A | 9 | U | 2 |
| 1 | A | 40 | A | 2 |
| 1 | A | 32 | A | 1 |
| 1 | A | 20 | A | 1 |

All unique RNA pucker outliers are listed below:

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 1 | A | 26 | U | 12 |
| 1 | A | 36 | U | 11 |
| 1 | A | 40 | A | 11 |
| 1 | A | 23 | A | 9 |
| 1 | A | 22 | U | 8 |
| 1 | A | 2 | G | 6 |
| 1 | A | 34 | A | 3 |
| 1 | A | 1 | G | 2 |

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided