



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:13 PM GMT

PDB ID : 4KYD
Title : Partial Structure of the C-terminal domain of the HPIV4B phosphoprotein, fused to MBP.
Authors : Yegambaram, K.; Bulloch, E.M.M.; Kingston, R.L
Deposited on : 2013-05-28
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	 8% 90% 5% • 5%
1	B	420	 8% 84% 10% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MPO	A	1503	-	-	-	X
3	MPO	A	1505	-	-	-	X
3	MPO	B	1503	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Maltose-binding periplasmic protein, Phosphoprotein, chimeric construct.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	0	4	0
			3105	1994	506	597	8			
1	B	400	Total	C	N	O	S	0	1	0
			3089	1984	505	592	8			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	82	ALA	ASP	ENGINEERED MUTATION	UNP P0AEX9
A	83	ALA	LYS	ENGINEERED MUTATION	UNP P0AEX9
A	359	ALA	GLU	ENGINEERED MUTATION	UNP P0AEX9
A	362	ALA	LYS	ENGINEERED MUTATION	UNP P0AEX9
A	363	ALA	ASP	ENGINEERED MUTATION	UNP P0AEX9
A	367	ASN	-	LINKER	UNP P0AEX9
A	368	ALA	-	LINKER	UNP P0AEX9
A	369	ALA	-	LINKER	UNP P0AEX9
A	370	ALA	-	LINKER	UNP P0AEX9
A	1368	SER	CYS	ENGINEERED MUTATION	UNP P21738
A	1400	TYR	-	EXPRESSION TAG	UNP P21738
B	82	ALA	ASP	ENGINEERED MUTATION	UNP P0AEX9
B	83	ALA	LYS	ENGINEERED MUTATION	UNP P0AEX9
B	359	ALA	GLU	ENGINEERED MUTATION	UNP P0AEX9
B	362	ALA	LYS	ENGINEERED MUTATION	UNP P0AEX9
B	363	ALA	ASP	ENGINEERED MUTATION	UNP P0AEX9
B	367	ASN	-	LINKER	UNP P0AEX9
B	368	ALA	-	LINKER	UNP P0AEX9
B	369	ALA	-	LINKER	UNP P0AEX9
B	370	ALA	-	LINKER	UNP P0AEX9
B	1368	SER	CYS	ENGINEERED MUTATION	UNP P21738
B	1400	TYR	-	EXPRESSION TAG	UNP P21738

- Molecule 2 is SUGAR (MALTOSE) (three-letter code: MAL) (formula: C₁₂H₂₂O₁₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			23	12	11		
2	B	1	Total	C	O	0	0
			23	12	11		

- Molecule 3 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (three-letter code: MPO) (formula: $C_7H_{15}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
3	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
3	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	148	Total	O	0	3
			151	151		
4	B	47	Total	O	0	1
			48	48		

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	142.83Å 142.83Å 114.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.10 – 2.21 38.10 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.4 (38.10-2.21) 99.5 (38.10-2.21)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.68 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.174 , 0.222 0.174 , 0.221	Depositor DCC
R_{free} test set	2138 reflections (5.21%)	DCC
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 54.5	EDS
Estimated twinning fraction	0.017 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 43151 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6517	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPO, MAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/3187	0.72	0/4325
1	B	0.51	0/3162	0.60	0/4291
All	All	0.59	0/6349	0.66	0/8616

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3105	0	3093	16	0
1	B	3089	0	3074	28	0
2	A	23	0	22	0	0
2	B	23	0	22	0	0
3	A	52	0	60	3	0
3	B	26	0	30	0	0
4	A	151	0	0	1	0
4	B	48	0	0	0	0
All	All	6517	0	6301	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:126:PRO:HD2	1:B:224:MET:HE3	1.37	1.04
1:A:1362:ARG:HH11	1:A:1362:ARG:HB3	1.31	0.94
1:A:44:GLU:OE1	1:A:44:GLU:N	2.09	0.85
1:A:157:THR:HG23	4:A:1710:HOH:O	1.81	0.79
1:B:126:PRO:HD2	1:B:224:MET:CE	2.14	0.76
1:B:115:LEU:HD21	1:B:224:MET:CE	2.18	0.74
1:B:311:LEU:HB3	1:B:317:ILE:HD13	1.77	0.67
1:B:115:LEU:HD21	1:B:224:MET:HE1	1.77	0.65
1:A:1362:ARG:HH11	1:A:1362:ARG:CB	2.08	0.64
1:B:41:ASP:O	1:B:46:LYS:HE3	1.99	0.63
1:A:1365:PHE:HB2	1:A:1371[B]:SER:OG	1.98	0.63
1:B:115:LEU:HD21	1:B:224:MET:HE2	1.82	0.61
1:B:10:TRP:HB3	1:B:43:LEU:HD11	1.82	0.61
1:A:205[B]:ASN:OD1	1:A:207[B]:ASP:OD1	2.20	0.60
1:B:325:GLN:OE1	1:B:325:GLN:HA	2.05	0.56
1:A:164:ASP:OD1	3:A:1505:MPO:H72	2.05	0.56
1:A:152:GLN:HB2	3:A:1503:MPO:H11	1.88	0.55
1:B:68:GLY:HA3	1:B:332:ASN:O	2.11	0.49
1:A:68:GLY:HA3	1:A:332:ASN:O	2.12	0.49
1:A:1362:ARG:NH1	1:A:1362:ARG:HB3	2.14	0.49
1:A:152:GLN:CB	3:A:1503:MPO:H11	2.43	0.49
1:B:229:PRO:O	1:B:298:PRO:HB2	2.11	0.49
1:A:168:ALA:O	1:A:181:VAL:HA	2.15	0.46
1:A:171:TYR:OH	1:A:174:GLY:HA2	2.16	0.45
1:B:278:GLU:HG3	1:B:282:ASN:ND2	2.31	0.45
1:B:62:TRP:HB3	1:B:67:PHE:HE1	1.80	0.45
1:B:72:GLN:OE1	1:B:1359:GLU:HG3	2.17	0.45
1:B:126:PRO:HG2	1:B:224:MET:HE1	1.99	0.44
1:B:51:ALA:HA	1:B:55:ASP:O	2.19	0.43
1:A:229:PRO:HA	1:A:232:TRP:CE2	2.54	0.43
1:B:6:LYS:HG2	1:B:34:LYS:HB3	1.99	0.43
1:B:11:ILE:O	1:B:39:HIS:HA	2.18	0.42
1:B:346:ALA:HB2	1:B:364:ALA:HB2	2.00	0.42
1:B:128:THR:O	1:B:131:GLU:HG2	2.19	0.42
1:B:229:PRO:HA	1:B:232:TRP:CE2	2.54	0.42
1:B:1374:LEU:O	1:B:1378:ILE:HD12	2.20	0.42
1:B:126:PRO:CD	1:B:224:MET:CE	2.93	0.42
1:A:6:LYS:HA	1:A:33:ILE:HG23	2.02	0.42
1:B:1361:ILE:CD1	1:B:1378:ILE:HD13	2.49	0.41
1:A:1:LYS:HB3	1:A:55:ASP:OD2	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:PRO:O	1:B:52:ALA:HB2	2.21	0.41
1:B:40:PRO:HD2	1:B:43:LEU:HD13	2.03	0.40
1:B:1:LYS:HB3	1:B:2:ILE:H	1.63	0.40
1:B:71:ALA:HB2	1:B:104:ILE:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/420 (96%)	396 (98%)	7 (2%)	0	100	100
1	B	399/420 (95%)	389 (98%)	10 (2%)	0	100	100
All	All	802/840 (96%)	785 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/335 (96%)	314 (98%)	7 (2%)	60	72
1	B	318/335 (95%)	310 (98%)	8 (2%)	55	67
All	All	639/670 (95%)	624 (98%)	15 (2%)	60	70

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLU
1	A	55	ASP
1	A	202	LYS
1	A	258	PHE
1	A	1362	ARG
1	A	1371[A]	SER
1	A	1371[B]	SER
1	B	31	THR
1	B	55	ASP
1	B	104	ILE
1	B	142	LYS
1	B	258	PHE
1	B	278	GLU
1	B	358	ASP
1	B	1367	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MAL	A	1501	-	24,24,24	1.14	2 (8%)	35,35,35	1.12	4 (11%)
3	MPO	A	1502	-	12,13,13	2.04	1 (8%)	15,17,17	2.46	4 (26%)
3	MPO	A	1503	-	12,13,13	2.01	1 (8%)	15,17,17	2.81	1 (6%)
3	MPO	A	1504	-	12,13,13	2.00	1 (8%)	15,17,17	1.78	3 (20%)
3	MPO	A	1505	-	12,13,13	1.96	1 (8%)	15,17,17	2.19	3 (20%)
2	MAL	B	1501	-	24,24,24	1.25	3 (12%)	35,35,35	1.26	4 (11%)
3	MPO	B	1502	-	12,13,13	2.04	1 (8%)	15,17,17	1.32	2 (13%)
3	MPO	B	1503	-	12,13,13	2.07	1 (8%)	15,17,17	2.18	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAL	A	1501	-	-	0/8/48/48	0/2/2/2
3	MPO	A	1502	-	-	0/7/15/15	0/1/1/1
3	MPO	A	1503	-	-	0/7/15/15	0/1/1/1
3	MPO	A	1504	-	-	0/7/15/15	0/1/1/1
3	MPO	A	1505	-	-	0/7/15/15	0/1/1/1
2	MAL	B	1501	-	-	0/8/48/48	0/2/2/2
3	MPO	B	1502	-	-	0/7/15/15	0/1/1/1
3	MPO	B	1503	-	-	0/7/15/15	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1501	MAL	O3'-C3'	2.00	1.47	1.43
2	B	1501	MAL	O5-C1	2.44	1.48	1.41
2	B	1501	MAL	O1-C1	2.49	1.48	1.41
2	A	1501	MAL	C1'-C2'	2.52	1.57	1.52
2	A	1501	MAL	O5'-C1'	2.67	1.48	1.43
3	A	1502	MPO	O3-S1	6.14	1.62	1.46
3	A	1505	MPO	O3-S1	6.45	1.63	1.46
3	A	1504	MPO	O3-S1	6.63	1.63	1.46
3	A	1503	MPO	O3-S1	6.71	1.63	1.46
3	B	1502	MPO	O3-S1	6.79	1.64	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1503	MPO	O3-S1	6.99	1.64	1.46

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1502	MPO	C3-C2-C1	-3.34	106.88	112.37
2	B	1501	MAL	C1-O1-C4'	-3.29	109.42	118.01
3	A	1504	MPO	O3-S1-O2	-2.57	105.62	111.61
2	A	1501	MAL	O1'-C1'-O5'	-2.42	103.63	110.25
3	A	1502	MPO	O2-S1-O1	-2.31	105.05	113.48
2	A	1501	MAL	O6-C6-C5	-2.04	104.59	111.33
2	A	1501	MAL	C1'-O5'-C5'	2.11	117.37	113.47
2	B	1501	MAL	O5-C5-C6	2.19	111.89	106.36
3	A	1504	MPO	C6-O4-C5	2.30	117.62	109.89
2	A	1501	MAL	O5'-C1'-C2'	2.36	113.57	109.80
3	B	1502	MPO	O2-S1-C1	2.44	108.98	106.91
2	B	1501	MAL	C1'-O5'-C5'	2.49	118.07	113.47
2	B	1501	MAL	C1'-C2'-C3'	2.50	114.15	110.43
3	B	1502	MPO	O1-S1-C1	2.56	109.09	106.91
3	A	1505	MPO	C3-C2-C1	2.80	116.98	112.37
3	B	1503	MPO	O1-S1-C1	3.33	109.75	106.91
3	A	1505	MPO	O2-S1-C1	3.49	109.88	106.91
3	A	1502	MPO	O1-S1-C1	4.82	111.02	106.91
3	A	1504	MPO	O2-S1-C1	5.67	111.74	106.91
3	A	1505	MPO	O1-S1-C1	5.92	111.96	106.91
3	B	1503	MPO	O2-S1-C1	6.46	112.42	106.91
3	A	1502	MPO	O2-S1-C1	6.95	112.84	106.91
3	A	1503	MPO	O1-S1-C1	10.29	115.68	106.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1503	MPO	2	0
3	A	1505	MPO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/420 (95%)	-0.17	1 (0%) 95 95	22, 37, 64, 94	0
1	B	400/420 (95%)	0.33	32 (8%) 15 14	41, 62, 111, 139	0
All	All	801/840 (95%)	0.08	33 (4%) 41 39	22, 50, 99, 139	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	31	THR	6.2
1	B	27	PHE	5.1
1	B	84	ALA	4.5
1	B	283	TYR	4.4
1	B	32	GLY	4.3
1	B	80	THR	4.2
1	B	33	ILE	4.0
1	B	141	ALA	3.4
1	B	1367	ASP	3.3
1	B	310	GLU	2.7
1	B	281	GLU	2.7
1	B	341	TYR	2.6
1	B	1	LYS	2.6
1	B	28	GLU	2.6
1	B	85	PHE	2.5
1	B	53	THR	2.5
1	B	13	GLY	2.5
1	B	1372	ARG	2.5
1	B	354	ARG	2.5
1	B	275	LEU	2.4
1	B	1369	ASP	2.3
1	B	285	LEU	2.3
1	B	52	ALA	2.3
1	B	6	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	7	LEU	2.2
1	B	278	GLU	2.2
1	B	282	ASN	2.2
1	B	54	GLY	2.2
1	A	29	LYS	2.1
1	B	304	LEU	2.1
1	B	81	PRO	2.1
1	B	43	LEU	2.0
1	B	79	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MPO	A	1503	13/13	0.87	0.43	33.63	55,92,123,125	0
3	MPO	A	1505	13/13	0.82	0.27	4.08	44,63,84,86	0
3	MPO	B	1503	13/13	0.91	0.20	3.23	72,79,91,91	0
2	MAL	A	1501	23/23	0.97	0.18	1.61	23,31,35,36	0
3	MPO	A	1504	13/13	0.89	0.22	1.16	77,81,92,93	0
3	MPO	A	1502	13/13	0.99	0.14	0.17	26,30,35,36	0
2	MAL	B	1501	23/23	0.94	0.14	-0.10	35,45,51,59	0
3	MPO	B	1502	13/13	0.96	0.12	-0.36	69,73,80,81	0

6.5 Other polymers [i](#)

There are no such residues in this entry.