wwPDB X-ray Structure Validation Summary Report (i)

Feb 1, 2016 – 10:11 AM GMT

PDB ID	:	3L4O
Title	:	Crystal Structure of the MauG/pre-Methylamine Dehydrogenase Complex After Treatment with Hydrogen Peroxide
Authors	:	Jensen, L.M.R.; Wilmot, C.M.
Deposited on	:	2009-12-21
Resolution	:	2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1192 (2.04-2.04)
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	2% 83% 10% • 5%
1	B	373	% 84% 10% • 5%
2	C	137	% 82% 9% 9%
2	E	137	78% 11% •• 9%
3	D	386	2% 86% 11% • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	386	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PG4	F	388	-	-	-	X

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 14721 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylamine utilization protein mauG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	353	2735	1708	490	526	11	0	1	0
1	B	355	2749	1716	492	530	11	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	368	HIS	-	EXPRESSION TAG	UNP Q51658
A	369	HIS	-	EXPRESSION TAG	UNP Q51658
A	370	HIS	-	EXPRESSION TAG	UNP Q51658
A	371	HIS	-	EXPRESSION TAG	UNP Q51658
A	372	HIS	-	EXPRESSION TAG	UNP Q51658
A	373	HIS	-	EXPRESSION TAG	UNP Q51658
B	368	HIS	-	EXPRESSION TAG	UNP Q51658
B	369	HIS	-	EXPRESSION TAG	UNP Q51658
B	370	HIS	-	EXPRESSION TAG	UNP Q51658
B	371	HIS	-	EXPRESSION TAG	UNP Q51658
B	372	HIS	-	EXPRESSION TAG	UNP Q51658
B	373	HIS	-	EXPRESSION TAG	UNP Q51658

- Molecule 2 is a protein called Methylamine dehydrogenase light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	125	958	592	161	191	14	0	1	0
2	E	125	958	592	161	191	14	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	132	HIS	-	EXPRESSION TAG	UNP Q51658
C	133	HIS	-	EXPRESSION TAG	UNP Q51658
C	134	HIS	-	EXPRESSION TAG	UNP Q51658
C	135	HIS	-	EXPRESSION TAG	UNP Q51658
C	136	HIS	-	EXPRESSION TAG	UNP Q51658
C	137	HIS	-	EXPRESSION TAG	UNP Q51658
E	132	HIS	-	EXPRESSION TAG	UNP Q51658
E	133	HIS	-	EXPRESSION TAG	UNP Q51658
E	134	HIS	-	EXPRESSION TAG	UNP Q51658
E	135	HIS	-	EXPRESSION TAG	UNP Q51658
E	136	HIS	-	EXPRESSION TAG	UNP Q51658
E	137	HIS	-	EXPRESSION TAG	UNP Q51658

- Molecule 3 is a protein called Methylamine dehydrogenase heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	376	2923	1853	502	560	8	0	0	0
			Total	C	N	O	S			
3	F	376	2923	1853	502	560	8	0	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	B	1	1	1	0	0
			Total	Ca		
4	A	1	1	1	0	0

- Molecule 5 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
5	A	1	43	34	1	4	4	0	0
			Total	C	Fe	N	O		
5	A	1	43	34	1	4	4	0	0
			Total	C	Fe	N	O		
5	B	1	43	34	1	4	4	0	0
			Total	C	Fe	N	O		
5	B	1	43	34	1	4	4	0	0

- Molecule 6 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	F	1	16	10	6	0	0

- Molecule 7 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	F	1	13	8	5	0	0

- Molecule 8 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	F	1	4	2	2	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	A	227	227	227	0	0
			Total	O		
9	B	288	288	288	0	0
			Total	O		
9	C	85	85	85	0	0
			Total	O		
9	D	228	228	228	0	0
			Total	O		
9	E	107	107	107	0	0
			Total	O		
9	F	333	333	333	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methylamine utilization protein mauG

Chain A:



- Molecule 1: Methylamine utilization protein mauG

Chain B:



- Molecule 2: Methylamine dehydrogenase light chain

Chain C:



- Molecule 2: Methylamine dehydrogenase light chain

Chain E:

- Molecule 3: Methylamine dehydrogenase heavy chain

Chain D:



- Molecule 3: Methylamine dehydrogenase heavy chain

Chain F:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	55.53Å 83.52Å 107.78Å 109.94° 91.54° 105.78°	Depositor
Resolution (Å)	38.20 – 2.05 38.20 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.0 (38.20-2.05) 85.0 (38.20-2.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	2.49 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.142 , 0.194 0.156 , 0.205	Depositor DCC
R_{free} test set	5304 reflections (5.25%)	DCC
Wilson B-factor (Å^2)	27.8	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 37.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 106311 reflections	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	14721	wwPDB-VP
Average B, all atoms (Å^2)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, HEC, 1PE, PG4, ACT, TRQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	1.02	1/2802 (0.0%)	1.03	13/3802 (0.3%)
1	B	1.12	3/2816 (0.1%)	1.16	13/3821 (0.3%)
2	C	0.98	0/969	0.91	1/1323 (0.1%)
2	E	1.17	0/969	0.95	2/1323 (0.2%)
3	D	0.96	0/3000	0.85	2/4088 (0.0%)
3	F	1.20	4/3000 (0.1%)	0.94	9/4088 (0.2%)
All	All	1.08	8/13556 (0.1%)	0.99	40/18445 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
2	E	0	1
All	All	0	3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	199	TRP	CB-CG	7.75	1.64	1.50
3	F	223	GLU	CG-CD	6.97	1.62	1.51
3	F	213	GLU	CG-CD	6.32	1.61	1.51
1	A	19	VAL	CB-CG2	5.46	1.64	1.52
1	B	53	VAL	CB-CG1	5.29	1.64	1.52

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	39	ARG	NE-CZ-NH2	-22.57	109.01	120.30
1	B	252	ARG	NE-CZ-NH2	-21.51	109.55	120.30
1	B	39	ARG	NE-CZ-NH1	19.15	129.88	120.30
1	B	252	ARG	NE-CZ-NH1	18.05	129.33	120.30
1	A	39	ARG	NE-CZ-NH2	-16.85	111.88	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	7	ASP	Peptide
1	B	39	ARG	Sidechain
2	E	130	ALA	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2735	0	2611	23	0
1	B	2749	0	2622	19	0
2	C	958	0	862	12	0
2	E	958	0	862	12	0
3	D	2923	0	2808	20	0
3	F	2923	0	2808	17	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	86	0	60	2	0
5	B	86	0	60	2	0
6	F	16	0	22	0	0
7	F	13	0	18	1	0
8	F	4	0	3	0	0
9	A	227	0	0	2	0
9	B	288	0	0	2	0
9	C	85	0	0	1	0
9	D	228	0	0	0	0
9	E	107	0	0	0	0
9	F	333	0	0	1	0
All	All	14721	0	12736	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:LYS:H	1:B:62:HIS:HE1	1.13	0.93
1:A:210:GLN:HE22	2:C:44:THR:HG21	1.42	0.85
1:B:48:LYS:H	1:B:62:HIS:CE1	1.96	0.81
1:A:206:MET:HE2	1:A:206:MET:HA	1.60	0.81
1:B:285:THR:HG22	1:B:286:ASP:OD1	1.80	0.80

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/373 (94%)	342 (97%)	10 (3%)	0	100	100
1	B	354/373 (95%)	347 (98%)	7 (2%)	0	100	100
2	C	123/137 (90%)	119 (97%)	4 (3%)	0	100	100
2	E	123/137 (90%)	120 (98%)	3 (2%)	0	100	100
3	D	374/386 (97%)	356 (95%)	16 (4%)	2 (0%)	34	22
3	F	374/386 (97%)	361 (96%)	12 (3%)	1 (0%)	46	36
All	All	1700/1792 (95%)	1645 (97%)	52 (3%)	3 (0%)	52	43

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	102	ILE
3	F	102	ILE
3	D	179	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/292 (95%)	261 (94%)	16 (6%)	25	14
1	B	278/292 (95%)	265 (95%)	13 (5%)	32	23
2	C	105/112 (94%)	103 (98%)	2 (2%)	65	60
2	E	105/112 (94%)	99 (94%)	6 (6%)	25	15
3	D	304/311 (98%)	292 (96%)	12 (4%)	39	30
3	F	304/311 (98%)	295 (97%)	9 (3%)	48	41
All	All	1373/1430 (96%)	1315 (96%)	58 (4%)	36	28

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	352	ARG
3	D	94	HIS
3	F	213	GLU
1	B	357	LEU
2	C	71	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	60	GLN
1	B	91	GLN
3	D	372	GLN
1	B	29	GLN
3	F	14	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	TRQ	C	57	2	14,17,18	1.53	1 (7%)	8,24,26	2.05	3 (37%)
2	TRQ	E	57	2	14,17,18	1.54	3 (21%)	8,24,26	1.78	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRQ	C	57	2	-	0/3/19/21	0/2/2/2
2	TRQ	E	57	2	-	0/3/19/21	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	57	TRQ	CD1-NE1	-2.65	1.32	1.36
2	E	57	TRQ	CD2-CE2	-2.13	1.37	1.40
2	E	57	TRQ	CZ3-CE3	3.95	1.40	1.34
2	C	57	TRQ	CZ3-CE3	4.92	1.42	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	E	57	TRQ	CB-CG-CD1	-3.35	123.83	127.97
2	C	57	TRQ	CB-CG-CD1	-3.10	124.14	127.97
2	E	57	TRQ	CD1-NE1-CE2	2.22	110.08	104.34
2	C	57	TRQ	CD1-NE1-CE2	2.29	110.25	104.34
2	C	57	TRQ	O7-CZ2-CE2	2.59	124.98	122.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	57	TRQ	3	0
2	E	57	TRQ	2	0

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	HEC	A	500	1	24,50,50	1.68	7 (29%)	19,82,82	3.35	10 (52%)
5	HEC	A	600	1	24,50,50	2.09	9 (37%)	19,82,82	3.66	9 (47%)
5	HEC	B	500	1	24,50,50	1.53	6 (25%)	19,82,82	3.03	7 (36%)
5	HEC	B	600	1	24,50,50	1.81	8 (33%)	19,82,82	3.80	7 (36%)
6	1PE	F	387	-	15,15,15	0.67	0	14,14,14	0.44	0
7	PG4	F	388	-	12,12,12	0.67	0	11,11,11	0.48	0
8	ACT	F	389	-	1,3,3	1.41	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEC	A	500	1	-	0/6/54/54	0/0/8/8
5	HEC	A	600	1	-	0/6/54/54	0/0/8/8

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEC	B	500	1	-	0/6/54/54	0/0/8/8
5	HEC	B	600	1	-	0/6/54/54	0/0/8/8
6	1PE	F	387	-	-	0/13/13/13	0/0/0/0
7	PG4	F	388	-	-	0/10/10/10	0/0/0/0
8	ACT	F	389	-	-	0/0/0/0	0/0/0/0

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	600	HEC	C4C-NC	-4.20	1.31	1.36
5	A	600	HEC	C4B-NB	-3.95	1.31	1.36
5	A	600	HEC	C1A-NA	-3.59	1.31	1.36
5	A	600	HEC	C4A-NA	-3.18	1.32	1.36
5	B	600	HEC	C3C-C2C	-3.13	1.37	1.40

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	600	HEC	CBB-CAB-C3B	-10.69	103.59	127.35
5	B	600	HEC	CBB-CAB-C3B	-9.93	105.28	127.35
5	A	500	HEC	CBB-CAB-C3B	-9.56	106.11	127.35
5	B	500	HEC	CBB-CAB-C3B	-8.59	108.27	127.35
5	B	600	HEC	CBC-CAC-C3C	-8.32	108.87	127.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	500	HEC	1	0
5	A	600	HEC	1	0
5	B	600	HEC	2	0
7	F	388	PG4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	353/373 (94%)	-0.37	6 (1%)	73	78	14, 21, 32, 48	0
1	B	355/373 (95%)	-0.44	4 (1%)	82	86	14, 20, 34, 55	0
2	C	124/137 (90%)	-0.06	1 (0%)	87	90	17, 21, 31, 53	0
2	E	124/137 (90%)	-0.24	0	100	100	12, 17, 27, 51	0
3	D	376/386 (97%)	-0.31	7 (1%)	70	75	13, 20, 30, 54	0
3	F	376/386 (97%)	-0.37	3 (0%)	87	90	13, 17, 28, 50	0
All	All	1708/1792 (95%)	-0.34	21 (1%)	81	84	12, 20, 31, 55	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	6	ALA	6.8
3	D	208	THR	4.8
1	B	7	ASP	4.8
3	F	208	THR	3.8
3	F	11	GLN	3.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
2	TRQ	C	57	16/17	0.96	0.12	-	24,29,36,44	0
2	TRQ	E	57	16/17	0.96	0.15	-	20,25,35,37	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
7	PG4	F	388	13/13	0.84	0.16	2.90	52,57,68,69	0
4	CA	A	400	1/1	0.99	0.10	1.91	25,25,25,25	0
4	CA	B	400	1/1	0.99	0.10	1.57	20,20,20,20	0
6	1PE	F	387	16/16	0.87	0.14	0.87	39,48,61,63	0
5	HEC	A	600	43/43	0.99	0.11	0.58	21,26,28,30	0
5	HEC	B	600	43/43	0.99	0.11	0.12	13,18,21,22	0
5	HEC	A	500	43/43	0.98	0.09	-0.19	23,26,28,29	0
5	HEC	B	500	43/43	0.99	0.08	-0.20	17,22,25,26	0
8	ACT	F	389	4/4	0.96	0.10	-	36,39,40,40	0

6.5 Other polymers (i)

There are no such residues in this entry.