



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 12:39 AM BST

PDB ID : 2L9S  
Title : Solution structure of Pf1 SID1-mSin3A PAH2 Complex  
Authors : Senthil Kumar, G.; Xie, T.; Zhang, Y.; Radhakrishnan, I.  
Deposited on : 2011-02-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 84%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	45	
2	B	94	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:208-A:223, B:302-B:343, B:351-B:380 (88)	0.49	11

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 4, 7, 10, 11, 12, 13, 18, 19
2	8, 9, 16
3	3, 17
4	6, 15
5	5, 14
Single-model clusters	20

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2196 atoms, of which 1080 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PHD finger protein 12.

Mol	Chain	Residues	Atoms						Trace
1	A	45	Total	C	H	N	O	S	0
			696	218	346	61	69	2	

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	197	SER	-	EXPRESSION TAG	UNP Q96QT6
A	198	ASN	-	EXPRESSION TAG	UNP Q96QT6
A	199	ALA	-	EXPRESSION TAG	UNP Q96QT6

- Molecule 2 is a protein called Paired amphipathic helix protein Sin3a.

Mol	Chain	Residues	Atoms					Trace
2	B	94	Total	C	H	N	O	0
			1500	482	734	134	150	

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	292	SER	-	EXPRESSION TAG	UNP Q60520
B	293	ASN	-	EXPRESSION TAG	UNP Q60520
B	294	ALA	-	EXPRESSION TAG	UNP Q60520

## 4 Residue-property plots


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PHD finger protein 12



- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2.2 Score per residue for model 2

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2.3 Score per residue for model 3

- Molecule 1: PHD finger protein 12



- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2.4 Score per residue for model 4

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2.5 Score per residue for model 5

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2.6 Score per residue for model 6

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a



### 4.2.7 Score per residue for model 7

- Molecule 1: PHD finger protein 12



- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2.8 Score per residue for model 8

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2.9 Score per residue for model 9

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




### 4.2.10 Score per residue for model 10

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a






#### 4.2.11 Score per residue for model 11 (medoid)

- Molecule 1: PHD finger protein 12



- Molecule 2: Paired amphipathic helix protein Sin3a




#### 4.2.12 Score per residue for model 12

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




#### 4.2.13 Score per residue for model 13

- Molecule 1: PHD finger protein 12



- Molecule 2: Paired amphipathic helix protein Sin3a



S385

#### 4.2.14 Score per residue for model 14

- Molecule 1: PHD finger protein 12

Chain A:  31% 64%



- Molecule 2: Paired amphipathic helix protein Sin3a


Chain B:  63% 14% 23%



#### 4.2.15 Score per residue for model 15

- Molecule 1: PHD finger protein 12

Chain A:  27% 9% 64%



- Molecule 2: Paired amphipathic helix protein Sin3a

Chain B:  68% 9% 23%




#### 4.2.16 Score per residue for model 16

- Molecule 1: PHD finger protein 12

Chain A:  24% 11% 64%




- Molecule 2: Paired amphipathic helix protein Sin3a




#### 4.2.17 Score per residue for model 17

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




#### 4.2.18 Score per residue for model 18

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




#### 4.2.19 Score per residue for model 19

- Molecule 1: PHD finger protein 12




- Molecule 2: Paired amphipathic helix protein Sin3a




4.2.20 Score per residue for model 20

- Molecule 1: PHD finger protein 12



- Molecule 2: Paired amphipathic helix protein Sin3a



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 80 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	structure solution	1.2
ARIA	refinement	1.2
ARIA	structure solution	1.2
ARIA	refinement	1.2

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2l9s_cs.str
Number of chemical shift lists	2
Total number of shifts	1547
Number of shifts mapped to atoms	1547
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	84%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality

### 6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	127	131	131	3±1
2	B	608	599	597	6±2
All	All	14700	14600	14560	165

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:214:ILE:HG12	2:B:332:LEU:HD11	0.80	1.52	10	8
1:A:214:ILE:HA	2:B:332:LEU:HD21	0.76	1.54	3	1
2:B:317:ARG:HD3	2:B:369:GLN:HG2	0.73	1.59	18	1
2:B:338:GLU:HB3	2:B:351:PRO:HB3	0.72	1.61	7	5
2:B:378:GLN:HA	2:B:378:GLN:HE21	0.72	1.44	3	1
2:B:321:GLN:HE21	2:B:321:GLN:HA	0.72	1.43	5	1
2:B:334:THR:HA	2:B:337:LYS:HE2	0.72	1.61	18	1
1:A:208:ARG:HB2	1:A:211:GLU:HB2	0.70	1.61	4	5
1:A:208:ARG:HB2	1:A:211:GLU:HG2	0.70	1.63	15	1
1:A:212:LEU:HD13	1:A:212:LEU:H	0.69	1.47	18	3
1:A:208:ARG:HB2	1:A:211:GLU:CB	0.66	2.20	16	3
1:A:212:LEU:HD11	2:B:304:PHE:HA	0.63	1.67	11	1
2:B:338:GLU:HG3	2:B:353:LEU:HB2	0.61	1.71	17	2

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:304:PHE:O	2:B:308:ILE:HG12	0.60	1.95	11	11
2:B:334:THR:O	2:B:338:GLU:HG2	0.59	1.97	14	3
2:B:375:GLU:HA	2:B:378:GLN:HE21	0.58	1.57	16	1
1:A:218:MET:HA	1:A:221:ASN:ND2	0.57	2.14	17	2
1:A:208:ARG:O	1:A:212:LEU:HB2	0.56	2.00	11	1
1:A:209:PRO:HB2	2:B:379:PHE:HA	0.56	1.78	2	1
1:A:219:GLU:HG3	1:A:220:ARG:HE	0.56	1.61	12	1
1:A:208:ARG:HB2	1:A:211:GLU:HB3	0.56	1.76	16	2
2:B:317:ARG:HE	2:B:317:ARG:HA	0.55	1.61	19	3
2:B:317:ARG:HA	2:B:317:ARG:HE	0.55	1.60	9	3
1:A:220:ARG:NE	1:A:220:ARG:HA	0.55	2.16	16	4
2:B:369:GLN:HB3	2:B:372:LEU:HB3	0.55	1.78	11	1
2:B:310:TYR:O	2:B:314:ILE:HG13	0.53	2.03	7	8
2:B:338:GLU:HG3	2:B:353:LEU:HD23	0.53	1.79	12	1
2:B:360:ALA:O	2:B:364:ARG:HG2	0.53	2.04	9	1
2:B:323:ASP:HA	2:B:326:LYS:HB3	0.52	1.81	8	1
1:A:212:LEU:HD21	2:B:304:PHE:HA	0.51	1.80	20	1
2:B:361:GLN:HA	2:B:364:ARG:HG2	0.51	1.83	7	1
2:B:310:TYR:CE2	2:B:314:ILE:HD11	0.51	2.41	20	12
2:B:313:LYS:HE2	2:B:371:ASP:OD2	0.51	2.06	8	1
2:B:371:ASP:O	2:B:375:GLU:HG2	0.51	2.05	19	2
2:B:327:ALA:O	2:B:331:ILE:HG13	0.51	2.06	3	6
1:A:208:ARG:HD3	1:A:211:GLU:HG3	0.50	1.82	12	2
1:A:220:ARG:HA	1:A:220:ARG:NE	0.50	2.22	1	1
1:A:220:ARG:HA	1:A:220:ARG:HE	0.49	1.67	18	1
1:A:212:LEU:HB2	2:B:304:PHE:HD1	0.49	1.67	6	1
2:B:362:VAL:HG11	2:B:376:PHE:CD2	0.49	2.43	10	1
2:B:360:ALA:O	2:B:364:ARG:HG3	0.49	2.06	18	1
1:A:209:PRO:HA	1:A:212:LEU:HD22	0.49	1.83	3	1
2:B:310:TYR:OH	2:B:376:PHE:HB2	0.48	2.08	12	4
2:B:315:LYS:O	2:B:319:GLN:HB2	0.48	2.08	14	1
2:B:314:ILE:HG12	2:B:372:LEU:HD11	0.48	1.85	13	3
1:A:212:LEU:HD23	1:A:213:LEU:N	0.47	2.24	13	2
2:B:328:PHE:O	2:B:332:LEU:HG	0.47	2.08	19	2
1:A:210:PHE:CD1	2:B:380:LEU:HD11	0.47	2.44	10	1
1:A:220:ARG:NH1	2:B:312:ASN:HA	0.47	2.25	13	1
1:A:212:LEU:HD22	1:A:213:LEU:N	0.47	2.25	1	2
2:B:358:VAL:O	2:B:362:VAL:HG23	0.46	2.10	3	5
1:A:208:ARG:HB2	1:A:211:GLU:CG	0.46	2.37	15	1
2:B:321:GLN:HE21	2:B:321:GLN:N	0.46	2.09	1	1
2:B:353:LEU:HD12	2:B:358:VAL:HG22	0.46	1.88	12	1

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:313:LYS:HE3	2:B:375:GLU:HG3	0.46	1.88	20	1
2:B:357:GLU:O	2:B:361:GLN:HG2	0.45	2.11	13	2
2:B:375:GLU:O	2:B:378:GLN:HG2	0.45	2.12	17	2
2:B:363:ALA:HA	2:B:373:LEU:HD21	0.44	1.89	18	1
2:B:353:LEU:HD11	2:B:361:GLN:OE1	0.44	2.12	7	1
2:B:357:GLU:O	2:B:361:GLN:HG3	0.43	2.13	1	1
2:B:306:HIS:HA	2:B:309:ASN:OD1	0.43	2.13	8	1
2:B:317:ARG:HA	2:B:317:ARG:NE	0.43	2.27	9	2
2:B:318:PHE:CE1	2:B:324:ILE:HG21	0.43	2.49	11	1
2:B:309:ASN:O	2:B:313:LYS:HB2	0.43	2.13	10	1
1:A:210:PHE:O	1:A:214:ILE:HG13	0.43	2.14	3	2
2:B:310:TYR:CE2	2:B:376:PHE:HD1	0.42	2.32	5	1
1:A:209:PRO:HB2	2:B:379:PHE:O	0.42	2.15	9	2
1:A:208:ARG:O	1:A:211:GLU:HB3	0.42	2.15	7	1
2:B:366:PHE:HB3	2:B:369:GLN:HB3	0.42	1.90	17	1
2:B:313:LYS:O	2:B:317:ARG:HG2	0.41	2.15	8	1
2:B:333:HIS:O	2:B:337:LYS:HG3	0.41	2.16	18	1
2:B:321:GLN:HE21	2:B:321:GLN:CA	0.41	2.23	5	1
1:A:208:ARG:CD	1:A:211:GLU:HG3	0.41	2.46	12	1
1:A:208:ARG:HD3	1:A:211:GLU:OE1	0.41	2.15	3	1
2:B:371:ASP:HA	2:B:374:SER:OG	0.41	2.15	4	1
1:A:209:PRO:O	1:A:212:LEU:HG	0.41	2.16	16	2
1:A:221:ASN:HB3	1:A:222:PRO:HD2	0.40	1.94	19	1
2:B:358:VAL:O	2:B:362:VAL:HG22	0.40	2.17	8	1
2:B:354:THR:O	2:B:358:VAL:HG23	0.40	2.16	4	1
2:B:311:VAL:HG13	2:B:325:TYR:OH	0.40	2.16	3	1
1:A:221:ASN:C	1:A:223:THR:H	0.40	2.20	1	1

## 6.3 Torsion angles ⓘ

### 6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	16/45 (36%)	14±1 (88±6%)	1±1 (7±6%)	1±1 (5±5%)	4	25
2	B	72/94 (77%)	70±1 (97±2%)	2±1 (3±2%)	0±0 (0±0%)	59	88

Continued on next page...



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1760/2780 (63%)	1677 (95%)	64 (4%)	19 (1%)	23 69

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	223	THR	13
1	A	209	PRO	4
2	B	319	GLN	1
2	B	367	LYS	1

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	13/39 (33%)	12±1 (90±8%)	1±1 (10±8%)	14 58
2	B	64/81 (79%)	62±1 (97±2%)	2±1 (3±2%)	48 88
All	All	1540/2400 (64%)	1471 (96%)	69 (4%)	38 82

All 26 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	380	LEU	11
1	A	212	LEU	9
1	A	213	LEU	6
1	A	218	MET	6
2	B	354	THR	5
2	B	355	GLU	4
2	B	371	ASP	3
2	B	336	GLN	2
2	B	321	GLN	2
1	A	210	PHE	2
2	B	316	ASN	2
2	B	343	LYS	2
2	B	317	ARG	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
2	B	370	GLU	1
1	A	219	GLU	1
2	B	353	LEU	1
1	A	223	THR	1
2	B	362	VAL	1
2	B	309	ASN	1
2	B	375	GLU	1
2	B	303	GLU	1
2	B	357	GLU	1
2	B	330	GLU	1
2	B	378	GLN	1
2	B	326	LYS	1
1	A	221	ASN	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 84% for the well-defined parts and 84% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2l9s\_cs.str

Chemical shift list name: *pah2pf1.star*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1037
Number of shifts mapped to atoms	1037
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	93	$-0.54 \pm 0.08$	Should be applied
$^{13}\text{C}_\beta$	89	$0.12 \pm 0.04$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	88	$-2.24 \pm 0.13$	Should be applied
$^{15}\text{N}$	88	$-0.08 \pm 0.19$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 69%, i.e. 814 atoms were assigned a chemical shift out of a possible 1177. 0 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	354/432 (82%)	142/172 (83%)	142/176 (81%)	70/84 (83%)
Sidechain	412/634 (65%)	257/373 (69%)	154/225 (68%)	1/36 (3%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	48/111 (43%)	48/59 (81%)	0/48 (0%)	0/4 (0%)
Overall	814/1177 (69%)	447/604 (74%)	296/449 (66%)	71/124 (57%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 57%, i.e. 994 atoms were assigned a chemical shift out of a possible 1748. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	450/677 (66%)	181/269 (67%)	181/278 (65%)	88/130 (68%)
Sidechain	492/935 (53%)	308/551 (56%)	183/332 (55%)	1/52 (2%)
Aromatic	52/136 (38%)	52/72 (72%)	0/60 (0%)	0/4 (0%)
Overall	994/1748 (57%)	541/892 (61%)	364/670 (54%)	89/186 (48%)


#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:



## 7.2 Chemical shift list 2

File name: 2l9s\_cs.str

Chemical shift list name: *pf1pah2.star*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	510
Number of shifts mapped to atoms	510
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

### 7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	44	$-0.09 \pm 0.14$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	42	$-0.01 \pm 0.14$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	36	$-1.86 \pm 0.08$	Should be applied
$^{15}\text{N}$	38	$-0.46 \pm 0.31$	None needed ( $< 0.5$ ppm)

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 15%, i.e. 178 atoms were assigned a chemical shift out of a possible 1177. 0 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	72/432 (17%)	30/172 (17%)	28/176 (16%)	14/84 (17%)
Sidechain	99/634 (16%)	62/373 (17%)	34/225 (15%)	3/36 (8%)
Aromatic	7/111 (6%)	5/59 (8%)	2/48 (4%)	0/4 (0%)
Overall	178/1177 (15%)	97/604 (16%)	64/449 (14%)	17/124 (14%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 27%, i.e. 478 atoms were assigned a chemical shift out of a possible 1748. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	200/677 (30%)	82/269 (30%)	80/278 (29%)	38/130 (29%)

*Continued on next page...*

Continued from previous page...

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Sidechain	258/935 (28%)	161/551 (29%)	88/332 (27%)	9/52 (17%)
Aromatic	20/136 (15%)	14/72 (19%)	6/60 (10%)	0/4 (0%)
Overall	478/1748 (27%)	257/892 (29%)	174/670 (26%)	47/186 (25%)

#### 7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

