wwPDB NMR Structure Validation Summary Report (i)

Apr 27, 2016 – 01:06 AM BST

PDB ID	:	2LAV
Title	:	NMR solution structure of human Vaccinia-Related Kinase 1
Authors	:	Shin, J.; Yoon, H.S.
Deposited on	:	2011-03-21

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 72%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	361	50% 34% • 14%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:23-A:44, A:49-A:61, A:66-A:334, A:351-A:356 (310)	0.58	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 12, 14, 17, 19
2	1, 13, 15, 18
3	5, 7, 10
4	3, 9
5	4, 6
Single-model clusters	8; 11; 16; 20

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 5859 atoms, of which 2949 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Vaccinia-related kinase 1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	361	5859	1853	2949	509	534	14	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Vaccinia-related kinase 1

Chain A: 50% 34% • 14%



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

- Molecule 1: Vaccinia-related kinase 1

Chain A: 51% 31% • 14%




E262
L265
K266
D267
V271
R272
D273
Y278
R279
I282
L285
M286
D287
K288
M303
E304
T305
V306
K307
L308
L309
D310
Y311
D322
I323
L324
L325
Q326
G327
L328
K329
D335
D336
G337
K338
L339
D340
L341
S342
V343
V344
E345
N346
G347
G348
L349
K350
T353
T354
T355
K356
K357
R358
K359
K360
E361

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	1.2

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2lav_cs.str
Number of chemical shift lists	1
Total number of shifts	3545
Number of shifts mapped to atoms	3545
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	72%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	2527	2553	2545	71±6
All	All	50540	51060	50900	1424

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

5 of 703 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:183:LEU:HD21	1:A:195:LEU:HD12	0.98	1.35	8	2
1:A:53:LEU:HD22	1:A:66:ALA:HB3	0.92	1.37	14	10
1:A:29:ILE:HD11	1:A:37:TRP:CZ2	0.89	2.03	2	2
1:A:201:ALA:HB3	1:A:353:THR:HG22	0.87	1.45	4	17
1:A:53:LEU:HD23	1:A:66:ALA:HB3	0.85	1.48	5	4

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	310/361 (86%)	273±4 (88±1%)	27±4 (9±1%)	10±2 (3±1%)	8 38
All	All	6200/7220 (86%)	5451 (88%)	541 (9%)	208 (3%)	8 38

5 of 33 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	354	ILE	20
1	A	353	THR	20
1	A	213	TYR	18
1	A	200	LEU	18
1	A	197	ASP	14

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	273/312 (88%)	222±6 (81±2%)	51±6 (19±2%)	5 38
All	All	5460/6240 (88%)	4443 (81%)	1017 (19%)	5 38

5 of 184 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	355	THR	20
1	A	202	TYR	20
1	A	273	ASP	18
1	A	310	ASP	18
1	A	82	THR	18

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 72% for the well-defined parts and 69% for the entire structure.

7.1 Chemical shift list 1

File name: 2lav_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	3545
Number of shifts mapped to atoms	3545
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	9

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	354	0.10 ± 0.05	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	317	0.61 ± 0.07	Should be applied
$^{13}C'$	343	0.18 ± 0.06	None needed (< 0.5 ppm)
^{15}N	327	-0.41 ± 0.16	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 72%, i.e. 2875 atoms were assigned a chemical shift out of a possible 4005. 34 out of 45 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	1481/1520 (97%)	578/605 (96%)	614/620 (99%)	289/295 (98%)
Sidechain	1278/2129 (60%)	860/1258 (68%)	413/770 (54%)	5/101 (5%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	116/356 (33%)	111/182 (61%)	0/153 (0%)	5/21 (24%)
Overall	2875/4005 (72%)	1549/2045 (76%)	1027/1543 (67%)	299/417 (72%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	29	ILE	HG23	-0.80	2.13 – -0.57	-5.8
1	A	29	ILE	HG22	-0.80	2.13 – -0.57	-5.8
1	A	29	ILE	HG21	-0.80	2.13 – -0.57	-5.8
1	A	29	ILE	HD11	-0.94	2.13 – -0.77	-5.6
1	A	29	ILE	HD12	-0.94	2.13 – -0.77	-5.6
1	A	29	ILE	HD13	-0.94	2.13 – -0.77	-5.6
1	A	130	ILE	HG22	-0.62	2.13 – -0.57	-5.2
1	A	130	ILE	HG23	-0.62	2.13 – -0.57	-5.2
1	A	130	ILE	HG21	-0.62	2.13 – -0.57	-5.2

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: