



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 03:50 PM BST

PDB ID : 1LFU  
Title : NMR Solution Structure of the Extended PBX Homeodomain Bound to DNA  
Authors : Sprules, T.; Green, N.; Featherstone, M.; Gehring, K.  
Deposited on : 2002-04-12

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 51%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | NMR archive<br>(#Entries) |
|-----------------------|-----------------------------|---------------------------|
| Clashscore            | 114402                      | 11133                     |
| Ramachandran outliers | 111179                      | 9975                      |
| Sidechain outliers    | 111093                      | 9958                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1   | A     | 14     | 100%             |
| 2   | B     | 14     | 100%             |
| 3   | P     | 82     | 21% 52% 10% 17%  |

## 2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

| Well-defined (core) protein residues |                       |                   |              |
|--------------------------------------|-----------------------|-------------------|--------------|
| Well-defined core                    | Residue range (total) | Backbone RMSD (Å) | Medoid model |
| 1                                    | P:6-P:70 (65)         | 0.22              | 10           |

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

| Cluster number | Models                                 |
|----------------|--|
| 1              | 2, 4, 6, 8, 10, 12, 13, 15, 17, 19, 20 |
| 2              | 1, 3, 5, 7, 9, 14, 16                  |
| 3              | 11, 18                                 |

### 3 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2239 atoms, of which 995 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(\*GP\*CP\*GP\*CP\*AP\*TP\*GP\*AP\*TP\*TP\*GP\*CP\*CP\*C)-3'.

| Mol | Chain | Residues | Atoms |     |     |    |    |    | Trace |
|-----|-------|----------|-------|-----|-----|----|----|----|-------|
| 1   | A     | 14       | Total | C   | H   | N  | O  | P  | 0     |
|     |       |          | 441   | 135 | 159 | 51 | 83 | 13 |       |

- Molecule 2 is a DNA chain called 5'-D(\*GP\*GP\*GP\*CP\*AP\*AP\*TP\*CP\*AP\*TP\*GP\*CP\*GP\*C)-3'.

| Mol | Chain | Residues | Atoms |     |     |    |    |    | Trace |
|-----|-------|----------|-------|-----|-----|----|----|----|-------|
| 2   | B     | 14       | Total | C   | H   | N  | O  | P  | 0     |
|     |       |          | 444   | 136 | 158 | 56 | 81 | 13 |       |

- Molecule 3 is a protein called homeobox protein PBX1.

| Mol | Chain | Residues | Atoms |     |     |     |     |   | Trace |
|-----|-------|----------|-------|-----|-----|-----|-----|---|-------|
| 3   | P     | 82       | Total | C   | H   | N   | O   | S | 0     |
|     |       |          | 1354  | 426 | 678 | 124 | 125 | 1 |       |

There are 2 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment               | Reference  |
|-------|---------|----------|--------|-----------------------|------------|
| P     | 0       | MET      | -      | INITIATING METHIONINE | UNP P41778 |
| P     | 38      | SER      | CYS    | ENGINEERED            | UNP P41778 |

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-D(\*GP\*CP\*GP\*CP\*AP\*TP\*GP\*AP\*TP\*TP\*GP\*CP\*CP\*C)-3'

Chain A:  100%

G1 C2 C3 C4 A5 T6 G7 A8 T9 T10 G11 C12 C13 C14

- Molecule 2: 5'-D(\*GP\*GP\*GP\*CP\*AP\*AP\*TP\*CP\*AP\*TP\*GP\*CP\*GP\*C)-3'

Chain B:  100%

G15 G16 G17 G18 C19 A20 T21 T22 C23 T24 G25 C26 G27 C28

- Molecule 3: homeobox protein PBX1

Chain P:  21% 52% 10% 17%

W0 A1 R2 R3 K4 R5 R6 R7 F8 R9 K10 Q11 A12 T13 T14 E15 I16 L17 N18 E19 Y20 Y21 Y22 S23 H24 L25 S26 N27 P28 E29 A30 K31 E32 E33 L34 A35 K36 K37 S38 G39 I40 T41 V42 S43 Q44 V45 S46 N47 W48 F49 K52 R53 I54 R55 Y56 K57

K58 N59 I60 G61 K62 F63 E66 A67 I68 I69 A70 Y71 A72 A73 T74 V75 V76 T77 A78

### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 10. Colouring as in section 4.1 above.

- Molecule 1: 5'-D(\*GP\*CP\*GP\*CP\*AP\*TP\*GP\*AP\*TP\*TP\*GP\*CP\*CP\*C)-3'


Chain A:  7% 93%

G1 C2 C3 C4 A5 T6 G7 A8 T9 T10 G11 C12 C13 C14

● Molecule 2: 5'-D(\*GP\*GP\*GP\*CP\*AP\*AP\*TP\*CP\*AP\*TP\*GP\*CP\*GP\*C)-3'

Chain B: 

100%



● Molecule 3: homeobox protein PBX1


Chain P: 

23%

46%

13%

17%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *dynamical annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *lowest energy with acceptable geometry*.

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version  |
|---------------|----------------|----------|
| CNS           | refinement     | 0.9, 1.1 |

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

|  |                 |
|--|-----------------|
| Chemical shift file(s)                       | BMRB entry 5349 |
| Number of chemical shift lists               | 2               |
| Total number of shifts                       | 1184            |
| Number of shifts mapped to atoms             | 941             |
| Number of unparsed shifts                    | 0               |
| Number of shifts with mapping errors         | 243             |
| Number of shifts with mapping warnings       | 0               |
| Assignment completeness (well-defined parts) | 51%             |

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | Chirality | Planarity |
|-----|-------|-----------|-----------|
| 2   | B     | 0.0±0.0   | 0.1±0.2   |
| All | All   | 0         | 1         |

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

| Mol | Chain | Res | Type | Group     | Models (Total) |
|-----|-------|-----|------|-----------|----------------|
| 2   | B     | 24  | DT   | Sidechain | 1              |

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|-----|-------|-------|----------|----------|---------|
| 1   | A     | 282   | 159      | 159      | 38±4    |
| 2   | B     | 286   | 158      | 158      | 52±5    |
| 3   | P     | 570   | 554      | 554      | 68±8    |
| All | All   | 22760 | 17420    | 17420    | 2972    |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 74.

5 of 613 unique clashes are listed below, sorted by their clash magnitude.



| Atom-1        | Atom-2          | Clash(Å) | Distance(Å) | Models |       |
|---------------|-----------------|----------|-------------|--------|-------|
|               |                 |          |             | Worst  | Total |
| 1:A:7:DG:H2'' | 1:A:8:DA:O5'    | 1.01     | 1.54        | 9      | 13    |
| 1:A:9:DT:C6   | 1:A:10:DT:H72   | 1.00     | 1.92        | 9      | 20    |
| 1:A:7:DG:H2'' | 1:A:8:DA:C8     | 0.96     | 1.94        | 19     | 19    |
| 2:B:18:DC:H1' | 2:B:19:DA:O4'   | 0.92     | 1.62        | 4      | 1     |
| 3:P:56:TYR:O  | 3:P:60:ILE:HD13 | 0.92     | 1.65        | 11     | 2     |

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

| Mol | Chain | Analysed        | Favoured     | Allowed     | Outliers   | Percentiles |    |
|-----|-------|-----------------|--------------|-------------|------------|-------------|----|
| 3   | P     | 68/82 (83%)     | 54±2 (79±2%) | 9±2 (12±2%) | 6±1 (9±1%) | 2           | 13 |
| All | All   | 1360/1640 (83%) | 1070 (79%)   | 170 (12%)   | 120 (9%)   | 2           | 13 |

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 3   | P     | 60  | ILE  | 20             |
| 3   | P     | 42  | VAL  | 20             |
| 3   | P     | 7   | ASN  | 19             |
| 3   | P     | 6   | ARG  | 18             |
| 3   | P     | 21  | TYR  | 16             |

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

| Mol | Chain | Analysed        | Rotameric    | Outliers   | Percentiles |    |
|-----|-------|-----------------|--------------|------------|-------------|----|
| 3   | P     | 61/70 (87%)     | 55±1 (91±2%) | 6±1 (9±2%) | 16          | 61 |
| All | All   | 1220/1400 (87%) | 1109 (91%)   | 111 (9%)   | 16          | 61 |

5 of 19 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 3   | P     | 36  | LYS  | 20             |
| 3   | P     | 47  | ASN  | 18             |
| 3   | P     | 31  | LYS  | 17             |
| 3   | P     | 58  | LYS  | 13             |
| 3   | P     | 10  | LYS  | 7              |

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 51% for the well-defined parts and 53% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 5349

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

|   |     |
|---|-----|
| Total number of shifts                  | 941 |
| Number of shifts mapped to atoms        | 941 |
| Number of unparsed shifts               | 0   |
| Number of shifts with mapping errors    | 0   |
| Number of shifts with mapping warnings  | 0   |
| Number of shift outliers (ShiftChecker) | 6   |

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

| Nucleus                | # values | Correction $\pm$ precision, ppm | Suggested action           |
|------------------------|----------|---------------------------------|----------------------------|
| $^{13}\text{C}_\alpha$ | 82       | $-0.62 \pm 0.10$                | Should be applied          |
| $^{13}\text{C}_\beta$  | 77       | $0.10 \pm 0.07$                 | None needed ( $< 0.5$ ppm) |
| $^{13}\text{C}'$       | 0        | —                               | —                          |
| $^{15}\text{N}$        | 78       | $-0.62 \pm 0.25$                | Should be applied          |

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 51%, i.e. 742 atoms were assigned a chemical shift out of a possible 1459. 2 out of 5 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

|           | Total         | $^1\text{H}$  | $^{13}\text{C}$ | $^{15}\text{N}$ |
|-----------|---------------|---------------|-----------------|-----------------|
| Backbone  | 266/336 (79%) | 133/134 (99%) | 68/136 (50%)    | 65/66 (98%)     |
| Sidechain | 385/469 (82%) | 246/279 (88%) | 129/162 (80%)   | 10/28 (36%)     |

*Continued on next page...*

Continued from previous page...

|          | Total          | <sup>1</sup> H | <sup>13</sup> C | <sup>15</sup> N |
|----------|----------------|----------------|-----------------|-----------------|
| Aromatic | 91/95 (96%)    | 47/50 (94%)    | 43/43 (100%)    | 1/2 (50%)       |
| Overall  | 742/1459 (51%) | 426/798 (53%)  | 240/528 (45%)   | 76/133 (57%)    |

### 7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

| Mol | Chain | Res | Type | Atom | Shift, ppm | Expected range, ppm | Z-score |
|-----|-------|-----|------|------|------------|---------------------|---------|
| ??? | P     | 52  | LYS  | HG3  | -0.85      | 2.76 – -0.04        | -7.9    |
| ??? | P     | 16  | LEU  | HB3  | -1.03      | 3.34 – -0.26        | -7.1    |
| ??? | P     | 52  | LYS  | HG2  | -0.42      | 2.67 – 0.07         | -6.9    |
| ??? | P     | 52  | LYS  | HB3  | -0.06      | 3.10 – 0.40         | -6.7    |
| ??? | P     | 52  | LYS  | HD3  | 0.12       | 2.75 – 0.45         | -6.4    |
| ??? | P     | 26  | PRO  | HG3  | 0.12       | 3.56 – 0.26         | -5.4    |

### 7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain P:



## 7.2 Chemical shift list 2

File name: BMRB entry 5349

Chemical shift list name: *assigned\_chem\_shift\_list\_2*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

|   |     |
|---|-----|
| Total number of shifts                  | 243 |
| Number of shifts mapped to atoms        | 0   |
| Number of unparsed shifts               | 0   |
| Number of shifts with mapping errors    | 243 |
| Number of shifts with mapping warnings  | 0   |
| Number of shift outliers (ShiftChecker) | 0   |

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. First 5 (of 243) occurrences are reported below.

| Chain    | Res | Type | Atom | Shift Data |             |           |
|----------|-----|------|------|------------|-------------|-----------|
|          |     |      |      | Value      | Uncertainty | Ambiguity |
| UNMAPPED | 5   | DA   | H3'  | 5.0        | 0.01        | 1         |
| UNMAPPED | 28  | DC   | H3'  | 4.52       | 0.01        | 1         |
| UNMAPPED | 13  | DC   | H6   | 7.56       | 0.01        | 1         |
| UNMAPPED | 6   | DT   | H1'  | 5.25       | 0.01        | 1         |
| UNMAPPED | 24  | DT   | H71  | 1.4        | 0.2         | 1         |

### 7.2.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 1459. 0 out of 5 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

|           | Total       | <sup>1</sup> H | <sup>13</sup> C | <sup>15</sup> N |
|-----------|-------------|----------------|-----------------|-----------------|
| Backbone  | 0/336 (0%)  | 0/134 (0%)     | 0/136 (0%)      | 0/66 (0%)       |
| Sidechain | 0/469 (0%)  | 0/279 (0%)     | 0/162 (0%)      | 0/28 (0%)       |
| Aromatic  | 0/95 (0%)   | 0/50 (0%)      | 0/43 (0%)       | 0/2 (0%)        |
| Overall   | 0/1459 (0%) | 0/798 (0%)     | 0/528 (0%)      | 0/133 (0%)      |

#### 7.2.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

#### 7.2.5 Random Coil Index (RCI) plots ⓘ

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned\_chem\_shift\_list\_2). RCI is only applicable to proteins.