



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 06:23 PM GMT

PDB ID : 4LFD
Title : Staphylococcus aureus sortase B-substrate complex
Authors : Jacobitz, A.W.; Sawaya, M.R.; Yi, S.W.; Amer, B.R.; Huang, G.L.; Nguyen, A.V.; Jung, M.E.; Clubb, R.T.
Deposited on : 2013-06-26
Resolution : 2.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.




Continued on next page...

Continued from previous page...



- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total O 4 4	0	0
4	B	9	Total O 9 9	0	0
4	F	1	Total O 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	7	Total O 7 7	0	0
4	D	6	Total O 6 6	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sortase B




- Molecule 1: Sortase B



- Molecule 1: Sortase B




- Molecule 1: Sortase B





- Molecule 2: (CBZ)NPQ(B27) PEPTIDE

Chain E:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: (CBZ)NPQ(B27) PEPTIDE

Chain F:  20% 80% 20%



- Molecule 2: (CBZ)NPQ(B27) PEPTIDE

Chain G:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: (CBZ)NPQ(B27) PEPTIDE

Chain H:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	102.60 Å 59.12 Å 72.49 Å 90.00° 90.02° 90.00°	Depositor
Resolution (Å)	72.49 – 2.49 59.21 – 2.49	Depositor EDS
% Data completeness (in resolution range)	91.4 (72.49-2.49) 91.4 (59.21-2.49)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.27 (at 2.48 Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R , R_{free}	0.215 , 0.271 0.257 , 0.279	Depositor DCC
R_{free} test set	1402 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	42.0	Xtriage
Anisotropy	0.820	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.4	EDS
Estimated twinning fraction	0.437 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	4 of 28147 reflections (0.014%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7451	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.39 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5368e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: B27, PHQ, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	0/1846	0.73	0/2478
1	B	0.50	0/1846	0.78	0/2478
1	C	0.52	0/1846	0.78	0/2478
1	D	0.50	0/1846	0.79	0/2478
2	E	0.59	0/24	0.73	0/32
2	F	0.55	0/24	0.61	0/32
2	G	0.52	0/24	0.69	0/32
2	H	0.50	0/24	0.72	0/32
All	All	0.49	0/7480	0.77	0/10040

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbit. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1812	0	1776	26	0
1	B	1812	0	1776	23	0
1	C	1812	0	1776	31	0
1	D	1812	0	1776	26	0
2	E	34	0	28	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	34	0	28	1	0
2	G	34	0	28	0	0
2	H	34	0	28	0	0
3	A	5	0	0	0	0
3	B	15	0	0	0	0
3	C	10	0	0	0	0
3	D	10	0	0	0	0
4	A	4	0	0	0	0
4	B	9	0	0	0	0
4	C	7	0	0	0	0
4	D	6	0	0	0	0
4	F	1	0	0	1	0
All	All	7451	0	7216	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:GLN:HE22	1:A:206:ASN:H	1.13	0.96
1:C:147:GLN:HE22	1:C:206:ASN:H	1.21	0.88
1:D:147:GLN:HE22	1:D:206:ASN:H	1.20	0.87
1:A:58:ARG:HH12	1:C:229:GLU:HG3	1.41	0.84
1:B:147:GLN:HE22	1:B:206:ASN:H	1.23	0.83
1:A:160:ASN:HD22	1:A:162:TYR:H	1.29	0.80
1:D:116:ASN:HD21	1:D:126:ILE:H	1.29	0.80
1:D:160:ASN:HD22	1:D:162:TYR:H	1.42	0.66
1:C:230:THR:HG23	1:C:232:LYS:HD3	1.78	0.66
1:D:116:ASN:ND2	1:D:126:ILE:H	1.93	0.65
1:C:116:ASN:HD21	1:C:126:ILE:H	1.43	0.64
1:C:124:ASN:OD1	1:C:219:THR:HG23	1.97	0.64
1:A:50:MET:SD	1:A:60:GLN:OE1	2.56	0.64
1:B:160:ASN:HD22	1:B:162:TYR:H	1.44	0.63
1:B:211:VAL:HG11	1:B:217:ILE:HD11	1.81	0.62
1:A:211:VAL:HG11	1:A:217:ILE:HD11	1.81	0.62
2:F:301:ASN:HB2	4:F:401:HOH:O	2.00	0.61
1:A:217:ILE:HG13	1:A:238:ALA:O	2.00	0.61
1:A:95:TYR:OH	1:A:102:ARG:HD3	2.00	0.61
1:B:116:ASN:HD21	1:B:126:ILE:H	1.48	0.61
1:C:116:ASN:ND2	1:C:126:ILE:H	1.99	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:LEU:HB2	1:B:236:VAL:HG22	1.81	0.61
1:A:116:ASN:ND2	1:A:126:ILE:H	1.99	0.60
1:C:211:VAL:HG11	1:C:217:ILE:HD11	1.83	0.60
1:B:155:ILE:CD1	1:B:166:GLN:HB3	2.32	0.60
1:A:147:GLN:NE2	1:A:206:ASN:H	1.92	0.59
1:B:155:ILE:HD13	1:B:166:GLN:HB3	1.84	0.59
1:A:116:ASN:HD21	1:A:126:ILE:H	1.51	0.59
1:D:155:ILE:HD13	1:D:166:GLN:HB3	1.83	0.59
1:D:155:ILE:CD1	1:D:166:GLN:HB3	2.34	0.58
1:B:116:ASN:ND2	1:B:126:ILE:H	2.01	0.58
1:D:53:HIS:HB2	1:D:60:GLN:HE21	1.69	0.58
1:D:106:ARG:HH12	1:D:130:HIS:CD2	2.21	0.58
1:D:211:VAL:HG11	1:D:217:ILE:HD11	1.85	0.57
1:C:219:THR:HG22	1:C:237:VAL:HG13	1.86	0.57
1:B:106:ARG:HH12	1:B:130:HIS:CD2	2.22	0.56
1:A:30:MET:O	1:A:34:GLN:HB2	2.05	0.56
1:D:30:MET:O	1:D:34:GLN:HB2	2.06	0.56
1:A:230:THR:HG23	1:A:232:LYS:HD3	1.88	0.55
1:C:30:MET:O	1:C:34:GLN:HB2	2.07	0.55
1:D:220:LEU:HB2	1:D:236:VAL:HG22	1.88	0.55
1:B:30:MET:O	1:B:34:GLN:HB2	2.08	0.54
1:C:128:TYR:HE1	1:C:182:ILE:HD11	1.73	0.53
1:B:115:ARG:HH22	1:B:182:ILE:HG23	1.74	0.53
1:A:146:LYS:N	1:A:146:LYS:HD2	2.24	0.53
1:C:53:HIS:HB2	1:C:60:GLN:HE21	1.74	0.52
1:A:147:GLN:HE22	1:A:206:ASN:N	1.96	0.52
1:C:116:ASN:HA	1:C:122:ASN:ND2	2.25	0.51
1:C:115:ARG:HH22	1:C:182:ILE:HG23	1.76	0.51
1:C:95:TYR:OH	1:C:102:ARG:HD3	2.11	0.51
1:B:120:ASN:ND2	1:C:147:GLN:HG2	2.26	0.51
1:C:243:VAL:HG12	1:D:243:VAL:HG12	1.92	0.50
1:C:146:LYS:C	1:C:205:ILE:HD11	2.33	0.49
1:A:239:LYS:HE3	1:A:241:ILE:HD11	1.94	0.49
1:A:115:ARG:HH22	1:A:182:ILE:HG23	1.77	0.49
1:B:116:ASN:HA	1:B:122:ASN:ND2	2.27	0.49
1:C:160:ASN:HD22	1:C:162:TYR:H	1.60	0.49
1:D:170:PHE:HE1	1:D:217:ILE:HD12	1.78	0.49
1:A:116:ASN:HA	1:A:122:ASN:ND2	2.28	0.48
1:C:181:TYR:HE2	1:C:221:SER:HB2	1.78	0.48
1:C:147:GLN:NE2	1:C:205:ILE:HG13	2.29	0.48
1:A:146:LYS:C	1:A:205:ILE:HD11	2.35	0.47

Continued on next page...

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	214/216 (99%)	206 (96%)	8 (4%)	0	100 100
1	B	214/216 (99%)	205 (96%)	8 (4%)	1 (0%)	34 55
1	C	214/216 (99%)	202 (94%)	11 (5%)	1 (0%)	34 55
1	D	214/216 (99%)	202 (94%)	11 (5%)	1 (0%)	34 55
2	E	2/5 (40%)	2 (100%)	0	0	100 100
2	F	2/5 (40%)	2 (100%)	0	0	100 100
2	G	2/5 (40%)	2 (100%)	0	0	100 100
2	H	2/5 (40%)	2 (100%)	0	0	100 100
All	All	864/884 (98%)	823 (95%)	38 (4%)	3 (0%)	46 68

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	32	ASP
1	C	32	ASP
1	D	32	ASP

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	204/205 (100%)	184 (90%)	20 (10%)	10 19
1	B	204/205 (100%)	180 (88%)	24 (12%)	6 12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	204/205 (100%)	177 (87%)	27 (13%)	5 9
1	D	204/205 (100%)	186 (91%)	18 (9%)	12 23
2	E	3/3 (100%)	3 (100%)	0	100 100
2	F	3/3 (100%)	2 (67%)	1 (33%)	0 0
2	G	3/3 (100%)	3 (100%)	0	100 100
2	H	3/3 (100%)	3 (100%)	0	100 100
All	All	828/832 (100%)	738 (89%)	90 (11%)	8 15

All (90) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	LYS
1	A	34	GLN
1	A	42	LEU
1	A	64	LEU
1	A	94	ASP
1	A	106	ARG
1	A	107	LYS
1	A	123	HIS
1	A	131	HIS
1	A	132	VAL
1	A	160	ASN
1	A	167	LEU
1	A	182	ILE
1	A	184	THR
1	A	205	ILE
1	A	209	VAL
1	A	210	ASN
1	A	217	ILE
1	A	224	GLU
1	A	232	LYS
1	B	34	GLN
1	B	64	LEU
1	B	72	VAL
1	B	87	LEU
1	B	90	LYS
1	B	94	ASP
1	B	106	ARG
1	B	107	LYS
1	B	123	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	126	ILE
1	B	131	HIS
1	B	132	VAL
1	B	140	VAL
1	B	160	ASN
1	B	167	LEU
1	B	182	ILE
1	B	184	THR
1	B	209	VAL
1	B	210	ASN
1	B	217	ILE
1	B	223	CYS
1	B	224	GLU
1	B	230	THR
1	B	232	LYS
2	F	301	ASN
1	C	34	GLN
1	C	38	ASN
1	C	42	LEU
1	C	64	LEU
1	C	87	LEU
1	C	90	LYS
1	C	91	THR
1	C	106	ARG
1	C	107	LYS
1	C	109	SER
1	C	123	HIS
1	C	126	ILE
1	C	131	HIS
1	C	132	VAL
1	C	140	VAL
1	C	167	LEU
1	C	182	ILE
1	C	184	THR
1	C	205	ILE
1	C	209	VAL
1	C	210	ASN
1	C	217	ILE
1	C	219	THR
1	C	223	CYS
1	C	224	GLU
1	C	229	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	232	LYS
1	D	34	GLN
1	D	64	LEU
1	D	90	LYS
1	D	106	ARG
1	D	107	LYS
1	D	123	HIS
1	D	126	ILE
1	D	131	HIS
1	D	132	VAL
1	D	140	VAL
1	D	160	ASN
1	D	167	LEU
1	D	184	THR
1	D	209	VAL
1	D	210	ASN
1	D	224	GLU
1	D	230	THR
1	D	232	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (33) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	60	GLN
1	A	116	ASN
1	A	122	ASN
1	A	147	GLN
1	A	153	HIS
1	A	160	ASN
1	A	194	GLN
1	B	60	GLN
1	B	116	ASN
1	B	122	ASN
1	B	130	HIS
1	B	147	GLN
1	B	153	HIS
1	B	160	ASN
1	B	194	GLN
1	C	38	ASN
1	C	44	GLN
1	C	60	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	116	ASN
1	C	122	ASN
1	C	135	ASN
1	C	147	GLN
1	C	153	HIS
1	C	160	ASN
1	C	193	GLN
1	D	60	GLN
1	D	116	ASN
1	D	122	ASN
1	D	147	GLN
1	D	153	HIS
1	D	160	ASN
1	D	194	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	B27	E	304	1,2	5,6,6	0.41	0	2,7,7	0.97	0
2	B27	F	304	1,2	5,6,6	0.34	0	2,7,7	0.87	0
2	B27	G	304	1,2	5,6,6	0.38	0	2,7,7	0.95	0
2	B27	H	304	1,2	5,6,6	0.56	0	2,7,7	1.64	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B27	E	304	1,2	-	0/4/6/6	0/0/0/0
2	B27	F	304	1,2	-	0/4/6/6	0/0/0/0
2	B27	G	304	1,2	-	0/4/6/6	0/0/0/0
2	B27	H	304	1,2	-	0/4/6/6	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	H	304	B27	CA-CX-SG	-2.23	109.47	114.51

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	301	-	4,4,4	0.59	0	6,6,6	0.41	0
3	SO4	B	301	-	4,4,4	0.21	0	6,6,6	0.20	0
3	SO4	B	302	-	4,4,4	0.11	0	6,6,6	0.23	0
3	SO4	B	303	-	4,4,4	0.66	0	6,6,6	0.39	0
3	SO4	C	301	-	4,4,4	0.68	0	6,6,6	0.34	0
3	SO4	C	302	-	4,4,4	0.14	0	6,6,6	0.19	0
3	SO4	D	301	-	4,4,4	0.62	0	6,6,6	0.63	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	D	302	-	4,4,4	0.28	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	301	-	-	0/0/0/0	0/0/0/0
3	SO4	B	301	-	-	0/0/0/0	0/0/0/0
3	SO4	B	302	-	-	0/0/0/0	0/0/0/0
3	SO4	B	303	-	-	0/0/0/0	0/0/0/0
3	SO4	C	301	-	-	0/0/0/0	0/0/0/0
3	SO4	C	302	-	-	0/0/0/0	0/0/0/0
3	SO4	D	301	-	-	0/0/0/0	0/0/0/0
3	SO4	D	302	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	216/216 (100%)	0.36	12 (5%)	28	31	52, 72, 101, 140	0
1	B	216/216 (100%)	0.17	10 (4%)	36	41	50, 68, 95, 111	0
1	C	216/216 (100%)	0.20	10 (4%)	36	41	45, 67, 98, 123	0
1	D	216/216 (100%)	0.25	14 (6%)	22	25	49, 70, 98, 123	0
2	E	3/5 (60%)	0.38	0	100	100	80, 80, 91, 98	0
2	F	3/5 (60%)	1.40	1 (33%)	0	0	82, 82, 92, 106	0
2	G	3/5 (60%)	0.40	0	100	100	76, 76, 88, 100	0
2	H	3/5 (60%)	0.28	0	100	100	87, 87, 89, 96	0
All	All	876/884 (99%)	0.25	47 (5%)	29	33	45, 70, 98, 140	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	32	ASP	12.2
1	A	32	ASP	8.7
1	D	32	ASP	7.2
1	B	32	ASP	7.1
1	A	31	GLU	6.2
1	C	31	GLU	5.5
1	D	31	GLU	4.8
1	C	225	ASP	4.7
1	D	54	GLN	4.5
1	B	31	GLU	4.0
1	A	244	SER	3.7
2	F	301	ASN	3.6
1	D	244	SER	3.5
1	B	106	ARG	3.4
1	D	55	GLU	3.2
1	B	225	ASP	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	46	PHE	3.2
1	A	30	MET	3.2
1	D	63	SER	3.1
1	B	244	SER	3.0
1	A	101	GLU	3.0
1	A	33	LYS	3.0
1	D	50	MET	3.0
1	D	51	SER	2.9
1	A	225	ASP	2.7
1	C	51	SER	2.7
1	B	54	GLN	2.6
1	C	106	ARG	2.5
1	C	101	GLU	2.4
1	B	197	ASP	2.4
1	C	48	MET	2.4
1	D	66	LYS	2.3
1	C	30	MET	2.3
1	D	48	MET	2.3
1	A	36	ARG	2.3
1	A	50	MET	2.3
1	D	30	MET	2.2
1	C	97	ASN	2.2
1	C	244	SER	2.2
1	D	225	ASP	2.2
1	D	177	THR	2.2
1	A	177	THR	2.2
1	B	48	MET	2.2
1	B	132	VAL	2.2
1	B	93	HIS	2.1
1	D	210	ASN	2.1
1	A	97	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	B27	G	304	7/7	0.92	0.18	-	82,84,90,91	0
2	B27	E	304	7/7	0.91	0.19	-	92,93,97,98	0
2	B27	H	304	7/7	0.92	0.13	-	89,90,92,94	0
2	B27	F	304	7/7	0.94	0.12	-	81,81,86,92	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SO4	D	302	5/5	0.84	0.22	1.41	137,137,138,139	0
3	SO4	B	302	5/5	0.86	0.17	0.49	111,114,114,115	0
3	SO4	C	302	5/5	0.93	0.13	-0.57	100,101,101,102	0
3	SO4	B	301	5/5	0.95	0.13	-0.64	102,102,103,104	0
3	SO4	D	301	5/5	0.95	0.15	-0.74	44,46,47,48	0
3	SO4	B	303	5/5	0.95	0.15	-0.89	45,50,51,55	0
3	SO4	C	301	5/5	0.98	0.12	-1.28	32,34,37,38	0
3	SO4	A	301	5/5	0.92	0.13	-1.42	55,58,59,61	0

6.5 Other polymers

There are no such residues in this entry.