



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2016 – 05:33 PM EDT

PDB ID : 5LIH
Title : Structure of a peptide-substrate bound to PKC α core kinase domain
Authors : Soriano, E.V.; Purkiss, A.G.; McDonald, N.Q.
Deposited on : 2016-07-14
Resolution : 3.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1624 (3.32-3.20)
Clashscore	102246	1806 (3.32-3.20)
Ramachandran outliers	100387	1773 (3.32-3.20)
Sidechain outliers	100360	1771 (3.32-3.20)
RSRZ outliers	91569	1632 (3.32-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div>2%</div> <div> <div></div> <div>73%</div> <div>21%</div> <div>7%</div> </div> </div>
1	B	349	<div> <div>2%</div> <div> <div></div> <div>71%</div> <div>21%</div> <div>8%</div> </div> </div>
2	F	16	<div> <div></div> <div> <div></div> <div>75%</div> <div>6%</div> <div>19%</div> </div> </div>
2	G	16	<div> <div></div> <div> <div></div> <div>56%</div> <div>6%</div> <div>38%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	AF3	A	602	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5298 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein kinase C iota type.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	P	S	0	1	0
			2527	1636	412	464	2	13			
1	B	320	Total	C	N	O	P	S	0	0	0
			2489	1610	403	461	2	13			

- Molecule 2 is a protein called PKC Epsilon pseudo substrate sequence.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	13	Total	C	N	O	S	0	0	0
			100	61	23	15	1			
2	G	10	Total	C	N	O		0	0	0
			76	47	17	12				

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	
			27	10	5	10	2	
3	B	1	Total	C	N	O	P	
			27	10	5	10	2	

- Molecule 4 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula: AlF_3).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Al	F		
			4	1	3	0	0
4	A	1	Total	Al	F		
			4	1	3	0	0
4	B	1	Total	Al	F		
			4	1	3	0	0
4	B	1	Total	Al	F		
			4	1	3	0	0

- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Mn		
			2	2	0	0
5	A	3	Total	Mn		
			3	3	0	0

- Molecule 6 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	S	0	0
			3	1	1	1		
6	B	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	13	Total	O	0	0
			13	13		
7	B	10	Total	O	0	0
			10	10		
7	F	2	Total	O	0	0
			2	2		

GLU	
ARG	
MET	
ARG	
PS	
R13	
R14	
ARG	
VAL	

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.98Å 84.23Å 111.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.28 – 3.25 67.28 – 3.25	Depositor EDS
% Data completeness (in resolution range)	99.1 (67.28-3.25) 92.4 (67.28-3.25)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 3.26Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, R_{free}	0.257 , 0.284 0.253 , 0.282	Depositor DCC
R_{free} test set	593 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	43.3	Xtriage
Anisotropy	0.662	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	5298	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.02 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.5117e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, SCN, AF3, ADP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/2569	0.47	0/3481
1	B	0.26	0/2526	0.45	0/3419
2	F	0.24	0/101	0.38	0/133
2	G	0.24	0/77	0.38	0/101
All	All	0.26	0/5273	0.46	0/7134

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2527	0	2343	40	0
1	B	2489	0	2325	44	0
2	F	100	0	94	1	0
2	G	76	0	71	2	0
3	A	27	0	12	1	0
3	B	27	0	12	2	0
4	A	8	0	0	0	0
4	B	8	0	0	1	0
5	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	2	0	0	0	0
6	A	3	0	0	0	0
6	B	3	0	0	0	0
7	A	13	0	0	0	0
7	B	10	0	0	0	0
7	F	2	0	0	1	0
All	All	5298	0	4857	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:306:PHE:HB3	1:A:318:LEU:HB2	1.56	0.87
1:B:300:GLN:O	1:B:304:HIS:ND1	2.13	0.81
1:B:306:PHE:HB3	1:B:318:LEU:HB2	1.70	0.72
1:A:426:ARG:NH2	1:A:499:LYS:O	2.26	0.68
1:A:417:PRO:HA	1:A:420:ILE:HD12	1.76	0.66
1:A:377:ARG:NH2	1:A:414:CYS:SG	2.68	0.66
1:B:375:ILE:HD11	1:B:404:LEU:HD11	1.79	0.65
1:B:266:ALA:HB2	1:B:285:VAL:HG22	1.78	0.64
1:B:327:ARG:NH1	1:B:564:TPO:O	2.30	0.64
1:B:401:LYS:HE3	1:B:410:THR:HG21	1.79	0.64
1:A:327:ARG:NH1	1:A:564:TPO:O	2.30	0.63
1:A:522:PHE:HB3	1:A:527:TRP:HZ2	1.64	0.63
1:A:482:PRO:HG2	1:A:485:LEU:HD13	1.81	0.63
3:B:601:ADP:O1B	4:B:605:AF3:F1	2.09	0.61
1:B:417:PRO:HA	1:B:420:ILE:HD12	1.83	0.60
1:A:321:CYS:HB2	1:A:584:TYR:HB3	1.85	0.59
1:A:266:ALA:HB2	1:A:285:VAL:HG22	1.86	0.57
1:B:378:ASP:O	1:B:383:ASN:ND2	2.35	0.56
1:A:309:ALA:HB1	1:A:315:LEU:HB3	1.88	0.54
1:B:399:MET:SD	2:G:13:ARG:NH1	2.81	0.54
1:B:391:HIS:CD2	1:B:537:PRO:HG3	2.42	0.53
1:B:366:LEU:HD11	1:B:379:LEU:HD22	1.91	0.53
1:A:344:MET:HE3	1:A:446:MET:HG2	1.90	0.53
1:B:375:ILE:O	1:B:400:CYS:HA	2.08	0.53
1:B:426:ARG:NH1	1:B:428:GLU:OE2	2.36	0.53
1:B:283:LYS:HB3	1:B:330:PHE:HB2	1.91	0.53
1:A:337:GLY:O	1:A:343:HIS:NE2	2.38	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:324:THR:HG22	1:A:325:GLU:H	1.75	0.52
1:B:368:TYR:OH	1:B:372:ARG:NH1	2.42	0.51
1:B:426:ARG:NH2	1:B:499:LYS:O	2.43	0.51
1:A:401:LYS:HG2	1:A:410:THR:HG21	1.90	0.51
1:A:508:HIS:ND1	1:A:515:ASP:OD2	2.44	0.51
1:A:257:LEU:HD12	1:A:270:LEU:HD22	1.92	0.51
1:A:283:LYS:HB3	1:A:330:PHE:HB2	1.92	0.51
1:A:378:ASP:O	1:A:383[B]:ASN:ND2	2.43	0.51
1:A:333:GLU:O	3:A:601:ADP:N6	2.43	0.50
1:A:324:THR:HG22	1:A:325:GLU:N	2.26	0.50
1:A:262:ARG:HB2	1:A:556:PHE:CE1	2.47	0.49
1:B:374:ILE:HA	1:B:401:LYS:O	2.12	0.49
1:B:251:LEU:HD12	1:B:324:THR:HG21	1.96	0.48
1:B:339:ASP:HB3	1:B:385:LEU:HD23	1.95	0.47
1:A:339:ASP:HA	1:A:385:LEU:HA	1.95	0.47
1:A:391:HIS:CD2	1:A:537:PRO:HG3	2.50	0.47
1:B:399:MET:HA	2:G:13:ARG:HH12	1.78	0.47
1:B:522:PHE:HB3	1:B:527:TRP:HZ2	1.79	0.47
1:A:366:LEU:HD11	1:A:379:LEU:HD22	1.97	0.47
1:B:284:VAL:CG1	1:B:327:ARG:HD3	2.45	0.47
1:B:495:SER:HB3	1:B:505:LEU:HB2	1.98	0.46
1:B:270:LEU:HD12	1:B:549:LEU:HD21	1.98	0.46
1:B:482:PRO:HG2	1:B:485:LEU:HD13	1.97	0.46
1:B:309:ALA:HB1	1:B:315:LEU:HB3	1.98	0.46
1:A:489:ALA:O	1:A:492:VAL:HG12	2.16	0.46
1:B:339:ASP:HA	1:B:385:LEU:HA	1.98	0.46
1:B:343:HIS:NE2	1:B:539:PHE:CE2	2.84	0.46
1:B:377:ARG:NH1	1:B:412:TPO:O1P	2.38	0.45
1:B:335:VAL:HG12	1:B:386:LEU:O	2.17	0.45
1:A:339:ASP:HB3	1:A:385:LEU:HD23	1.97	0.45
1:B:328:LEU:HD11	1:B:579:PHE:CE1	2.51	0.45
1:B:279:ILE:HG12	1:B:547:PHE:CE2	2.50	0.45
1:A:284:VAL:CG1	1:A:327:ARG:HD3	2.46	0.45
1:B:508:HIS:O	1:B:512:GLY:N	2.50	0.45
1:A:335:VAL:HG12	1:A:386:LEU:O	2.17	0.45
1:A:422:PRO:HD3	1:A:437:TRP:CE2	2.53	0.44
1:B:410:THR:HG22	1:B:411:SER:H	1.82	0.44
1:A:270:LEU:HD12	1:A:549:LEU:HD21	2.00	0.44
1:A:513:PHE:O	1:A:517:GLN:HG3	2.17	0.44
1:B:355:ALA:O	1:B:359:SER:OG	2.25	0.43
1:B:489:ALA:O	1:B:492:VAL:HG12	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:376:TYR:OH	1:B:395:THR:O	2.33	0.43
1:A:258:ARG:HG3	1:A:259:VAL:N	2.34	0.43
1:A:375:ILE:HG22	1:A:377:ARG:HG3	2.01	0.42
1:B:312:HIS:CG	1:B:313:PRO:HD2	2.54	0.42
1:A:464:GLN:N	1:A:468:ASP:HB2	2.34	0.42
1:B:391:HIS:HE2	1:B:537:PRO:HD3	1.84	0.42
2:F:13:ARG:NH2	7:F:101:HOH:O	2.50	0.42
1:B:257:LEU:HD12	1:B:270:LEU:HD22	2.01	0.42
1:A:391:HIS:HD2	1:A:537:PRO:HG3	1.84	0.42
1:A:272:ARG:HB2	1:A:279:ILE:HD13	2.00	0.42
1:B:279:ILE:HG12	1:B:547:PHE:HE2	1.85	0.42
1:A:375:ILE:O	1:A:400:CYS:HA	2.20	0.41
1:B:404:LEU:HD12	1:B:432:PHE:CE1	2.55	0.41
1:A:253:ASP:O	1:A:273:LEU:HD12	2.21	0.41
1:B:333:GLU:O	3:B:601:ADP:N6	2.52	0.41
1:A:340:LEU:HA	1:A:340:LEU:HD23	1.94	0.40
1:A:377:ARG:NH1	1:A:401:LYS:HB2	2.36	0.40
1:B:262:ARG:NH2	1:B:564:TPO:O2P	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/349 (91%)	300 (94%)	18 (6%)	1 (0%)	46	83
1	B	312/349 (89%)	296 (95%)	16 (5%)	0	100	100
2	F	11/16 (69%)	10 (91%)	1 (9%)	0	100	100
2	G	8/16 (50%)	8 (100%)	0	0	100	100
All	All	650/730 (89%)	614 (94%)	35 (5%)	1 (0%)	52	87

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	299	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/311 (79%)	242 (98%)	5 (2%)	63	87
1	B	248/311 (80%)	245 (99%)	3 (1%)	78	92
2	F	9/15 (60%)	9 (100%)	0	100	100
2	G	7/15 (47%)	7 (100%)	0	100	100
All	All	511/652 (78%)	503 (98%)	8 (2%)	70	89

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	255	ASP
1	A	277	ASP
1	A	418	ASN
1	A	496	PHE
1	A	524	ASN
1	B	255	ASP
1	B	418	ASN
1	B	496	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	412	1	7,10,11	1.68	1 (14%)	10,14,16	1.34	1 (10%)
1	TPO	A	564	1	7,10,11	1.69	1 (14%)	10,14,16	1.42	2 (20%)
1	TPO	B	412	1	7,10,11	1.67	1 (14%)	10,14,16	1.66	2 (20%)
1	TPO	B	564	1	7,10,11	1.16	0	10,14,16	1.37	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	412	1	-	0/8/11/13	0/0/0/0
1	TPO	A	564	1	-	0/8/11/13	0/0/0/0
1	TPO	B	412	1	-	0/8/11/13	0/0/0/0
1	TPO	B	564	1	-	0/8/11/13	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	412	TPO	P-O1P	3.25	1.61	1.50
1	B	412	TPO	P-O1P	3.30	1.61	1.50
1	A	564	TPO	P-O1P	3.34	1.61	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	412	TPO	P-OG1-CB	-3.43	106.42	121.42
1	A	564	TPO	P-OG1-CB	-3.35	106.74	121.42
1	B	564	TPO	P-OG1-CB	-3.19	107.47	121.42
1	A	412	TPO	P-OG1-CB	-2.81	109.13	121.42
1	A	564	TPO	O-C-CA	-2.06	120.07	125.69
1	B	412	TPO	C-CA-N	2.17	114.74	109.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	564	TPO	1	0
1	B	412	TPO	1	0
1	B	564	TPO	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 5 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	A	601	5	24,29,29	1.00	1 (4%)	23,45,45	1.66	1 (4%)
4	AF3	A	602	-	0,3,3	0.00	-	0,3,3	0.00	-
4	AF3	A	605	5	0,3,3	0.00	-	0,3,3	0.00	-
6	SCN	A	606	-	2,2,2	2.02	1 (50%)	1,1,1	0.41	0
3	ADP	B	601	5	24,29,29	1.00	1 (4%)	23,45,45	1.69	1 (4%)
4	AF3	B	602	-	0,3,3	0.00	-	0,3,3	0.00	-
4	AF3	B	605	5	0,3,3	0.00	-	0,3,3	0.00	-
6	SCN	B	606	-	2,2,2	2.01	1 (50%)	1,1,1	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	601	5	-	0/12/32/32	0/3/3/3
4	AF3	A	602	-	-	0/0/0/0	0/0/0/0
4	AF3	A	605	5	-	0/0/0/0	0/0/0/0
6	SCN	A	606	-	-	0/0/0/0	0/0/0/0
3	ADP	B	601	5	-	0/12/32/32	0/3/3/3
4	AF3	B	602	-	-	0/0/0/0	0/0/0/0
4	AF3	B	605	5	-	0/0/0/0	0/0/0/0
6	SCN	B	606	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	606	SCN	C-S	2.80	1.80	1.63
6	A	606	SCN	C-S	2.81	1.80	1.63
3	B	601	ADP	C5-C4	3.11	1.47	1.40
3	A	601	ADP	C5-C4	3.15	1.47	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	ADP	N3-C2-N1	-6.50	123.76	128.87
3	A	601	ADP	N3-C2-N1	-6.47	123.79	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	ADP	1	0
3	B	601	ADP	2	0
4	B	605	AF3	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/349 (92%)	0.17	6 (1%) 70 60	27, 40, 54, 64	0
1	B	318/349 (91%)	0.30	6 (1%) 70 60	32, 46, 63, 83	0
2	F	13/16 (81%)	0.54	0 100 100	39, 43, 61, 62	0
2	G	10/16 (62%)	0.38	0 100 100	42, 47, 58, 59	0
All	All	665/730 (91%)	0.24	12 (1%) 71 62	27, 43, 60, 83	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	465	ASN	4.1
1	A	466	THR	3.5
1	A	248	SER	3.0
1	B	529	MET	2.5
1	A	524	ASN	2.5
1	A	276	THR	2.4
1	B	577	SER	2.3
1	B	576	GLN	2.2
1	B	487	VAL	2.1
1	B	538	PRO	2.1
1	A	290	VAL	2.0
1	B	386	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	TPO	A	564	11/12	0.96	0.19	-	32,33,35,35	0
1	TPO	B	564	11/12	0.93	0.16	-	44,46,49,50	0
1	TPO	A	412	11/12	0.93	0.23	-	37,40,44,45	0
1	TPO	B	412	11/12	0.94	0.26	-	53,57,58,59	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	AF3	A	602	4/4	0.89	0.43	7.83	37,37,37,37	0
6	SCN	B	606	3/3	0.85	0.27	0.37	65,65,65,65	0
4	AF3	B	602	4/4	0.95	0.26	0.27	36,36,36,36	0
6	SCN	A	606	3/3	0.88	0.19	-0.74	37,37,37,37	0
4	AF3	B	605	4/4	0.90	0.19	-1.02	26,39,59,68	0
3	ADP	B	601	27/27	0.93	0.20	-1.06	32,44,48,49	0
3	ADP	A	601	27/27	0.96	0.18	-1.72	25,31,44,58	0
4	AF3	A	605	4/4	0.94	0.18	-1.86	22,34,54,63	0
5	MN	B	604	1/1	0.96	0.12	-2.61	40,40,40,40	0
5	MN	A	604	1/1	0.98	0.14	-3.30	38,38,38,38	0
5	MN	A	603	1/1	0.97	0.10	-4.46	31,31,31,31	0
5	MN	A	607	1/1	0.90	0.12	-	56,56,56,56	0
5	MN	B	603	1/1	0.97	0.07	-	50,50,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.