



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 27, 2016 – 02:42 AM BST

PDB ID : 2LMO
Title : Structural Model for a 40-Residue Beta-Amyloid Fibril with Two-Fold Symmetry, Negative Stagger
Authors : Tycko, R.; Petkova, A.
Deposited on : 2011-12-08

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLID-STATE NMR

The overall completeness of chemical shifts assignment is 2%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	40	40% 20% 5% 15% 20%
1	B	40	35% 30% 13% 20%
1	C	40	25% 35% 20% 20%
1	D	40	20% 30% 8% 23% 20%
1	E	40	35% 35% 8% 20%
1	F	40	48% 30% 20%
1	G	40	33% 30% 5% 13% 20%
1	H	40	25% 30% 5% 20% 20%
1	I	40	25% 38% 10% 8% 20%

Continued on next page...

Mol	Chain	Length	Quality of chain
1	J	40	
1	K	40	
1	L	40	

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:11-A:23, A:28-A:40, B:9-B:23, B:29-B:40, C:12-C:23, C:29-C:40, D:12-D:23, D:30-D:40, E:9-E:26, E:30-E:40, F:9-F:40, G:13-G:39, H:12-H:24, H:30-H:40, I:12-I:40, J:12-J:39, K:11-K:39, L:12-L:40 (327)	1.36	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 5, 6, 7, 8, 9, 10
Single-model clusters	4

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 5700 atoms, of which 2844 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Beta-amyloid protein 40.


Mol	Chain	Residues	Atoms						Trace
1	A	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	B	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	C	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	D	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	E	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	F	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	G	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	H	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	I	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	J	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	K	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	
1	L	32	Total	C	H	N	O	S	0
			475	154	237	40	43	1	

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Beta-amyloid protein 40




- Molecule 1: Beta-amyloid protein 40




- Molecule 1: Beta-amyloid protein 40



- Molecule 1: Beta-amyloid protein 40




- Molecule 1: Beta-amyloid protein 40



- Molecule 1: Beta-amyloid protein 40

Chain F: 




- Molecule 1: Beta-amyloid protein 40

Chain G: 





- Molecule 1: Beta-amyloid protein 40

Chain H: 




- Molecule 1: Beta-amyloid protein 40

Chain I: 




- Molecule 1: Beta-amyloid protein 40

Chain J: 



- Molecule 1: Beta-amyloid protein 40

Chain K: 



- Molecule 1: Beta-amyloid protein 40

Chain L: 



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.

- Molecule 1: Beta-amyloid protein 40

Chain A: 




- Molecule 1: Beta-amyloid protein 40

Chain B: 




- Molecule 1: Beta-amyloid protein 40

Chain C: 





- Molecule 1: Beta-amyloid protein 40

Chain D: 



- Molecule 1: Beta-amyloid protein 40

Chain E: 




- Molecule 1: Beta-amyloid protein 40

Chain F: 





- Molecule 1: Beta-amyloid protein 40


Chain G: 




- Molecule 1: Beta-amyloid protein 40




- Molecule 1: Beta-amyloid protein 40



- Molecule 1: Beta-amyloid protein 40



- Molecule 1: Beta-amyloid protein 40



- Molecule 1: Beta-amyloid protein 40



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 40 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2lmo_cs.str
Number of chemical shift lists	1
Total number of shifts	110
Number of shifts mapped to atoms	110
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	2%

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.72±0.03	0±0/200 (0.0±0.0%)	0.89±0.13	0±1/266 (0.2±0.5%)
1	B	0.83±0.10	0±0/208 (0.1±0.2%)	0.94±0.13	1±1/278 (0.3±0.4%)
1	C	0.82±0.14	0±0/182 (0.1±0.2%)	0.90±0.09	0±1/243 (0.1±0.2%)
1	D	0.77±0.09	0±0/178 (0.1±0.2%)	0.89±0.08	0±1/238 (0.2±0.3%)
1	E	0.73±0.02	0±0/221 (0.0±0.0%)	0.79±0.07	0±0/296 (0.0±0.1%)
1	F	0.69±0.02	0±0/242 (0.0±0.0%)	0.83±0.06	0±0/323 (0.0±0.0%)
1	G	0.72±0.08	0±0/202 (0.0±0.1%)	0.86±0.07	0±0/271 (0.0±0.1%)
1	H	0.71±0.06	0±0/185 (0.0±0.0%)	0.86±0.06	0±0/248 (0.1±0.2%)
1	I	0.76±0.05	0±0/216 (0.1±0.2%)	0.94±0.09	0±1/288 (0.1±0.2%)
1	J	0.78±0.09	0±0/209 (0.0±0.1%)	1.04±0.16	1±1/281 (0.3±0.4%)
1	K	0.71±0.04	0±0/218 (0.0±0.0%)	0.79±0.05	0±0/293 (0.0±0.0%)
1	L	0.62±0.03	0±0/216 (0.0±0.0%)	0.78±0.03	0±0/288 (0.0±0.0%)
All	All	0.74	10/24770 (0.0%)	0.88	34/33130 (0.1%)

5 of 7 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	29	GLY	CA-C	-7.86	1.39	1.51	2	3
1	J	30	ALA	N-CA	-6.00	1.34	1.46	7	1
1	I	29	GLY	CA-C	-5.86	1.42	1.51	6	2
1	C	29	GLY	CA-C	-5.63	1.42	1.51	4	1
1	D	30	ALA	N-CA	-5.57	1.35	1.46	5	1

5 of 25 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	J	29	GLY	CA-C-N	-8.98	97.44	117.20	7	3
1	B	30	ALA	N-CA-C	-8.74	87.40	111.00	10	1
1	J	30	ALA	N-CA-C	-7.82	89.89	111.00	3	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	29	GLY	CA-C-N	-7.42	100.88	117.20	10	1
1	B	30	ALA	N-CA-CB	7.36	120.41	110.10	2	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	197	202	202	9±3
1	B	204	201	200	15±4
1	C	179	183	183	15±5
1	D	175	180	180	17±5
1	E	217	215	214	20±7
1	F	238	237	236	14±2
1	G	198	201	201	17±12
1	H	182	189	189	19±12
1	I	213	219	219	27±5
1	J	205	210	210	30±7
1	K	214	216	216	24±7
1	L	213	219	219	16±6
All	All	24350	24720	24689	1500

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

5 of 1125 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:I:26:SER:O	1:I:27:ASN:O	1.00	1.78	8	4
1:C:29:GLY:O	1:D:30:ALA:HB1	0.99	1.56	4	1
1:G:18:VAL:C	1:H:18:VAL:O	0.97	2.03	3	2
1:K:17:LEU:O	1:L:17:LEU:C	0.94	2.05	8	1
1:B:31:ILE:N	1:B:31:ILE:HD13	0.91	1.78	9	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	25/40 (62%)	19±1 (78±3%)	4±1 (15±4%)	2±1 (7±3%)	3	17
1	B	25/40 (62%)	21±1 (83±3%)	3±1 (13±4%)	1±1 (4±3%)	6	30
1	C	23/40 (58%)	18±1 (80±6%)	3±1 (15±3%)	1±1 (5±4%)	5	25
1	D	22/40 (55%)	17±1 (78±4%)	3±1 (15±4%)	2±1 (8±4%)	2	16
1	E	27/40 (68%)	21±1 (77±3%)	4±1 (14±4%)	2±1 (9±3%)	2	13
1	F	30/40 (75%)	22±1 (72±4%)	5±1 (16±5%)	4±1 (12±3%)	1	8
1	G	27/40 (68%)	18±1 (67±2%)	5±1 (20±4%)	4±1 (13±4%)	1	6
1	H	23/40 (58%)	18±1 (79±5%)	3±1 (14±5%)	2±1 (7±4%)	3	18
1	I	28/40 (70%)	20±1 (70±3%)	5±1 (16±4%)	4±1 (14±3%)	1	5
1	J	28/40 (70%)	19±2 (69±6%)	5±2 (19±7%)	4±1 (12±4%)	1	7
1	K	29/40 (72%)	20±1 (70±3%)	5±1 (18±5%)	4±1 (12±4%)	1	7
1	L	28/40 (70%)	19±1 (69±4%)	6±1 (20±3%)	3±1 (11±3%)	1	8
All	All	3150/4800 (66%)	2325 (74%)	516 (16%)	309 (10%)	2	11

5 of 88 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	D	37	GLY	8
1	G	25	GLY	8
1	L	28	LYS	8
1	K	25	GLY	8
1	L	25	GLY	7

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	20/31 (65%)	18±1 (89±4%)	2±1 (11±4%)	12	56
1	B	20/31 (65%)	18±1 (91±4%)	2±1 (10±4%)	15	60
1	C	18/31 (58%)	16±1 (87±8%)	2±1 (13±8%)	10	51
1	D	18/31 (58%)	16±1 (86±8%)	3±1 (14±8%)	8	49
1	E	22/31 (71%)	20±1 (92±4%)	2±1 (8±4%)	19	64
1	F	24/31 (77%)	23±1 (94±2%)	2±1 (6±2%)	27	72
1	G	20/31 (65%)	18±1 (88±4%)	2±1 (12±4%)	11	53
1	H	19/31 (61%)	16±1 (86±3%)	3±1 (14±3%)	9	49
1	I	22/31 (71%)	19±1 (87±5%)	3±1 (13±5%)	10	51
1	J	21/31 (68%)	18±1 (85±5%)	3±1 (15±5%)	8	47
1	K	22/31 (71%)	20±1 (91±5%)	2±1 (9±5%)	18	63
1	L	22/31 (71%)	20±1 (93±3%)	2±1 (7±3%)	22	68
All	All	2480/3720 (67%)	2214 (89%)	266 (11%)	13	56

5 of 88 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	G	36	VAL	10
1	L	31	ILE	10
1	K	31	ILE	10
1	H	36	VAL	10
1	D	36	VAL	9

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 2% for the well-defined parts and 2% for the entire structure.

7.1 Chemical shift list 1

File name: 2lmo_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	110
Number of shifts mapped to atoms	110
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	33	0.14 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	27	-1.34 ± 0.17	Should be applied
$^{13}\text{C}'$	33	1.36 ± 0.18	Should be applied
^{15}N	17	-1.00 ± 0.51	None needed (imprecise)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 2%, i.e. 81 atoms were assigned a chemical shift out of a possible 3753. 0 out of 88 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	61/1635 (4%)	0/654 (0%)	48/654 (7%)	13/327 (4%)
Sidechain	20/1710 (1%)	0/972 (0%)	20/701 (3%)	0/37 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	0/408 (0%)	0/228 (0%)	0/156 (0%)	0/24 (0%)
Overall	81/3753 (2%)	0/1854 (0%)	68/1511 (5%)	13/388 (3%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

