



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:20 AM GMT

PDB ID : 3LP1
Title : HIV-1 reverse transcriptase with inhibitor
Authors : Yan, Y.; Munshi, S.K.; Prasad, G.S.; Su, H.P.
Deposited on : 2010-02-04
Resolution : 2.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1611 (2.26-2.22)
Clashscore	102246	1764 (2.26-2.22)
Ramachandran outliers	100387	1724 (2.26-2.22)
Sidechain outliers	100360	1724 (2.26-2.22)
RSRZ outliers	91569	1616 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	563	<div> <div>6%</div> <div>84%</div> <div>14%</div> <div>••</div> </div>
2	B	443	<div> <div>11%</div> <div>77%</div> <div>12%</div> <div>• 10%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8029 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase/ribonuclease H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	554	Total	C	N	O	S	0	0	0
			4503	2911	751	833	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	EXPRESSION TAG	UNP P04585
A	-1	ASN	-	EXPRESSION TAG	UNP P04585
A	0	SER	-	EXPRESSION TAG	UNP P04585


- Molecule 2 is a protein called p51 RT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	400	Total	C	N	O	S	0	0	0
			3311	2155	548	602	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	MET	-	EXPRESSION TAG	UNP P04585
B	-1	ASN	-	EXPRESSION TAG	UNP P04585
B	0	SER	-	EXPRESSION TAG	UNP P04585

- Molecule 3 is 3-CYCLOPENTYL-1,4-DIHYDROXY-1,8-NAPHTHYRIDIN-2(1H)-ONE (three-letter code: LP8) (formula: C₁₃H₁₄N₂O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			18	13	2	3		

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mn	0	0
			2	2		

- Molecule 5 is 11-CYCLOPROPYL-5,11-DIHYDRO-4-METHYL-6H-DIPYRIDO[3,2-B:2',3'-E][1,4]DIAZEPIN-6-ONE (three-letter code: NVP) (formula: C₁₅H₁₄N₄O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			20	15	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	108	Total	O	0	0
			108	108		
6	B	67	Total	O	0	0
			67	67		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Reverse transcriptase/ribonuclease H



- Molecule 2: p51 RT



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	118.78Å 155.26Å 155.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.23 43.06 – 2.23	Depositor EDS
% Data completeness (in resolution range)	96.2 (50.00-2.23) 96.2 (43.06-2.23)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.28 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.268 , 0.292 0.265 , 0.290	Depositor DCC
R_{free} test set	3399 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	41.6	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 67665 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8029	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LP8, MN, NVP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/4619	0.50	0/6279
2	B	0.33	0/3404	0.49	0/4627
All	All	0.33	0/8023	0.50	0/10906

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4503	0	4540	38	0
2	B	3311	0	3326	28	0
3	A	18	0	12	0	0
4	A	2	0	0	0	0
5	A	20	0	14	1	0
6	A	108	0	0	2	0
6	B	67	0	0	0	0
All	All	8029	0	7892	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:GLN:HE22	1:A:340:GLN:HE22	1.28	0.80
2:B:396:GLU:O	2:B:400:THR:HG22	1.84	0.78
2:B:337:TRP:HE1	2:B:367:GLN:HE21	1.32	0.75
2:B:12:LEU:HD23	2:B:17:ASP:HA	1.78	0.65
1:A:91:GLN:HG3	1:A:93:GLY:O	1.97	0.65
1:A:175:ASN:HB3	1:A:178:ILE:HD12	1.78	0.64
2:B:373:GLN:O	2:B:377:THR:HG23	1.98	0.64
2:B:266:TRP:HH2	2:B:427:TYR:CZ	2.16	0.62
1:A:369:THR:CG2	1:A:398:TRP:HZ3	2.12	0.62
2:B:214:LEU:HD23	2:B:214:LEU:H	1.67	0.60
1:A:199:ARG:HA	1:A:202:ILE:HD12	1.84	0.58
1:A:373:GLN:HG3	6:A:562:HOH:O	2.04	0.57
2:B:388:LYS:HE2	2:B:415:GLU:HG3	1.86	0.57
1:A:91:GLN:HG3	1:A:93:GLY:H	1.69	0.56
1:A:181:TYR:CE2	1:A:183:TYR:HB2	2.40	0.56
1:A:46:LYS:HE3	1:A:116:PHE:HB3	1.87	0.56
1:A:365:VAL:O	1:A:369:THR:HG23	2.06	0.55
1:A:94:ILE:HD13	1:A:230:MET:HG2	1.90	0.54
1:A:369:THR:HG22	1:A:398:TRP:HZ3	1.72	0.54
2:B:209:LEU:HG	2:B:214:LEU:HD12	1.89	0.53
1:A:399:GLU:HA	1:A:402:TRP:CE3	2.43	0.53
1:A:91:GLN:CG	1:A:93:GLY:O	2.57	0.53
1:A:369:THR:HG22	1:A:398:TRP:CZ3	2.44	0.53
1:A:106:VAL:HG12	1:A:236:PRO:HB3	1.91	0.53
2:B:13:LYS:HE3	2:B:85:GLN:HB3	1.91	0.52
1:A:544:GLY:HA2	2:B:286:THR:HG22	1.91	0.52
2:B:373:GLN:HE22	2:B:407:GLN:H	1.58	0.51
1:A:183:TYR:OH	1:A:230:MET:HE1	2.11	0.51
2:B:377:THR:HG22	2:B:410:TRP:HZ2	1.76	0.51
2:B:373:GLN:NE2	2:B:407:GLN:H	2.09	0.50
1:A:244:ILE:HD13	1:A:267:ALA:HB2	1.92	0.50
1:A:179:VAL:HG23	5:A:701:NVP:HCC1	1.93	0.50
2:B:345:PRO:O	2:B:346:PHE:HB2	2.10	0.49
2:B:362:THR:HG22	2:B:367:GLN:HG3	1.93	0.49
2:B:362:THR:HG23	2:B:366:LYS:HE3	1.95	0.49
2:B:169:GLU:HB3	2:B:170:PRO:HD3	1.95	0.49
1:A:303:LEU:O	1:A:307:ARG:HG3	2.14	0.48
1:A:183:TYR:OH	1:A:230:MET:CE	2.62	0.48
1:A:354:TYR:HD1	1:A:374:LYS:HD2	1.78	0.47
1:A:91:GLN:HG2	6:A:573:HOH:O	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:406:TRP:CH2	2:B:418:ASN:HA	2.49	0.46
1:A:169:GLU:HB3	1:A:170:PRO:HD3	1.98	0.46
2:B:13:LYS:O	2:B:16:MET:HG2	2.16	0.45
2:B:250:ASP:O	2:B:251:SER:HB3	2.17	0.45
1:A:328:GLU:HG2	1:A:390:LYS:HB2	1.97	0.45
2:B:180:ILE:HG12	2:B:189:VAL:HG13	1.99	0.45
1:A:136:ASN:HB3	1:A:137:ASN:H	1.63	0.45
2:B:58:THR:HG23	2:B:76:ASP:O	2.17	0.44
2:B:167:ILE:HG12	2:B:212:TRP:CD2	2.54	0.43
1:A:108:VAL:HG22	1:A:188:TYR:CE2	2.54	0.43
1:A:402:TRP:C	1:A:402:TRP:CD1	2.92	0.42
1:A:452:LEU:HD23	1:A:470:THR:HA	2.01	0.42
1:A:230:MET:HA	1:A:230:MET:HE2	2.01	0.42
1:A:547:GLN:HA	1:A:550:LYS:HE2	2.00	0.42
2:B:157:PRO:HG3	2:B:184:MET:HA	2.01	0.42
1:A:536:VAL:HB	1:A:542:ILE:HD13	2.01	0.41
1:A:54:ASN:HB3	1:A:143:ARG:HH21	1.86	0.41
2:B:175:ASN:HD21	2:B:201:LYS:NZ	2.18	0.41
2:B:365:VAL:O	2:B:369:THR:HG23	2.21	0.41
1:A:485:ALA:O	1:A:489:SER:HB3	2.21	0.41
1:A:330:GLN:NE2	1:A:340:GLN:HE22	2.07	0.41
2:B:94:ILE:HA	2:B:95:PRO:HD3	1.91	0.41
1:A:411:ILE:HG22	1:A:412:PRO:O	2.20	0.41
2:B:246:LEU:HA	2:B:247:PRO:HD3	1.92	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	550/563 (98%)	532 (97%)	14 (2%)	4 (1%)	26 25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	392/443 (88%)	377 (96%)	14 (4%)	1 (0%)	46	51
All	All	942/1006 (94%)	909 (96%)	28 (3%)	5 (0%)	34	34

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	GLU
1	A	543	GLY
2	B	296	THR
1	A	345	PRO
1	A	287	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	492/503 (98%)	460 (94%)	32 (6%)	21	19
2	B	364/403 (90%)	344 (94%)	20 (6%)	27	26
All	All	856/906 (94%)	804 (94%)	52 (6%)	23	22

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
1	A	6	GLU
1	A	7	THR
1	A	21	VAL
1	A	26	LEU
1	A	118	VAL
1	A	139	THR
1	A	161	GLN
1	A	173	LYS
1	A	205	LEU
1	A	210	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	211	ARG
1	A	220	LYS
1	A	228	LEU
1	A	248	GLU
1	A	260	LEU
1	A	276	VAL
1	A	279	LEU
1	A	284	ARG
1	A	287	LYS
1	A	297	GLU
1	A	303	LEU
1	A	368	LEU
1	A	373	GLN
1	A	402	TRP
1	A	448	ARG
1	A	479	LEU
1	A	496	VAL
1	A	503	LEU
1	A	523	GLU
1	A	547	GLN
1	A	548	VAL
2	B	6	GLU
2	B	11	LYS
2	B	12	LEU
2	B	26	LEU
2	B	72	ARG
2	B	80	LEU
2	B	91	GLN
2	B	111	VAL
2	B	116	PHE
2	B	120	LEU
2	B	205	LEU
2	B	209	LEU
2	B	212	TRP
2	B	214	LEU
2	B	297	GLU
2	B	347	LYS
2	B	349	LEU
2	B	362	THR
2	B	400	THR
2	B	425	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	103	ASN
1	A	147	ASN
1	A	198	HIS
1	A	258	GLN
1	A	330	GLN
1	A	407	GLN
1	A	500	GLN
1	A	507	GLN
1	A	509	GLN
1	A	519	ASN
1	A	520	GLN
1	A	524	GLN
2	B	147	ASN
2	B	161	GLN
2	B	175	ASN
2	B	208	HIS
2	B	258	GLN
2	B	278	GLN
2	B	306	ASN
2	B	367	GLN
2	B	373	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	LP8	A	601	4	17,20,20	1.38	2 (11%)	19,29,29	1.37	2 (10%)
5	NVP	A	701	-	18,23,23	1.29	2 (11%)	18,34,34	1.54	5 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LP8	A	601	4	-	0/4/11/11	0/3/3/3
5	NVP	A	701	-	-	0/0/6/6	0/2/4/4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	LP8	C-N	-2.59	1.36	1.40
5	A	701	NVP	CC-CA	2.55	1.53	1.48
5	A	701	NVP	CB-CA	2.59	1.53	1.48
3	A	601	LP8	O3-C	3.80	1.33	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	LP8	C5-C4-N	-4.20	116.62	121.79
5	A	701	NVP	C4-C5-C6	-3.32	117.59	120.13
5	A	701	NVP	C10-C15-N14	-2.37	120.72	123.53
5	A	701	NVP	CC-CA-N1	-2.04	116.15	118.25
5	A	701	NVP	C5-C4-N3	-2.03	121.36	124.56
5	A	701	NVP	C13-N14-C15	2.05	119.48	116.93
3	A	601	LP8	O10-C1-C2	2.30	124.63	121.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	701	NVP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	554/563 (98%)	0.55	36 (6%) 22 23	27, 44, 56, 62	0
2	B	400/443 (90%)	0.70	48 (12%) 6 6	29, 40, 80, 83	0
All	All	954/1006 (94%)	0.61	84 (8%) 12 13	27, 43, 71, 83	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	301	LEU	7.1
2	B	90	VAL	7.1
2	B	362	THR	5.6
2	B	361	HIS	5.4
2	B	231	GLY	4.3
2	B	6	GLU	4.2
2	B	299	ALA	4.1
1	A	52	PRO	4.0
2	B	298	GLU	3.9
2	B	69	THR	3.9
1	A	286	THR	3.9
2	B	346	PHE	3.7
2	B	277	ARG	3.7
2	B	292	VAL	3.7
2	B	251	SER	3.6
2	B	14	PRO	3.5
1	A	69	THR	3.5
1	A	360	ALA	3.4
2	B	314	VAL	3.3
1	A	357	MET	3.3
2	B	13	LYS	3.3
1	A	356	ARG	3.3
1	A	137	ASN	3.2
2	B	309	ILE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	305	GLU	3.2
1	A	195	ILE	3.2
1	A	402	TRP	3.2
2	B	232	TYR	3.2
2	B	283	LEU	3.2
1	A	514	GLU	3.2
2	B	310	LEU	3.1
1	A	118	VAL	3.1
2	B	266	TRP	3.1
2	B	302	GLU	3.1
2	B	250	ASP	3.0
1	A	221	HIS	2.9
1	A	220	LYS	2.9
2	B	70	LYS	2.8
1	A	334	GLN	2.8
1	A	193	LEU	2.8
2	B	257	ILE	2.7
2	B	240	THR	2.6
2	B	290	THR	2.6
2	B	271	TYR	2.6
2	B	293	ILE	2.6
1	A	203	GLU	2.5
1	A	24	TRP	2.5
2	B	92	LEU	2.5
1	A	63	ILE	2.5
1	A	173	LYS	2.4
2	B	297	GLU	2.4
1	A	491	LEU	2.4
2	B	276	VAL	2.4
2	B	24	TRP	2.4
2	B	272	PRO	2.4
2	B	311	LYS	2.3
1	A	219	LYS	2.3
1	A	550	LYS	2.3
1	A	196	GLY	2.3
1	A	426	TRP	2.3
2	B	274	ILE	2.3
1	A	199	ARG	2.3
2	B	279	LEU	2.3
1	A	72	ARG	2.3
1	A	526	ILE	2.2
2	B	286	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	90	VAL	2.2
1	A	543	GLY	2.2
2	B	16	MET	2.2
2	B	295	LEU	2.2
2	B	252	TRP	2.2
1	A	435	VAL	2.1
2	B	212	TRP	2.1
2	B	22	LYS	2.1
1	A	14	PRO	2.1
2	B	288	ALA	2.1
1	A	2	ILE	2.1
1	A	43	LYS	2.0
2	B	426	TRP	2.0
1	A	207	GLN	2.0
1	A	138	GLU	2.0
2	B	313	PRO	2.0
2	B	282	LEU	2.0
1	A	140	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	NVP	A	701	20/20	0.93	0.14	-0.60	35,35,36,36	0
3	LP8	A	601	18/18	0.92	0.12	-1.02	44,44,45,45	0
4	MN	A	602	1/1	0.97	0.14	-1.11	44,44,44,44	0
4	MN	A	603	1/1	0.99	0.10	-	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.