



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:30 PM GMT

PDB ID : 4LU1  
Title : Crystal structure of the uncharacterized Maf protein YceF from E. coli, mutant D69A  
Authors : Dong, A.; Xu, X.; Cui, H.; Tchigvintsev, A.; Flick, R.; Brown, G.; Popovic, A.; Yakunin, A.F.; Savchenko, A.; Structural Genomics Consortium (SGC)  
Deposited on : 2013-07-24  
Resolution : 1.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	5832 (1.94-1.90)
Clashscore	102246	6540 (1.94-1.90)
Ramachandran outliers	100387	6464 (1.94-1.90)
Sidechain outliers	100360	6465 (1.94-1.90)
RSRZ outliers	91569	5846 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	<div> <div>4%</div> <div>81% 7% 11%</div> </div>
1	B	215	<div> <div>10%</div> <div>79% • 17%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UNX	A	201	-	-	-	X
2	UNX	B	201	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maf-like protein YceF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	3	0
			1476	934	260	277	5			
1	B	178	Total	C	N	O	S	0	2	0
			1361	866	240	250	5			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP C9QYM5
A	-19	GLY	-	EXPRESSION TAG	UNP C9QYM5
A	-18	SER	-	EXPRESSION TAG	UNP C9QYM5
A	-17	SER	-	EXPRESSION TAG	UNP C9QYM5
A	-16	HIS	-	EXPRESSION TAG	UNP C9QYM5
A	-15	HIS	-	EXPRESSION TAG	UNP C9QYM5
A	-14	HIS	-	EXPRESSION TAG	UNP C9QYM5
A	-13	HIS	-	EXPRESSION TAG	UNP C9QYM5
A	-12	HIS	-	EXPRESSION TAG	UNP C9QYM5
A	-11	HIS	-	EXPRESSION TAG	UNP C9QYM5
A	-10	SER	-	EXPRESSION TAG	UNP C9QYM5
A	-9	SER	-	EXPRESSION TAG	UNP C9QYM5
A	-8	GLY	-	EXPRESSION TAG	UNP C9QYM5
A	-7	ARG	-	EXPRESSION TAG	UNP C9QYM5
A	-6	GLU	-	EXPRESSION TAG	UNP C9QYM5
A	-5	ASN	-	EXPRESSION TAG	UNP C9QYM5
A	-4	LEU	-	EXPRESSION TAG	UNP C9QYM5
A	-3	TYR	-	EXPRESSION TAG	UNP C9QYM5
A	-2	PHE	-	EXPRESSION TAG	UNP C9QYM5
A	-1	GLN	-	EXPRESSION TAG	UNP C9QYM5
A	0	GLY	-	EXPRESSION TAG	UNP C9QYM5
A	69	ALA	ASP	ENGINEERED MUTATION	UNP C9QYM5
B	-20	MET	-	EXPRESSION TAG	UNP C9QYM5
B	-19	GLY	-	EXPRESSION TAG	UNP C9QYM5
B	-18	SER	-	EXPRESSION TAG	UNP C9QYM5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	SER	-	EXPRESSION TAG	UNP C9QYM5
B	-16	HIS	-	EXPRESSION TAG	UNP C9QYM5
B	-15	HIS	-	EXPRESSION TAG	UNP C9QYM5
B	-14	HIS	-	EXPRESSION TAG	UNP C9QYM5
B	-13	HIS	-	EXPRESSION TAG	UNP C9QYM5
B	-12	HIS	-	EXPRESSION TAG	UNP C9QYM5
B	-11	HIS	-	EXPRESSION TAG	UNP C9QYM5
B	-10	SER	-	EXPRESSION TAG	UNP C9QYM5
B	-9	SER	-	EXPRESSION TAG	UNP C9QYM5
B	-8	GLY	-	EXPRESSION TAG	UNP C9QYM5
B	-7	ARG	-	EXPRESSION TAG	UNP C9QYM5
B	-6	GLU	-	EXPRESSION TAG	UNP C9QYM5
B	-5	ASN	-	EXPRESSION TAG	UNP C9QYM5
B	-4	LEU	-	EXPRESSION TAG	UNP C9QYM5
B	-3	TYR	-	EXPRESSION TAG	UNP C9QYM5
B	-2	PHE	-	EXPRESSION TAG	UNP C9QYM5
B	-1	GLN	-	EXPRESSION TAG	UNP C9QYM5
B	0	GLY	-	EXPRESSION TAG	UNP C9QYM5
B	69	ALA	ASP	ENGINEERED MUTATION	UNP C9QYM5

- Molecule 2 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total X 1 1	0	0
2	A	1	Total X 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	82	Total O 82 82	0	0
3	B	57	Total O 57 57	0	0



- Molecule 1: Maf-like protein YceF



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.65Å 52.83Å 129.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.42 – 1.92 33.39 – 1.92	Depositor EDS
% Data completeness (in resolution range)	98.8 (33.42-1.92) 98.8 (33.39-1.92)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.72 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.199 , 0.249 0.208 , 0.258	Depositor DCC
$R_{free}$ test set	858 reflections (3.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.9	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 55.2	EDS
Estimated twinning fraction	0.024 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 27031 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2978	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/1514	0.71	0/2053
1	B	0.43	0/1390	0.61	0/1883
All	All	0.46	0/2904	0.66	0/3936

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1476	0	1454	9	0
1	B	1361	0	1338	6	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	82	0	0	1	0
3	B	57	0	0	3	0
All	All	2978	0	2792	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below, sorted by their



clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:LEU:HD13	1:A:100:VAL:HG22	1.76	0.67
1:A:46:LEU:CG	1:A:120:VAL:CG2	2.76	0.62
1:A:46:LEU:CG	1:A:120:VAL:HG23	2.29	0.62
1:A:75:ASP:OD2	3:A:375:HOH:O	2.18	0.54
1:A:164:LEU:HD12	1:B:169:PRO:HD2	1.93	0.51

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/215 (89%)	188 (98%)	4 (2%)	0	100	100
1	B	174/215 (81%)	168 (97%)	6 (3%)	0	100	100
All	All	366/430 (85%)	356 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/184 (85%)	154 (99%)	2 (1%)	76	73
1	B	140/184 (76%)	138 (99%)	2 (1%)	74	70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	296/368 (80%)	292 (99%)	4 (1%)	74	70

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	SER
1	A	154	GLU
1	B	90	LEU
1	B	193	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	191/215 (88%)	0.25	8 (4%) 40 44	17, 30, 49, 61	0
1	B	178/215 (82%)	0.71	21 (11%) 6 7	17, 35, 64, 91	0
All	All	369/430 (85%)	0.47	29 (7%) 15 17	17, 31, 58, 91	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	90	LEU	7.7
1	B	131	SER	4.3
1	B	75	ASP	4.1
1	B	135	ILE	4.0
1	A	37	SER	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	UNX	A	201	1/1	0.83	0.24	8.11	30,30,30,30	0
2	UNX	B	201	1/1	0.88	0.27	4.72	30,30,30,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.