wwPDB NMR Structure Validation Summary Report (i)

Apr 27, 2016 – 02:48 AM BST

PDB ID	:	2M7W
Title	:	Independently verified structure of gp41-M-MAT, a membrane associated MPER trimer from HIV-1 gp41
Authors	:	Martin, J.W.; Reardon, P.N.; Sage, H.S.; Moses, D.S.; Munir, A.S.; Haynes, B.F.; Spicer, L.D.; Donald, B.R.
Deposited on	:	2013-05-01

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 19%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	59	66% 32% •
1	B	59	66% 31% •
1	C	59	66% 31% •

2 Ensemble composition and analysis (i)

This entry contains 18 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:1-A:59, B:1-B:59, C:1-C:59 (177)	1.16	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: Inconsistent models

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2913 atoms, of which 1413 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Envelope glycoprotein.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	59	971	329	471	80	91	0
1	B	59	971	329	471	80	91	0
1	C	59	971	329	471	80	91	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Envelope glycoprotein

Chain A:

- Molecule 1: Envelope glycoprotein

Chain B:

- Molecule 1: Envelope glycoprotein

Chain C:

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.

- Molecule 1: Envelope glycoprotein

Chain A:



- Molecule 1: Envelope glycoprotein

Chain B:



- Molecule 1: Envelope glycoprotein

Chain C:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *complete search, simulated annealing.*

Of the 200 calculated structures, 18 were deposited, based on the following criterion: *structures with the least restraint violations.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DISCO	structure solution	1
X-PLOR NIH	structure solution	2.33
X-PLOR NIH	refinement	2.33

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2m7w_cs.str
Number of chemical shift lists	1
Total number of shifts	485
Number of shifts mapped to atoms	485
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	19%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.34±0.09	0±0/518 (0.0±0.1%)	1.08±0.04	0±1/706 (0.1±0.2%)
1	B	1.34±0.09	0±0/518 (0.0±0.1%)	1.08±0.04	0±1/706 (0.1±0.2%)
1	C	1.34±0.09	0±0/518 (0.0±0.1%)	1.08±0.04	0±1/706 (0.1±0.2%)
All	All	1.35	7/27972 (0.0%)	1.08	24/38124 (0.1%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	38	GLU	CB-CG	6.18	1.63	1.52	8	2
1	B	38	GLU	CB-CG	6.15	1.63	1.52	8	2
1	C	38	GLU	CB-CG	6.13	1.63	1.52	8	2
1	C	35	GLU	CD-OE1	-5.12	1.20	1.25	2	1

5 of 24 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	B	26	PHE	CB-CG-CD1	8.20	126.54	120.80	6	1
1	C	26	PHE	CB-CG-CD1	8.07	126.45	120.80	6	1
1	A	26	PHE	CB-CG-CD1	8.05	126.43	120.80	6	1
1	B	13	TYR	CB-CG-CD2	-6.68	116.99	121.00	11	1
1	C	13	TYR	CB-CG-CD2	-6.67	117.00	121.00	11	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	500	471	471	13±5
1	B	500	471	471	14±6
1	C	500	471	471	14±5
All	All	27000	25434	25434	643

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

5 of 345 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:6:ALA:HB1	1:B:8:ARG:HH12	0.83	1.33	10	2
1:A:6:ALA:HB1	1:A:8:ARG:HH12	0.83	1.33	10	2
1:A:51:ILE:HD13	1:A:52:THR:N	0.82	1.90	18	2
1:C:6:ALA:HB1	1:C:8:ARG:HH12	0.81	1.33	10	2
1:B:51:ILE:HD13	1:B:52:THR:N	0.81	1.90	18	2

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	57/59 (97%)	48±2 (84±3%)	7±2 (13±3%)	2±1 (3±1%)	9	42
1	B	57/59 (97%)	48±2 (84±4%)	7±2 (13±3%)	2±1 (3±1%)	9	42
1	C	57/59 (97%)	48±2 (84±3%)	7±2 (13±3%)	2±1 (3±2%)	9	43
All	All	3078/3186 (97%)	2594 (84%)	392 (13%)	92 (3%)	9	42

5 of 21 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	3	ILE	13
1	A	3	ILE	13
1	C	3	ILE	13
1	B	29	SER	6
1	A	29	SER	6

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	51/51 (100%)	41±3 (81±6%)	10±3 (19±6%)	5 38
1	B	51/51 (100%)	41±3 (81±6%)	10±3 (19±6%)	5 38
1	C	51/51 (100%)	41±3 (81±6%)	10±3 (19±6%)	5 38
All	All	2754/2754 (100%)	2238 (81%)	516 (19%)	5 38

5 of 126 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	C	48	TRP	12
1	A	48	TRP	12
1	B	48	TRP	12
1	A	51	ILE	11
1	C	51	ILE	11

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 19% for the well-defined parts and 19% for the entire structure.

7.1 Chemical shift list 1

File name: 2m7w_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	485
Number of shifts mapped to atoms	485
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	55	0.13 ± 0.41	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	49	0.64 ± 0.18	Should be applied
$^{13}C'$	42	-0.45 ± 0.16	None needed (< 0.5 ppm)
^{15}N	55	-0.29 ± 0.21	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 19%, i.e. 434 atoms were assigned a chemical shift out of a possible 2289. 0 out of 30 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	250/873 (29%)	98/348 (28%)	97/354 (27%)	55/171 (32%)
Sidechain	172/1074 (16%)	116/630 (18%)	53/399 (13%)	3/45 (7%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	12/342 (4%)	6/174 (3%)	0/150 (0%)	6/18 (33%)
Overall	434/2289 (19%)	220/1152 (19%)	150/903 (17%)	64/234 (27%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	6	ALA	HB2	-0.42	2.61 – 0.11	-7.1
1	A	6	ALA	HB3	-0.42	2.61 – 0.11	-7.1
1	A	6	ALA	HB1	-0.42	2.61 – 0.11	-7.1

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: