



Full wwPDB NMR Structure Validation Report i

Sep 26, 2016 – 09:56 PM EDT

PDB ID : 2MA9
Title : HIV-1 Vif SOCS-box and Elongin BC solution structure
Authors : Lu, Z.; Bergeron, J.R.; Atkinson, R.A.; Schaller, T.; Veselkov, D.A.; Oregioni, A.; Yang, Y.; Matthews, S.J.; Malim, M.H.; Sanderson, M.R.
Deposited on : 2013-07-01

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:


Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbit	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027939
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
SOLUTION NMR


The overall completeness of chemical shifts assignment is 27%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$



2 Ensemble composition and analysis

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:6-A:37, B:1-B:118, C:1-C:91 (241)	0.20	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 3 single-model clusters were found.

Cluster number	Models
1	2, 3, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20
Single-model clusters	1; 4; 5

3 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 2307 atoms, of which 412 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Virion infectivity factor.

Mol	Chain	Residues	Atoms					Trace
1	A	32	Total	C	H	N	O	0
			316	169	59	45	43	

- Molecule 2 is a protein called Transcription elongation factor B polypeptide 2.

Mol	Chain	Residues	Atoms						Trace
2	B	118	Total	C	H	N	O	S	0
			1124	575	204	156	184	5	

- Molecule 3 is a protein called Transcription elongation factor B polypeptide 1.

Mol	Chain	Residues	Atoms						Trace
3	C	91	Total	C	H	N	O	S	0
			867	460	149	117	137	4	

4 Residue-property plots [\(i\)](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.2 Score per residue for model 2 (medoid)


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1



4.2.3 Score per residue for model 3

- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1



4.2.4 Score per residue for model 4

- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.5 Score per residue for model 5

- Molecule 1: Virion infectivity factor




- Molecule 2: Transcription elongation factor B polypeptide 2





- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.6 Score per residue for model 6


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.7 Score per residue for model 7


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.8 Score per residue for model 8


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2




- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.9 Score per residue for model 9


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.10 Score per residue for model 10


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.11 Score per residue for model 11

- Molecule 1: Virion infectivity factor




- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1





4.2.12 Score per residue for model 12

- Molecule 1: Virion infectivity factor

Chain A: 47% 33% 8% 11%



- Molecule 2: Transcription elongation factor B polypeptide 2

Chain B: 77% 22%



- Molecule 3: Transcription elongation factor B polypeptide 1

Chain C: 66% 34%



4.2.13 Score per residue for model 13

- Molecule 1: Virion infectivity factor

Chain A: 47% 33% 6% • 11%



- Molecule 2: Transcription elongation factor B polypeptide 2

Chain B: 76% 22% •




- Molecule 3: Transcription elongation factor B polypeptide 1

Chain C: 82% 18%




4.2.14 Score per residue for model 14


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2




- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.15 Score per residue for model 15


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.16 Score per residue for model 16


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2




- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.17 Score per residue for model 17


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2




- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.18 Score per residue for model 18


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.19 Score per residue for model 19


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2




- Molecule 3: Transcription elongation factor B polypeptide 1




4.2.20 Score per residue for model 20


- Molecule 1: Virion infectivity factor



- Molecule 2: Transcription elongation factor B polypeptide 2



- Molecule 3: Transcription elongation factor B polypeptide 1



5 Refinement protocol and experimental data overview i

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CS-Rosetta	structure solution	
CS-Rosetta	refinement	
HADDOCK	structure solution	
HADDOCK	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section [7](#) of this report.

Chemical shift file(s)	2ma9_cs.cif
Number of chemical shift lists	3
Total number of shifts	827
Number of shifts mapped to atoms	827
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	27%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [\(i\)](#)

6.1 Standard geometry [\(i\)](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	257	59	282	8±3
2	B	920	204	908	10±2
3	C	718	149	714	10±4
All	All	37900	8240	38080	434

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:23:LYS:HD3	1:A:24:PRO:HD2	0.92	1.41	15	11
1:A:23:LYS:HD2	1:A:35:ASP:HB3	0.90	1.43	2	6
1:A:23:LYS:HG3	1:A:35:ASP:HB3	0.90	1.41	11	14
2:B:17:ASP:HB3	3:C:14:LYS:HD2	0.83	1.50	4	13
1:A:23:LYS:HG2	1:A:24:PRO:HD2	0.82	1.50	8	2
2:B:71:ALA:HB2	2:B:111:SER:HB3	0.78	1.54	4	7
1:A:25:PRO:HA	1:A:32:LEU:HA	0.76	1.57	12	19
1:A:16:LEU:HA	3:C:77:ILE:HG22	0.76	1.57	20	9
2:B:27:LEU:HD11	2:B:57:LEU:HD21	0.75	1.58	12	17
2:B:101:ASP:HA	2:B:104:LYS:HD3	0.74	1.59	9	1
3:C:2:LYS:HE3	3:C:10:GLU:HB3	0.71	1.61	7	2
3:C:65:TYR:HA	3:C:69:SER:HB2	0.70	1.64	4	8

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:22:ILE:HD11	3:C:83:LEU:HD23	0.69	1.63	17	4
2:B:5:LEU:HG	2:B:16:THR:HB	0.69	1.65	9	18
2:B:24:VAL:HB	2:B:53:ASP:HA	0.66	1.68	12	5
2:B:4:PHE:H	2:B:67:ALA:HB1	0.65	1.48	4	7
1:A:21:GLN:HB2	1:A:34:GLU:HG3	0.65	1.68	15	3
1:A:8:LEU:HD21	3:C:85:LEU:HB3	0.64	1.70	11	14
2:B:106:GLN:HG2	2:B:117:VAL:HA	0.63	1.69	9	2
2:B:104:LYS:H	2:B:105:PRO:HD2	0.62	1.52	9	4
1:A:29:VAL:HG13	2:B:102:VAL:HB	0.62	1.71	5	2
2:B:107:ASP:HB2	2:B:118:GLN:HE22	0.61	1.54	4	4
2:B:23:THR:HA	2:B:56:THR:HA	0.61	1.70	8	17
2:B:32:GLU:HG3	2:B:39:PRO:HD3	0.60	1.72	6	7
1:A:18:LYS:HD2	3:C:80:GLU:HB2	0.60	1.72	9	2
2:B:28:LYS:HB3	2:B:39:PRO:HB3	0.60	1.71	12	3
3:C:72:ILE:HB	3:C:73:PRO:HD2	0.59	1.75	12	11
2:B:92:PRO:HA	2:B:95:SER:O	0.59	1.98	16	17
1:A:23:LYS:HD3	1:A:33:THR:HB	0.59	1.75	2	2
2:B:14:ILE:HG12	3:C:12:ILE:HD12	0.58	1.74	17	1
3:C:4:ILE:HD11	3:C:41:GLU:HB3	0.58	1.73	2	4
1:A:16:LEU:HD23	3:C:77:ILE:HA	0.58	1.73	5	1
2:B:111:SER:HB2	2:B:114:GLU:HG3	0.58	1.76	6	1
1:A:16:LEU:HB3	3:C:82:ALA:HB3	0.58	1.73	12	6
1:A:6:GLY:HA3	3:C:27:MET:SD	0.57	2.39	4	6
3:C:2:LYS:HB3	3:C:41:GLU:HG2	0.57	1.75	6	1
3:C:74:GLU:HG3	3:C:75:PHE:N	0.57	2.15	17	3
1:A:16:LEU:HG	3:C:77:ILE:HA	0.57	1.75	9	2
2:B:107:ASP:HB3	2:B:116:ALA:HB3	0.56	1.78	14	1
2:B:10:HIS:HB2	2:B:90:ILE:HG12	0.55	1.78	20	7
1:A:16:LEU:HB2	3:C:82:ALA:HB2	0.54	1.79	5	1
3:C:78:ALA:HB3	3:C:79:PRO:HD3	0.54	1.80	9	13
2:B:43:ARG:HB2	2:B:78:ALA:HB3	0.54	1.80	4	2
1:A:23:LYS:HB3	1:A:33:THR:HB	0.54	1.80	10	5
3:C:15:ARG:O	3:C:19:LEU:HG	0.53	2.03	11	2
3:C:27:MET:HG3	3:C:91:PHE:HA	0.53	1.78	18	4
2:B:9:ARG:HB3	2:B:12:THR:HB	0.53	1.80	7	1
3:C:27:MET:SD	3:C:30:GLY:HA3	0.52	2.44	5	1
3:C:2:LYS:HD2	3:C:10:GLU:HB3	0.52	1.81	12	2
1:A:18:LYS:HE3	1:A:20:LYS:HD2	0.51	1.82	5	1
3:C:44:PHE:HB3	3:C:47:ILE:HD12	0.50	1.83	11	5
2:B:103:MET:HA	2:B:106:GLN:NE2	0.49	2.23	5	5
2:B:6:MET:HG3	2:B:72:PRO:HB2	0.49	1.85	1	7

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:18:LYS:HD2	3:C:80:GLU:HB3	0.49	1.84	1	2
2:B:104:LYS:HB3	2:B:105:PRO:HD3	0.49	1.83	7	8
1:A:32:LEU:HG	3:C:84:GLU:HG3	0.48	1.86	17	2
2:B:17:ASP:CB	3:C:14:LYS:HD2	0.48	2.36	17	1
3:C:2:LYS:HE2	3:C:4:ILE:HG12	0.48	1.84	8	2
1:A:11:LEU:HB3	3:C:58:TYR:CE2	0.48	2.42	10	4
1:A:26:LEU:H	1:A:26:LEU:HD23	0.48	1.69	10	1
1:A:22:ILE:HG23	1:A:32:LEU:HD22	0.48	1.84	17	1
2:B:17:ASP:HB3	3:C:14:LYS:CD	0.48	2.39	12	2
3:C:3:LEU:HB2	3:C:11:PHE:HB2	0.48	1.83	14	4
1:A:23:LYS:HD2	1:A:35:ASP:CB	0.47	2.39	17	1
1:A:7:SER:OG	3:C:26:ALA:HA	0.47	2.10	2	1
1:A:26:LEU:HG	1:A:31:LYS:O	0.47	2.09	9	1
2:B:103:MET:HA	2:B:106:GLN:HE22	0.47	1.69	10	1
2:B:25:PHE:HB2	2:B:53:ASP:HB3	0.47	1.86	8	2
2:B:111:SER:HB2	2:B:114:GLU:HB2	0.46	1.87	10	2
2:B:32:GLU:O	2:B:36:LYS:HA	0.46	2.10	19	2
2:B:116:ALA:HA	3:C:81:ILE:HD11	0.46	1.87	2	1
1:A:9:GLN:HA	3:C:86:LEU:HG	0.46	1.87	12	1
3:C:38:GLU:O	3:C:42:VAL:HG22	0.46	2.11	12	6
2:B:100:PRO:HB2	2:B:103:MET:HB2	0.46	1.88	8	1
1:A:16:LEU:CD2	3:C:80:GLU:HA	0.45	2.41	20	5
2:B:90:ILE:O	2:B:92:PRO:HD3	0.45	2.12	13	2
3:C:3:LEU:HB3	3:C:44:PHE:HE2	0.45	1.71	12	1
1:A:31:LYS:HD2	2:B:117:VAL:HB	0.45	1.88	13	1
1:A:18:LYS:HE3	1:A:20:LYS:CD	0.44	2.42	14	2
3:C:74:GLU:HG3	3:C:75:PHE:H	0.44	1.72	17	1
2:B:14:ILE:HA	3:C:12:ILE:HB	0.44	1.88	10	5
1:A:32:LEU:CD2	3:C:84:GLU:HA	0.44	2.42	17	5
1:A:18:LYS:HD3	3:C:83:LEU:CB	0.44	2.42	2	2
2:B:104:LYS:H	2:B:105:PRO:CD	0.44	2.24	9	3
3:C:15:ARG:HA	3:C:18:ALA:HB3	0.44	1.89	16	2
1:A:32:LEU:HD23	3:C:84:GLU:HA	0.44	1.89	4	5
2:B:107:ASP:HB2	2:B:118:GLN:NE2	0.44	2.28	16	1
1:A:8:LEU:HA	1:A:11:LEU:HB2	0.44	1.90	6	1
1:A:18:LYS:HB3	3:C:83:LEU:HD22	0.44	1.89	12	1
1:A:17:ILE:O	1:A:19:PRO:HD3	0.44	2.12	13	2
2:B:7:ILE:HD13	2:B:75:VAL:HB	0.44	1.90	20	1
2:B:46:LYS:HD2	2:B:62:PHE:CZ	0.44	2.48	7	1
3:C:2:LYS:HD3	3:C:41:GLU:OE2	0.44	2.13	1	1
3:C:7:ASP:OD2	3:C:48:PRO:HA	0.44	2.12	5	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:111:SER:O	2:B:114:GLU:HB2	0.44	2.12	6	1
1:A:22:ILE:HD11	3:C:83:LEU:CD2	0.44	2.41	17	1
1:A:30:ARG:CZ	3:C:48:PRO:HB3	0.43	2.43	2	2
2:B:3:VAL:CG2	2:B:20:GLU:HG3	0.43	2.43	9	1
3:C:20:THR:O	3:C:24:ILE:HG13	0.43	2.13	16	1
2:B:46:LYS:HB3	2:B:51:LEU:HD21	0.43	1.89	12	1
3:C:2:LYS:HD3	3:C:12:ILE:HD11	0.43	1.89	6	1
2:B:57:LEU:O	2:B:62:PHE:HB2	0.43	2.14	18	1
1:A:25:PRO:CA	1:A:32:LEU:HA	0.43	2.38	12	1
1:A:7:SER:HB3	3:C:26:ALA:HA	0.43	1.91	17	4
1:A:8:LEU:HD23	3:C:85:LEU:HB3	0.42	1.91	18	1
3:C:88:ALA:HB1	3:C:91:PHE:HD2	0.42	1.75	16	1
3:C:51:VAL:HB	3:C:85:LEU:HD21	0.42	1.92	12	1
2:B:70:GLN:HG3	3:C:61:TYR:CD1	0.42	2.50	2	1
2:B:71:ALA:HB1	2:B:112:ALA:H	0.41	1.75	14	1
1:A:7:SER:HB3	3:C:25:LYS:O	0.41	2.15	20	1
1:A:16:LEU:HB3	3:C:82:ALA:CB	0.41	2.45	7	2
3:C:2:LYS:HD3	3:C:41:GLU:HG2	0.41	1.92	7	1
3:C:69:SER:O	3:C:72:ILE:HG12	0.41	2.15	9	1
2:B:91:GLU:HG2	2:B:93:PHE:H	0.41	1.76	5	1
1:A:25:PRO:HD3	3:C:87:MET:SD	0.41	2.56	12	1
2:B:106:GLN:HA	2:B:118:GLN:OE1	0.41	2.15	8	1
1:A:11:LEU:HB3	3:C:58:TYR:HE2	0.41	1.76	20	1
1:A:18:LYS:HD2	3:C:80:GLU:CB	0.41	2.46	1	1
1:A:32:LEU:HB3	3:C:87:MET:SD	0.40	2.56	7	1
2:B:38:PRO:HA	2:B:39:PRO:HD3	0.40	1.81	7	1
3:C:59:PHE:O	3:C:63:VAL:HG23	0.40	2.17	16	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	30/36 (83%)	23±1 (77±5%)	4±2 (12±5%)	3±1 (10±3%)	1 10
2	B	116/118 (98%)	110±1 (95±1%)	4±1 (4±1%)	2±1 (1±1%)	20 66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	C	89/91 (98%)	78±2 (88±2%)	10±2 (11±2%)	2±1 (2±1%)	15 58
All	All	4700/4900 (96%)	4224 (90%)	353 (8%)	123 (3%)	11 47

All 16 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	36	ARG	20
1	A	29	VAL	19
2	B	10	HIS	18
3	C	77	ILE	16
1	A	35	ASP	12
2	B	104	LYS	11
3	C	78	ALA	5
1	A	30	ARG	4
1	A	24	PRO	4
3	C	82	ALA	4
3	C	67	ASN	3
3	C	31	PRO	2
1	A	19	PRO	2
3	C	89	ALA	1
3	C	32	GLY	1
2	B	2	ASP	1

6.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	28/32 (88%)	21±2 (76±6%)	7±2 (24±6%)	3 28
2	B	103/103 (100%)	94±2 (91±2%)	9±2 (9±2%)	16 61
3	C	80/80 (100%)	72±2 (90±2%)	8±2 (10±2%)	14 59
All	All	4220/4300 (98%)	3743 (89%)	477 (11%)	12 55

All 74 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	36	ARG	20
3	C	70	THR	20
1	A	7	SER	19
2	B	5	LEU	19
1	A	28	SER	19
2	B	66	THR	18
3	C	66	THR	17
3	C	50	HIS	16
3	C	68	SER	16
2	B	70	GLN	15
3	C	9	HIS	15
1	A	26	LEU	15
2	B	74	THR	15
2	B	21	SER	14
2	B	110	SER	14
1	A	37	TRP	13
1	A	23	LYS	13
1	A	33	THR	12
2	B	113	ASN	12
1	A	16	LEU	11
3	C	29	SER	11
3	C	91	PHE	9
3	C	20	THR	8
2	B	2	ASP	8
3	C	6	SER	8
1	A	35	ASP	7
2	B	64	SER	7
2	B	107	ASP	6
2	B	83	ASP	6
2	B	63	THR	6
3	C	86	LEU	6
3	C	80	GLU	4
3	C	38	GLU	4
2	B	94	SER	4
3	C	53	SER	3
3	C	5	SER	3
2	B	118	GLN	3
2	B	111	SER	3
3	C	69	SER	3
2	B	32	GLU	3
2	B	47	ASP	3
2	B	1	MET	3
2	B	53	ASP	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	B	37	ARG	2
2	B	40	ASP	2
2	B	68	ARG	2
2	B	84	THR	2
2	B	13	THR	2
3	C	7	ASP	2
3	C	39	THR	2
2	B	12	THR	2
1	A	29	VAL	2
3	C	49	SER	2
2	B	108	SER	2
2	B	16	THR	2
3	C	23	THR	1
2	B	89	CYS	1
3	C	25	LYS	1
3	C	37	ASN	1
2	B	25	PHE	1
3	C	21	SER	1
3	C	42	VAL	1
1	A	34	GLU	1
3	C	34	PHE	1
2	B	48	ASP	1
2	B	115	GLN	1
1	A	30	ARG	1
3	C	33	GLN	1
1	A	19	PRO	1
2	B	82	ASP	1
2	B	114	GLU	1
3	C	71	GLU	1
3	C	67	ASN	1
2	B	101	ASP	1

6.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

6.7 Other polymers [\(i\)](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

7 Chemical shift validation i

The completeness of assignment taking into account all chemical shift lists is 27% for the well-defined parts and 27% for the entire structure.

7.1 Chemical shift list 1

File name: 2ma9_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping i

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	152
Number of shifts mapped to atoms	152
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing i

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	33	0.45 ± 0.89	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	32	0.60 ± 0.74	None needed (imprecise)
$^{13}\text{C}'$	0	—	—
^{15}N	26	-0.34 ± 0.67	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments i

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 5%, i.e. 137 atoms were assigned a chemical shift out of a possible 2979. 0 out of 36 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	109/1169 (9%)	55/464 (12%)	30/482 (6%)	24/223 (11%)
Sidechain	28/1613 (2%)	0/951 (0%)	28/594 (5%)	0/68 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	0/197 (0%)	0/107 (0%)	0/85 (0%)	0/5 (0%)
Overall	137/2979 (5%)	55/1522 (4%)	58/1161 (5%)	24/296 (8%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 5%, i.e. 137 atoms were assigned a chemical shift out of a possible 2979. 0 out of 36 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	109/1169 (9%)	55/464 (12%)	30/482 (6%)	24/223 (11%)
Sidechain	28/1613 (2%)	0/951 (0%)	28/594 (5%)	0/68 (0%)
Aromatic	0/197 (0%)	0/107 (0%)	0/85 (0%)	0/5 (0%)
Overall	137/2979 (5%)	55/1522 (4%)	58/1161 (5%)	24/296 (8%)

7.1.4 Statistically unusual chemical shifts [\(i\)](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	20	LYS	CB	41.78	41.68 – 23.88	5.1

7.1.5 Random Coil Index (RCI) plots [\(i\)](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: 2ma9_cs.cif

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping [\(i\)](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	468
Number of shifts mapped to atoms	468
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.2.2 Chemical shift referencing [\(i\)](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	106	-0.13 \pm 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	46	-0.21 \pm 0.18	None needed (< 0.5 ppm)
$^{13}\text{C}'$	97	0.51 \pm 0.16	Should be applied
^{15}N	91	-0.79 \pm 0.37	Should be applied

7.2.3 Completeness of resonance assignments [\(i\)](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 16%, i.e. 463 atoms were assigned a chemical shift out of a possible 2979. 0 out of 36 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	417/1169 (36%)	123/464 (27%)	203/482 (42%)	91/223 (41%)
Sidechain	46/1613 (3%)	0/951 (0%)	46/594 (8%)	0/68 (0%)
Aromatic	0/197 (0%)	0/107 (0%)	0/85 (0%)	0/5 (0%)
Overall	463/2979 (16%)	123/1522 (8%)	249/1161 (21%)	91/296 (31%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 16%, i.e. 463 atoms were assigned a chemical shift out of a possible 2979. 0 out of 36 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	417/1169 (36%)	123/464 (27%)	203/482 (42%)	91/223 (41%)
Sidechain	46/1613 (3%)	0/951 (0%)	46/594 (8%)	0/68 (0%)
Aromatic	0/197 (0%)	0/107 (0%)	0/85 (0%)	0/5 (0%)
Overall	463/2979 (16%)	123/1522 (8%)	249/1161 (21%)	91/296 (31%)


7.2.4 Statistically unusual chemical shifts [\(i\)](#)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots [\(i\)](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:



7.3 Chemical shift list 3

File name: 2ma9_cs.cif

Chemical shift list name: assigned_chem_shift_list_3

7.3.1 Bookkeeping [\(i\)](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	207
Number of shifts mapped to atoms	207

Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.3.2 Chemical shift referencing [\(i\)](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	53	-0.57 ± 0.67	None needed (imprecise)
$^{13}\text{C}_\beta$	16	2.04 ± 0.35	Should be applied
$^{13}\text{C}'$	34	-0.03 ± 0.28	None needed (< 0.5 ppm)
^{15}N	43	-0.23 ± 0.29	None needed (< 0.5 ppm)

7.3.3 Completeness of resonance assignments [\(i\)](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 7%, i.e. 204 atoms were assigned a chemical shift out of a possible 2979. 0 out of 36 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	188/1169 (16%)	58/464 (12%)	87/482 (18%)	43/223 (19%)
Sidechain	16/1613 (1%)	0/951 (0%)	16/594 (3%)	0/68 (0%)
Aromatic	0/197 (0%)	0/107 (0%)	0/85 (0%)	0/5 (0%)
Overall	204/2979 (7%)	58/1522 (4%)	103/1161 (9%)	43/296 (15%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 7%, i.e. 204 atoms were assigned a chemical shift out of a possible 2979. 0 out of 36 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	188/1169 (16%)	58/464 (12%)	87/482 (18%)	43/223 (19%)
Sidechain	16/1613 (1%)	0/951 (0%)	16/594 (3%)	0/68 (0%)
Aromatic	0/197 (0%)	0/107 (0%)	0/85 (0%)	0/5 (0%)
Overall	204/2979 (7%)	58/1522 (4%)	103/1161 (9%)	43/296 (15%)

7.3.4 Statistically unusual chemical shifts [\(i\)](#)

There are no statistically unusual chemical shifts.

7.3.5 Random Coil Index (RCI) plots [\(i\)](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain C:

