



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:42 PM GMT

PDB ID : 4MGH  
Title : Importance of Hydrophobic Cavities in Allosteric Regulation of Formylglycinamide Synthetase: Insight from Xenon Trapping and Statistical Coupling Analysis  
Authors : Tanwar, A.S.; Goyal, V.D.; Choudhary, D.; Panjekar, S.; Anand, R.  
Deposited on : 2013-08-28  
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3152 (2.70-2.62)
Clashscore	102246	3524 (2.70-2.62)
Ramachandran outliers	100387	3469 (2.70-2.62)
Sidechain outliers	100360	3469 (2.70-2.62)
RSRZ outliers	91569	3161 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1303	<div> <div>2%</div> <div>84%</div> <div>12% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	1305	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	A	1310	-	-	-	X
5	SO4	A	1315	-	-	-	X
6	ACT	A	1320	-	-	X	X
6	ACT	A	1322	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 10514 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphoribosylformylglycinamide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1284	9879	6204	1757	1870	48	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	EXPRESSION TAG	UNP P74881
A	-6	LEU	-	EXPRESSION TAG	UNP P74881
A	-5	VAL	-	EXPRESSION TAG	UNP P74881
A	-4	PRO	-	EXPRESSION TAG	UNP P74881
A	-3	ARG	-	EXPRESSION TAG	UNP P74881
A	-2	GLY	-	EXPRESSION TAG	UNP P74881
A	-1	SER	-	EXPRESSION TAG	UNP P74881
A	0	HIS	-	EXPRESSION TAG	UNP P74881

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Mg	0	0
			4	4		

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mn	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is XENON (three-letter code: XE) (formula: Xe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	4	Total	Xe	0	0
			4	4		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	518	Total	O	0	0
			518	518		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoribosylformylglycinamide synthase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	146.45Å 146.45Å 141.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.99 – 2.65 19.99 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.99-2.65) 100.0 (19.99-2.65)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.42 (at 2.67Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.153 , 0.203 0.161 , 0.206	Depositor DCC
$R_{free}$ test set	1176 reflections (2.41%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.1	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 43.5	EDS
Estimated twinning fraction	0.018 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 49942 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10514	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP, MN, XE, CYG, SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	2/10076 (0.0%)	0.91	22/13675 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1152	GLU	CG-CD	5.62	1.60	1.51
1	A	211	GLN	CG-CD	5.26	1.63	1.51

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	487	ARG	NE-CZ-NH2	-11.21	114.69	120.30
1	A	932	ARG	NE-CZ-NH1	9.47	125.03	120.30
1	A	382	ARG	NE-CZ-NH1	9.02	124.81	120.30
1	A	332	LEU	CA-CB-CG	7.30	132.09	115.30
1	A	1115	ARG	NE-CZ-NH2	-6.75	116.92	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9879	0	9656	107	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	27	0	12	0	0
3	A	4	0	0	0	0
4	A	1	0	0	0	0
5	A	65	0	0	4	0
6	A	16	0	12	5	0
7	A	4	0	0	0	0
8	A	518	0	0	18	0
All	All	10514	0	9680	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:782:LYS:NZ	8:A:1548:HOH:O	1.92	1.02
1:A:80:ARG:CD	8:A:1716:HOH:O	2.19	0.88
1:A:820:SER:H	1:A:930:GLN:HE22	1.30	0.79
1:A:829:ILE:HD13	1:A:829:ILE:N	2.00	0.77
1:A:218:ARG:CZ	1:A:220:LYS:HE2	2.15	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1279/1303 (98%)	1232 (96%)	44 (3%)	3 (0%)	52 77

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	623	GLY
1	A	969	ASN
1	A	594	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1023/1040 (98%)	960 (94%)	63 (6%)	23 45

5 of 63 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	429	GLU
1	A	596	GLN
1	A	1190	MET
1	A	445	ASN
1	A	508	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	584	GLN
1	A	746	HIS
1	A	1125	HIS
1	A	585	HIS
1	A	592	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CYG	A	1135	1	10,14,15	1.92	2 (20%)	6,17,19	7.89	5 (83%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CYG	A	1135	1	-	0/10/16/18	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1135	CYG	OE2-CD1	3.57	1.26	1.21
1	A	1135	CYG	CG1-CD1	4.69	1.56	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1135	CYG	OE2-CD1-CG1	-15.58	113.22	123.94
1	A	1135	CYG	CB1-CG1-CD1	-5.12	105.56	113.12
1	A	1135	CYG	O-C-CA	-2.33	119.41	125.49
1	A	1135	CYG	OE2-CD1-SG	2.40	124.74	122.83
1	A	1135	CYG	CG1-CD1-SG	9.64	122.03	113.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 9 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	ADP	A	1301	3	22,29,29	1.18	4 (18%)	27,45,45	1.83	5 (18%)
5	SO4	A	1307	-	4,4,4	0.55	0	6,6,6	0.41	0
5	SO4	A	1308	-	4,4,4	0.76	0	6,6,6	0.81	0
5	SO4	A	1309	-	4,4,4	0.68	0	6,6,6	0.56	0
5	SO4	A	1310	-	4,4,4	0.40	0	6,6,6	1.17	1 (16%)
5	SO4	A	1311	-	4,4,4	0.52	0	6,6,6	0.41	0
5	SO4	A	1312	-	4,4,4	0.68	0	6,6,6	0.24	0
5	SO4	A	1313	-	4,4,4	1.01	0	6,6,6	0.70	0
5	SO4	A	1314	-	4,4,4	0.34	0	6,6,6	0.38	0
5	SO4	A	1315	-	4,4,4	0.64	0	6,6,6	0.56	0
5	SO4	A	1316	-	4,4,4	0.29	0	6,6,6	0.14	0
5	SO4	A	1317	-	4,4,4	0.55	0	6,6,6	0.35	0
5	SO4	A	1318	-	4,4,4	0.55	0	6,6,6	1.16	1 (16%)
5	SO4	A	1319	-	4,4,4	0.61	0	6,6,6	0.60	0
6	ACT	A	1320	-	1,3,3	3.85	1 (100%)	0,3,3	0.00	-
6	ACT	A	1321	-	1,3,3	3.31	1 (100%)	0,3,3	0.00	-
6	ACT	A	1322	-	1,3,3	5.56	1 (100%)	0,3,3	0.00	-
6	ACT	A	1323	-	1,3,3	0.21	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1301	3	-	0/12/32/32	0/3/3/3
5	SO4	A	1307	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1308	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1309	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1310	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1311	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1312	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1313	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1314	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1315	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1316	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1317	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1318	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1319	-	-	0/0/0/0	0/0/0/0
6	ACT	A	1320	-	-	0/0/0/0	0/0/0/0
6	ACT	A	1321	-	-	0/0/0/0	0/0/0/0
6	ACT	A	1322	-	-	0/0/0/0	0/0/0/0
6	ACT	A	1323	-	-	0/0/0/0	0/0/0/0

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1301	ADP	C5-N7	-2.17	1.32	1.39
2	A	1301	ADP	PB-O1B	2.16	1.58	1.51
2	A	1301	ADP	C5-C4	2.49	1.46	1.40
2	A	1301	ADP	O4'-C1'	2.51	1.44	1.41
6	A	1321	ACT	CH3-C	3.31	1.53	1.48

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1301	ADP	N3-C2-N1	-4.45	125.49	128.89
2	A	1301	ADP	PA-O3A-PB	-3.52	120.87	132.67
2	A	1301	ADP	C2'-C1'-N9	-3.39	109.12	114.29
2	A	1301	ADP	O3A-PA-O5'	-2.92	95.20	102.94
5	A	1318	SO4	O4-S-O3	-2.70	98.02	108.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1307	SO4	1	0
5	A	1310	SO4	1	0
5	A	1315	SO4	1	0
5	A	1317	SO4	1	0
6	A	1320	ACT	4	0
6	A	1323	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1283/1303 (98%)	-0.54	22 (1%) 73 72	9, 18, 37, 76	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	25	ALA	4.9
1	A	789	ASN	4.4
1	A	-1	SER	4.1
1	A	276	ASN	4.0
1	A	117	SER	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CYG	A	1135	15/16	0.97	0.10	-	11,13,14,15	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	ACT	A	1320	4/4	0.80	0.45	20.47	47,50,52,53	0
6	ACT	A	1322	4/4	0.83	0.26	10.83	21,22,26,26	0
5	SO4	A	1315	5/5	0.95	0.30	5.28	47,51,53,58	0
3	MG	A	1305	1/1	0.98	0.20	4.95	2,2,2,2	0
5	SO4	A	1310	5/5	0.99	0.17	2.76	23,25,26,28	0
5	SO4	A	1309	5/5	0.98	0.22	1.56	27,28,29,30	0
5	SO4	A	1307	5/5	0.98	0.17	0.71	29,32,34,34	0
5	SO4	A	1314	5/5	0.99	0.12	0.62	25,25,26,27	0
5	SO4	A	1308	5/5	0.99	0.12	-0.02	20,20,23,24	0
2	ADP	A	1301	27/27	0.98	0.09	-1.14	7,9,10,11	0
3	MG	A	1302	1/1	0.95	0.07	-1.97	15,15,15,15	0
3	MG	A	1303	1/1	0.96	0.05	-2.18	15,15,15,15	0
7	XE	A	1326	1/1	0.99	0.10	-2.41	29,29,29,29	1
7	XE	A	1324	1/1	1.00	0.06	-3.05	15,15,15,15	0
7	XE	A	1325	1/1	1.00	0.07	-3.10	21,21,21,21	1
5	SO4	A	1311	5/5	0.99	0.15	-	26,29,31,31	0
5	SO4	A	1313	5/5	0.86	0.38	-	46,48,59,60	0
3	MG	A	1304	1/1	0.96	0.07	-	16,16,16,16	0
6	ACT	A	1321	4/4	0.96	0.10	-	18,20,21,22	0
5	SO4	A	1317	5/5	0.98	0.22	-	40,41,43,47	0
6	ACT	A	1323	4/4	0.91	0.19	-	33,33,36,40	0
5	SO4	A	1319	5/5	0.99	0.24	-	28,30,32,33	0
4	MN	A	1306	1/1	0.98	0.09	-	47,47,47,47	0
7	XE	A	1327	1/1	1.00	0.03	-	24,24,24,24	1
5	SO4	A	1312	5/5	0.96	0.37	-	46,46,50,51	0
5	SO4	A	1318	5/5	0.97	0.31	-	34,35,43,46	0
5	SO4	A	1316	5/5	0.96	0.17	-	14,14,15,15	5

## 6.5 Other polymers ⓘ

There are no such residues in this entry.