



# Full wwPDB NMR Structure Validation Report i

Apr 27, 2016 – 03:03 AM BST

PDB ID : 2MKJ  
Title : Solution structure of tandem RRM domains of cytoplasmic polyadenylation element binding protein 4 (CPEB4) in free state  
Authors : Afroz, T.; Skrisovska, L.; Belloc, E.; Boixet, J.G.; Mendez, R.; Allain, F.H.-T.  
Deposited on : 2014-02-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:


Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbit	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*SOLUTION NMR*


The overall completeness of chemical shifts assignment is 65%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$



## 2 Ensemble composition and analysis

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:62-A:95, A:109-A:140, A:145-A:254 (176)	1.01	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: Inconsistent models

### 3 Entry composition [\(i\)](#)

There is only 1 type of molecule in this entry. The entry contains 3182 atoms, of which 1582 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Cytoplasmic polyadenylation element-binding protein 4.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	202	3182	1023	1582	273	298	6	0

## 4 Residue-property plots [\(i\)](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4



#### 4.2.2 Score per residue for model 2 (medoid)


- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4






#### 4.2.3 Score per residue for model 3

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.4 Score per residue for model 4

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.5 Score per residue for model 5


- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4



#### 4.2.6 Score per residue for model 6

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4





#### 4.2.7 Score per residue for model 7

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4



Chain A: 80% 7% 13%




#### 4.2.8 Score per residue for model 8

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4


Chain A: 78% 6% • 13%






#### 4.2.11 Score per residue for model 11

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.12 Score per residue for model 12

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.13 Score per residue for model 13

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.14 Score per residue for model 14

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.15 Score per residue for model 15

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.16 Score per residue for model 16

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.17 Score per residue for model 17

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.18 Score per residue for model 18

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4




#### 4.2.19 Score per residue for model 19

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4



#### 4.2.20 Score per residue for model 20

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 4



## 5 Refinement protocol and experimental data overview i

The models were refined using the following method: *simulated annealing*.

Of the 30 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	
CYANA	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section [7](#) of this report.

Chemical shift file(s)	2mkj_cs.str
Number of chemical shift lists	1
Total number of shifts	1720
Number of shifts mapped to atoms	1720
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	65%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [\(i\)](#)

### 6.1 Standard geometry [\(i\)](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1392	1381	1381	5±3
All	All	27840	27620	27620	99

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:156:LEU:H	1:A:156:LEU:HD13	0.77	1.39	10	7
1:A:156:LEU:HD13	1:A:156:LEU:H	0.73	1.42	8	5
1:A:196:LEU:CD1	1:A:197:TYR:H	0.63	2.07	11	2
1:A:156:LEU:CD1	1:A:156:LEU:H	0.62	2.08	20	4
1:A:156:LEU:H	1:A:156:LEU:CD1	0.57	2.12	10	2
1:A:156:LEU:HD13	1:A:156:LEU:N	0.57	2.14	16	2
1:A:156:LEU:N	1:A:156:LEU:HD13	0.53	2.16	17	1
1:A:189:LEU:HD23	1:A:189:LEU:C	0.52	2.24	3	4
1:A:88:PHE:CD1	1:A:115:PHE:CZ	0.52	2.98	11	4
1:A:161:PHE:CZ	1:A:203:ALA:HB3	0.50	2.40	18	3
1:A:88:PHE:CE1	1:A:115:PHE:CZ	0.50	3.00	5	3
1:A:196:LEU:HD12	1:A:197:TYR:H	0.49	1.65	11	5
1:A:154:TRP:CG	1:A:155:ASN:N	0.48	2.81	1	2
1:A:196:LEU:HD13	1:A:197:TYR:CD2	0.48	2.43	1	4

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:189:LEU:HD13	1:A:189:LEU:C	0.47	2.30	2	2
1:A:241:HIS:CG	1:A:242:GLY:N	0.47	2.83	11	1
1:A:94:ASP:CB	1:A:112:PHE:H	0.46	2.22	20	1
1:A:192:ILE:O	1:A:196:LEU:CD2	0.45	2.64	18	1
1:A:173:LYS:HE2	1:A:173:LYS:HA	0.45	1.87	18	1
1:A:223:SER:HA	1:A:228:TYR:CD2	0.45	2.46	8	3
1:A:88:PHE:CZ	1:A:124:LEU:HD12	0.45	2.47	5	1
1:A:182:ARG:N	1:A:183:PRO:CD	0.44	2.81	9	3
1:A:154:TRP:CD1	1:A:155:ASN:N	0.44	2.85	19	1
1:A:156:LEU:N	1:A:156:LEU:CD1	0.44	2.81	12	2
1:A:63:TYR:N	1:A:63:TYR:CD1	0.43	2.85	15	1
1:A:64:SER:HB2	1:A:156:LEU:HD12	0.43	1.90	17	2
1:A:223:SER:HA	1:A:228:TYR:CG	0.43	2.48	4	1
1:A:94:ASP:HA	1:A:112:PHE:O	0.43	2.13	20	1
1:A:161:PHE:CZ	1:A:203:ALA:CB	0.43	3.01	17	2
1:A:88:PHE:CE2	1:A:115:PHE:CE2	0.43	3.07	8	1
1:A:184:LEU:CD2	1:A:186:ALA:H	0.43	2.27	19	1
1:A:246:LYS:H	1:A:246:LYS:HE2	0.43	1.74	11	1
1:A:189:LEU:C	1:A:189:LEU:HD23	0.43	2.34	12	2
1:A:156:LEU:CD1	1:A:156:LEU:N	0.42	2.82	7	3
1:A:241:HIS:CG	1:A:242:GLY:H	0.42	2.32	11	1
1:A:68:PHE:C	1:A:68:PHE:CD1	0.42	2.92	10	1
1:A:63:TYR:CD1	1:A:63:TYR:N	0.42	2.87	14	1
1:A:180:VAL:N	1:A:181:PRO:CD	0.42	2.82	10	1
1:A:94:ASP:CB	1:A:112:PHE:N	0.42	2.82	20	1
1:A:173:LYS:HA	1:A:173:LYS:HE2	0.42	1.90	5	1
1:A:196:LEU:N	1:A:196:LEU:CD2	0.41	2.84	18	1
1:A:238:GLN:C	1:A:239:LEU:HD22	0.41	2.36	17	1
1:A:161:PHE:CD1	1:A:186:ALA:CB	0.41	3.04	2	1
1:A:156:LEU:HD12	1:A:156:LEU:N	0.41	2.30	7	1
1:A:196:LEU:HD12	1:A:197:TYR:N	0.41	2.30	20	1
1:A:184:LEU:HD12	1:A:184:LEU:C	0.41	2.36	18	1
1:A:124:LEU:CD1	1:A:137:LEU:HD21	0.41	2.46	19	1
1:A:62:ARG:CD	1:A:62:ARG:N	0.41	2.84	3	1
1:A:88:PHE:CG	1:A:115:PHE:CZ	0.41	3.09	11	1
1:A:205:ILE:O	1:A:207:THR:N	0.40	2.54	10	1
1:A:189:LEU:C	1:A:189:LEU:HD13	0.40	2.36	6	1
1:A:161:PHE:CE2	1:A:186:ALA:HB3	0.40	2.52	13	1
1:A:92:ILE:HD12	1:A:92:ILE:N	0.40	2.31	17	1
1:A:129:ILE:HD12	1:A:129:ILE:N	0.40	2.31	15	1

## 6.3 Torsion angles [\(i\)](#)

### 6.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	175/203 (86%)	152±3 (87±2%)	19±4 (11±2%)	4±2 (2±1%)	13 53
All	All	3500/4060 (86%)	3042 (87%)	383 (11%)	75 (2%)	13 53

All 30 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	153	PRO	6
1	A	246	LYS	6
1	A	182	ARG	5
1	A	139	VAL	5
1	A	206	ASP	5
1	A	165	GLY	4
1	A	159	SER	4
1	A	217	ALA	3
1	A	179	GLY	3
1	A	70	GLY	3
1	A	222	PHE	3
1	A	78	GLU	3
1	A	145	LYS	2
1	A	239	LEU	2
1	A	237	VAL	2
1	A	164	ASP	2
1	A	140	SER	2
1	A	235	ARG	2
1	A	247	ARG	2
1	A	219	ARG	1
1	A	86	ARG	1
1	A	89	GLY	1
1	A	110	TYR	1
1	A	94	ASP	1
1	A	243	GLU	1
1	A	166	SER	1
1	A	87	ARG	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	184	LEU	1
1	A	236	PHE	1
1	A	63	TYR	1

### 6.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	150/175 (86%)	136±2 (91±1%)	14±2 (9±1%)	16 61
All	All	3000/3500 (86%)	2726 (91%)	274 (9%)	16 61

All 49 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	76	ILE	20
1	A	228	TYR	20
1	A	81	ILE	20
1	A	196	LEU	17
1	A	156	LEU	17
1	A	191	MET	13
1	A	88	PHE	11
1	A	224	ASN	11
1	A	137	LEU	10
1	A	246	LYS	10
1	A	115	PHE	8
1	A	145	LYS	7
1	A	114	LEU	7
1	A	235	ARG	7
1	A	134	LYS	6
1	A	62	ARG	6
1	A	222	PHE	6
1	A	122	GLN	6
1	A	251	LYS	6
1	A	91	LEU	5
1	A	184	LEU	5
1	A	207	THR	4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	172	ARG	4
1	A	219	ARG	3
1	A	155	ASN	3
1	A	176	PHE	3
1	A	241	HIS	3
1	A	215	LYS	3
1	A	239	LEU	3
1	A	195	ARG	3
1	A	95	TRP	2
1	A	205	ILE	2
1	A	238	GLN	2
1	A	130	GLU	2
1	A	124	LEU	2
1	A	237	VAL	2
1	A	173	LYS	2
1	A	63	TYR	2
1	A	197	TYR	1
1	A	211	LEU	1
1	A	136	TYR	1
1	A	68	PHE	1
1	A	189	LEU	1
1	A	193	MET	1
1	A	182	ARG	1
1	A	151	ILE	1
1	A	66	LYS	1
1	A	147	LYS	1
1	A	188	GLU	1

### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 6.7 Other polymers [\(i\)](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 65% for the well-defined parts and 60% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2mkj\_cs.str

Chemical shift list name: *CPEB4RRM12\_assigned\_chemical\_shifts*

#### 7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1720
Number of shifts mapped to atoms	1720
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	8

#### 7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	137	$2.96 \pm 0.16$	Should be applied
$^{13}\text{C}_\beta$	124	$2.92 \pm 0.10$	Should be applied
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	170	$-0.52 \pm 0.21$	Should be applied

#### 7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 65%, i.e. 1431 atoms were assigned a chemical shift out of a possible 2186. 26 out of 30 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	601/856 (70%)	309/340 (91%)	133/352 (38%)	159/164 (97%)
Sidechain	704/1154 (61%)	439/679 (65%)	257/420 (61%)	8/55 (15%)

*Continued on next page...*

*Continued from previous page...*

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	126/176 (72%)	81/93 (87%)	43/80 (54%)	2/3 (67%)
Overall	1431/2186 (65%)	829/1112 (75%)	433/852 (51%)	169/222 (76%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 60%, i.e. 1504 atoms were assigned a chemical shift out of a possible 2515. 27 out of 31 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	631/978 (65%)	324/388 (84%)	137/404 (34%)	170/186 (91%)
Sidechain	734/1330 (55%)	457/787 (58%)	268/480 (56%)	9/63 (14%)
Aromatic	139/207 (67%)	89/110 (81%)	48/92 (52%)	2/5 (40%)
Overall	1504/2515 (60%)	870/1285 (68%)	453/976 (46%)	181/254 (71%)

#### 7.1.4 Statistically unusual chemical shifts [\(i\)](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	76	ILE	CG1	11.00	36.54 – 18.94	-9.5
1	A	214	PRO	HD3	0.44	5.52 – 1.72	-8.4
1	A	214	PRO	HD2	0.91	5.45 – 1.85	-7.6
1	A	205	ILE	CG1	14.50	36.54 – 18.94	-7.5
1	A	214	PRO	HG3	-0.38	3.56 – 0.26	-6.9
1	A	175	ILE	CG1	17.13	36.54 – 18.94	-6.0
1	A	214	PRO	HG2	0.18	3.48 – 0.38	-5.6
1	A	134	LYS	CE	37.29	46.00 – 37.80	-5.6

#### 7.1.5 Random Coil Index (RCI) plots [\(i\)](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

