



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:42 AM GMT

PDB ID : 3MW0  
Title : X-ray structure of the doubly hydroxylated iron complex-NikA species, NikA1/O2  
Authors : Cavazza, C.; Bochot, C.; Rousselot-Pailley, P.; Carpentier, P.; Cherrier, M.V.; Martin, L.; Marchi-Delapierre, C.; Fontecilla-Camps, J.C.; Menage, S.  
Deposited on : 2010-05-05  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	502	
1	B	502	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	DTD	A	505	-	-	-	X
4	DTD	B	504	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8443 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nickel-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	498	Total 3957	C 2537	N 668	O 742	S 10	6	2	0
1	B	497	Total 3933	C 2521	N 666	O 736	S 10	5	1	0

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Fe 1	0	0
2	A	1	Total 1	Fe 1	0	0

- Molecule 3 is 2-[2-[CARBOXYMETHYL-[(2-HYDROXYPHENYL)METHYL]AMINO]ETHYL-[(2,3-DIHYDROXYPHENYL)METHYL]AMINO]ETHANOIC ACID (three-letter code: BHR) (formula: C<sub>20</sub>H<sub>24</sub>N<sub>2</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			29	20	2	7		
3	B	1	Total	C	N	O	0	0
			29	20	2	7		

- Molecule 4 is DITHIANE DIOL (three-letter code: DTD) (formula:  $C_4H_8O_2S_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			8	4	2	2		
4	B	1	Total	C	O	S	0	0
			8	4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	273	Total	O	0	0
			273	273		
5	B	204	Total	O	0	0
			204	204		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nickel-binding periplasmic protein



- Molecule 1: Nickel-binding periplasmic protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.98Å 94.51Å 125.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.26 – 2.30 47.26 – 2.28	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.26-2.30) 99.8 (47.26-2.28)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.03 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.174 , 0.244 0.256 , 0.299	Depositor DCC
$R_{free}$ test set	2326 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.7	Xtrriage
Anisotropy	0.131	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 32.8	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	0 of 47530 reflections	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	8443	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BHR, FE, DTD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/4068	0.65	1/5544 (0.0%)
1	B	0.54	0/4039	0.63	2/5502 (0.0%)
All	All	0.57	0/8107	0.64	3/11046 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	396	ARG	NE-CZ-NH2	-5.91	117.35	120.30
1	B	396	ARG	NE-CZ-NH1	-5.86	117.37	120.30
1	B	224	LEU	CA-CB-CG	5.47	127.89	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	495[B]	GLU	Mainchain

## 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3957	0	3898	42	1
1	B	3933	0	3862	35	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	29	0	20	0	0
3	B	29	0	20	1	0
4	A	8	0	8	1	0
4	B	8	0	8	0	0
5	A	273	0	0	7	2
5	B	204	0	0	4	0
All	All	8443	0	7816	78	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:GLU:CB	1:A:59:ASP:O	1.66	1.44
1:A:58:GLU:HB2	1:A:59:ASP:O	0.76	0.89
1:A:292:THR:CG2	1:A:294:PHE:O	2.20	0.89
1:B:105:ASN:HB3	5:B:664:HOH:O	1.76	0.85
3:B:503:BHR:H15	5:B:551:HOH:O	1.75	0.83
1:A:10:TRP:HE1	1:A:26:GLN:HE21	1.31	0.75
1:A:58:GLU:CG	1:A:59:ASP:O	2.34	0.74
1:A:445:THR:HG21	5:B:618:HOH:O	1.87	0.72
1:B:438:VAL:HG12	1:B:450:LEU:HB2	1.72	0.72
1:A:188:LYS:H	1:A:188:LYS:HE3	1.55	0.72
1:B:227:ASP:HB3	1:B:284:TYR:CE2	2.26	0.70
1:A:87:ASN:HD21	1:A:142:ILE:H	1.43	0.64
1:A:292:THR:HG21	1:A:294:PHE:O	1.96	0.63
1:A:58:GLU:CD	1:A:59:ASP:O	2.37	0.63
1:A:250:MET:HE1	1:A:397:THR:HG23	1.81	0.62
1:A:446:GLN:HA	1:A:446:GLN:HE21	1.65	0.61
1:A:453:ASP:O	1:A:457:ARG:HG3	2.00	0.61

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:250:MET:CE	1:B:396:ARG:HA	2.31	0.60
1:B:438:VAL:HG12	1:B:450:LEU:CB	2.31	0.60
1:A:131:GLN:HG2	5:A:606:HOH:O	2.02	0.60
1:A:219:GLY:HA3	4:A:505:DTD:H12	1.83	0.59
1:A:10:TRP:HE1	1:A:26:GLN:NE2	2.01	0.59
1:B:414:PRO:O	1:B:415:SER:HB2	2.04	0.58
1:B:10:TRP:HE1	1:B:26:GLN:HE21	1.52	0.57
1:A:292:THR:HG22	1:A:294:PHE:O	2.03	0.57
1:A:453:ASP:OD2	1:A:457:ARG:HD2	2.05	0.57
1:B:215:ASP:OD1	1:B:477:LYS:HE2	2.05	0.56
1:B:442:HIS:CD2	1:B:442:HIS:H	2.21	0.56
1:B:171:LYS:HB3	1:B:174:GLN:HB2	1.88	0.56
1:B:203:THR:CG2	5:B:606:HOH:O	2.54	0.55
1:B:457:ARG:O	1:B:461:GLU:HG3	2.06	0.55
1:B:438:VAL:CG1	1:B:450:LEU:HB3	2.37	0.55
1:B:296:PRO:HB3	1:B:302:ASN:HD22	1.72	0.55
1:B:52:LYS:HE3	1:B:68:ARG:HG2	1.88	0.54
1:A:221:GLU:OE2	1:A:396:ARG:NH2	2.30	0.54
1:A:327:PRO:O	1:A:328:ALA:C	2.47	0.53
1:A:27:MET:HG3	5:A:777:HOH:O	2.09	0.52
1:B:7:THR:HG22	1:B:214:ILE:HG22	1.92	0.52
1:A:327:PRO:O	1:A:328:ALA:O	2.28	0.52
1:B:414:PRO:HA	1:B:419:PHE:CD1	2.45	0.51
1:B:103:LEU:HD21	1:B:133:LEU:HD13	1.92	0.51
1:B:340:LEU:HD13	1:B:369:ALA:HB2	1.93	0.51
1:B:5:GLU:HG2	1:B:193:LYS:HB3	1.92	0.50
1:B:336:ASN:HD22	1:B:336:ASN:N	2.10	0.50
1:A:477:LYS:HE3	5:A:599:HOH:O	2.11	0.50
1:A:58:GLU:OE2	1:A:59:ASP:O	2.29	0.50
1:A:87:ASN:ND2	1:A:142:ILE:H	2.10	0.49
1:A:495[B]:GLU:H	1:A:495[B]:GLU:CD	2.16	0.48
1:B:438:VAL:CG1	1:B:450:LEU:CB	2.90	0.48
1:A:446:GLN:HA	1:A:446:GLN:NE2	2.29	0.48
1:A:265:VAL:O	1:A:269:LEU:HG	2.13	0.47
1:B:250:MET:HE3	1:B:396:ARG:HA	1.95	0.47
1:B:221:GLU:OE1	1:B:396:ARG:NH1	2.45	0.47
1:B:10:TRP:HE1	1:B:26:GLN:NE2	2.13	0.46
1:A:288:GLN:O	1:A:468:SER:HB2	2.16	0.46
1:A:248:THR:HB	1:A:250:MET:CE	2.46	0.45
1:A:279:ILE:HG12	1:A:283:LEU:HD12	1.99	0.45
1:B:19:PRO:HG3	1:B:142:ILE:HB	1.97	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:441:THR:O	1:A:447:ARG:NH1	2.38	0.45
1:A:378:GLU:HG2	1:A:382:TYR:CE2	2.51	0.45
1:B:171:LYS:HD3	1:B:174:GLN:HE21	1.81	0.44
1:B:97:ARG:HB3	1:B:97:ARG:HH21	1.81	0.44
1:A:224:LEU:HD23	1:A:473:MET:CE	2.47	0.43
1:B:282:ALA:O	1:B:283:LEU:HD23	2.19	0.43
1:B:414:PRO:O	1:B:415:SER:CB	2.67	0.42
1:A:447:ARG:NH2	5:A:517:HOH:O	2.48	0.42
1:A:336:ASN:N	5:A:656:HOH:O	2.51	0.42
1:B:427:ASP:OD1	1:B:461:GLU:OE1	2.37	0.42
1:A:296:PRO:HB3	1:A:302:ASN:HD22	1.85	0.41
1:B:328:ALA:HA	1:B:329:GLY:HA2	1.65	0.41
1:A:248:THR:HB	1:A:250:MET:HE3	2.03	0.41
1:B:235:ASN:HA	1:B:236:PRO:HD2	1.88	0.41
1:A:3:PRO:HB2	5:A:736:HOH:O	2.20	0.41
1:B:201:ASP:HA	1:B:202:PRO:HD2	1.96	0.41
1:A:338:GLN:HA	1:A:339:PRO:HD3	1.96	0.40
1:A:445:THR:HG23	5:A:719:HOH:O	2.21	0.40
1:B:385:GLN:HG2	1:B:393:ILE:HD13	2.02	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:700:HOH:O	5:A:770:HOH:O[3_555]	1.96	0.24
1:A:387:ASP:OD2	1:B:460:ASP:OD2[3_545]	2.09	0.11
5:A:654:HOH:O	5:A:671:HOH:O[3_545]	2.10	0.10

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/502 (99%)	481 (97%)	16 (3%)	1 (0%)	52	64
1	B	496/502 (99%)	480 (97%)	16 (3%)	0	100	100
All	All	994/1004 (99%)	961 (97%)	32 (3%)	1 (0%)	56	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	328	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	422/425 (99%)	404 (96%)	18 (4%)	35	47
1	B	417/425 (98%)	397 (95%)	20 (5%)	31	42
All	All	839/850 (99%)	801 (96%)	38 (4%)	35	46

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	127	TYR
1	A	174	GLN
1	A	187	GLU
1	A	188	LYS
1	A	203	THR
1	A	216	LEU
1	A	250	MET
1	A	292	THR
1	A	314	LYS
1	A	319	LEU
1	A	352	LEU
1	A	354	LYS
1	A	373	LEU
1	A	439	LEU
1	A	442[A]	HIS

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	442[B]	HIS
1	A	447	ARG
1	A	473	MET
1	B	23	THR
1	B	50	LEU
1	B	58	GLU
1	B	97	ARG
1	B	127	TYR
1	B	133	LEU
1	B	172	LEU
1	B	203	THR
1	B	216	LEU
1	B	224	LEU
1	B	250	MET
1	B	276	LYS
1	B	313	GLN
1	B	319	LEU
1	B	331	ASP
1	B	340	LEU
1	B	373	LEU
1	B	439	LEU
1	B	472	MET
1	B	498	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	26	GLN
1	A	87	ASN
1	A	131	GLN
1	A	174	GLN
1	A	197	ASN
1	A	302	ASN
1	A	446	GLN
1	B	25	ASN
1	B	26	GLN
1	B	174	GLN
1	B	302	ASN
1	B	313	GLN
1	B	336	ASN
1	B	442	HIS
1	B	482	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	496	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	BHR	A	504	2	24,30,30	0.62	0	34,40,40	1.46	6 (17%)
4	DTD	A	505	-	6,8,8	0.60	0	6,10,10	1.03	0
3	BHR	B	503	2	24,30,30	0.72	0	34,40,40	1.67	6 (17%)
4	DTD	B	504	-	6,8,8	0.85	0	6,10,10	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BHR	A	504	2	-	0/17/21/21	0/2/2/2

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTD	A	505	-	-	0/0/11/11	0/0/1/1
3	BHR	B	503	2	-	0/17/21/21	0/2/2/2
4	DTD	B	504	-	-	0/0/11/11	0/0/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	A	504	BHR	C7-N8-C9	-3.51	103.94	111.28
3	A	504	BHR	C19-N8-C7	-3.48	104.80	110.73
3	B	503	BHR	C20-C19-N8	-2.96	109.15	113.53
3	B	503	BHR	C7-C1-C2	-2.87	114.54	120.16
3	A	504	BHR	C22-C21-N11	-2.35	110.06	113.53
3	A	504	BHR	C7-C1-C2	-2.25	115.76	120.16
3	B	503	BHR	C12-C13-C14	-2.11	116.04	120.16
3	B	503	BHR	C12-N11-C10	-2.03	107.04	111.28
3	A	504	BHR	C12-C13-C18	3.16	123.66	120.48
3	A	504	BHR	C7-C1-C6	3.46	125.22	120.39
3	B	503	BHR	C7-C1-C6	4.68	126.92	120.39
3	B	503	BHR	C12-C13-C18	4.93	125.44	120.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	505	DTD	1	0
3	B	503	BHR	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	498/502 (99%)	0.83	39 (7%) 16 22	9, 19, 33, 45	2 (0%)
1	B	497/502 (99%)	1.11	66 (13%) 4 7	11, 26, 46, 60	1 (0%)
All	All	995/1004 (99%)	0.97	105 (10%) 8 12	9, 22, 42, 60	3 (0%)

All (105) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	328	ALA	7.6
1	B	113	LEU	7.5
1	B	442	HIS	6.8
1	A	2	ALA	6.3
1	A	328	ALA	5.9
1	A	3	PRO	5.9
1	A	59	ASP	5.9
1	B	382	TYR	4.8
1	B	499	PRO	4.4
1	B	58	GLU	4.4
1	A	499	PRO	4.0
1	B	116	THR	4.0
1	B	336	ASN	4.0
1	A	442[A]	HIS	3.9
1	A	172	LEU	3.9
1	B	81	ALA	3.8
1	B	120	ILE	3.6
1	B	121	THR	3.6
1	B	62	THR	3.6
1	B	117	GLU	3.6
1	B	85	ALA	3.6
1	B	60	GLY	3.5
1	B	330	LYS	3.5
1	B	318	LEU	3.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	59	ASP	3.4
1	B	108	VAL	3.4
1	B	67	LEU	3.4
1	B	112	ALA	3.4
1	A	180	ARG	3.3
1	A	177	VAL	3.3
1	B	337	GLY	3.2
1	A	76	GLY	3.1
1	B	64	THR	3.1
1	B	145	SER	3.1
1	B	82	GLU	3.0
1	B	63	TRP	3.0
1	A	115	LYS	3.0
1	B	414	PRO	2.9
1	B	105	ASN	2.9
1	B	327	PRO	2.9
1	B	69	ASP	2.9
1	B	109	ASP	2.9
1	B	3	PRO	2.9
1	A	175	TYR	2.8
1	B	389[A]	ARG	2.8
1	B	125	ALA	2.7
1	B	53	SER	2.6
1	B	338	GLN	2.6
1	A	21	LEU	2.6
1	B	129	PHE	2.6
1	B	329	GLY	2.5
1	B	374	ILE	2.5
1	A	382	TYR	2.5
1	A	143	ALA	2.5
1	A	174	GLN	2.5
1	A	144	PRO	2.5
1	B	441	THR	2.5
1	B	150	HIS	2.5
1	A	213	ASP	2.4
1	A	153	MET	2.4
1	B	79	PHE	2.4
1	B	66	THR	2.4
1	B	37	LEU	2.4
1	B	185	TRP	2.4
1	B	123	LYS	2.4
1	B	88	PHE	2.4

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	415	SER	2.3
1	B	325	THR	2.3
1	A	166	ILE	2.3
1	A	494	PHE	2.3
1	A	179	VAL	2.3
1	B	55	THR	2.3
1	B	451	TYR	2.3
1	A	170	SER	2.3
1	A	32	MET	2.3
1	B	52	LYS	2.2
1	A	4	ASP	2.2
1	A	58	GLU	2.2
1	B	430	LEU	2.2
1	A	114	SER	2.2
1	A	330	LYS	2.2
1	B	306	LYS	2.2
1	B	96	GLN	2.2
1	B	446	GLN	2.2
1	B	334	GLU	2.1
1	B	107	ILE	2.1
1	A	341	ARG	2.1
1	A	209	PHE	2.1
1	A	229	PHE	2.1
1	A	162	THR	2.1
1	B	284	TYR	2.1
1	B	57	SER	2.1
1	A	198	VAL	2.1
1	B	61	LYS	2.1
1	A	441	THR	2.1
1	B	86	GLU	2.1
1	B	92	LEU	2.1
1	A	192	LYS	2.1
1	A	29	ALA	2.0
1	A	75	ASN	2.0
1	B	456	THR	2.0
1	B	149	ASN	2.0
1	A	85	ALA	2.0
1	A	190	ALA	2.0
1	B	65	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	DTD	A	505	8/8	0.59	0.40	11.11	75,75,75,76	0
4	DTD	B	504	8/8	0.77	0.33	7.73	79,79,79,79	0
3	BHR	A	504	29/29	0.88	0.20	0.35	23,27,29,31	0
3	BHR	B	503	29/29	0.90	0.17	-0.98	19,27,31,32	0
2	FE	B	505	1/1	0.90	0.07	-	23,23,23,23	0
2	FE	A	503	1/1	0.95	0.08	-	20,20,20,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.