



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:50 AM GMT

PDB ID : 3N5X
Title : Structure of neuronal nitric oxide synthase heme domain in complex with 4-(2-(5-(2-(6-amino-4-methylpyridin-2-yl)ethyl)pyridin-3-yl)ethyl)-6-methylpyridin-2-amine
Authors : Li, H.; Delker, S.L.; Poulos, T.L.
Deposited on : 2010-05-25
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>10%</div> <div> <div></div> <div>91%</div> <div>5% . .</div> </div> </div>
1	B	422	<div> <div>5%</div> <div> <div></div> <div>94%</div> <div>. .</div> </div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7537 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	6	0
			3342	2142	567	611	22			
1	B	411	Total	C	N	O	S	0	6	0
			3370	2158	575	615	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 6-(2-{5-[2-(2-AMINO-6-METHYLPYRIDIN-4-YL)ETHYL]PYRIDIN-3-YL}ETHYL)-4-METHYLPYRIDIN-2-AMINE (three-letter code: XFK) (formula: $C_{21}H_{25}N_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			26	21	5		
3	A	1	Total	C	N	0	0
			26	21	5		
3	B	1	Total	C	N	0	0
			26	21	5		
3	B	1	Total	C	N	0	0
			26	21	5		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Cl 1 1	0	0
5	A	1	Total Cl 1 1	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	3	Total Zn 3 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	271	Total O 271 271	0	0
7	B	351	Total O 351 351	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase



- Molecule 1: Nitric oxide synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.29Å 111.10Å 164.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.62 – 1.80 38.62 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (38.62-1.80) 99.6 (38.62-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.173 , 0.198 0.189 , 0.216	Depositor DCC
R_{free} test set	4440 reflections (5.23%)	DCC
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.242	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 89269 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7537	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, ACT, XFK, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3454	0.58	0/4686
1	B	0.59	0/3483	0.60	0/4722
All	All	0.57	0/6937	0.59	0/9408

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3342	0	3257	14	0
1	B	3370	0	3290	8	0
2	A	43	0	30	3	0
2	B	43	0	30	4	0
3	A	52	0	50	2	0
3	B	52	0	50	3	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	271	0	0	3	0
7	B	351	0	0	2	0
All	All	7537	0	6713	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:750:HEM:O2D	3:B:800:XFK:H21	1.68	0.92
1:A:299:ARG:HB3	1:A:318:LEU:HD21	1.51	0.92
2:A:750:HEM:O2D	3:A:800:XFK:H21	1.72	0.89
1:A:299:ARG:HA	1:A:299:ARG:HE	1.54	0.73
2:B:750:HEM:HBB2	2:B:750:HEM:HHC	1.87	0.57
1:A:596:ARG:NH2	7:A:1042:HOH:O	2.40	0.53
1:A:569:ASN:HD22	1:A:569:ASN:H	1.56	0.52
1:A:380:ARG:HD3	1:A:400:GLU:OE1	2.08	0.52
1:A:686:SER:HA	1:A:691:PHE:CG	2.45	0.52
1:A:628:GLN:HG2	1:B:631:VAL:HG11	1.92	0.51
1:B:535:GLN:HG3	7:B:1514:HOH:O	2.10	0.51
1:A:609:GLU:HG3	7:A:1054:HOH:O	2.10	0.50
2:A:750:HEM:HHC	2:A:750:HEM:HBB2	1.92	0.50
1:B:686:SER:HA	1:B:691:PHE:CG	2.47	0.49
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.03	0.46
1:B:325:GLY:O	1:B:332:MET:HG3	2.16	0.45
1:B:364:GLN:NE2	7:B:1498:HOH:O	2.49	0.45
2:A:750:HEM:C1C	3:A:800:XFK:H07B	2.53	0.44
1:A:329:HIS:C	1:A:330:ILE:HG13	2.37	0.44
2:B:750:HEM:HBA1	2:B:750:HEM:HMA2	2.00	0.44
2:B:750:HEM:C1C	3:B:800:XFK:H07B	2.53	0.43
1:A:569:ASN:N	1:A:569:ASN:HD22	2.15	0.43
1:A:631:VAL:HG11	1:B:628:GLN:HG2	1.99	0.43
1:A:307:GLU:HG2	1:A:692:HIS:CG	2.54	0.43
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.56	0.40
1:B:485:TYR:CE1	1:B:514:ARG:HA	2.56	0.40
1:A:361:PHE:O	1:A:364:GLN:HG2	2.21	0.40
7:A:1150:HOH:O	3:B:805:XFK:H16	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/422 (97%)	399 (98%)	10 (2%)	0	100	100
1	B	413/422 (98%)	407 (98%)	6 (2%)	0	100	100
All	All	822/844 (97%)	806 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/377 (98%)	360 (98%)	9 (2%)	57	41
1	B	372/377 (99%)	369 (99%)	3 (1%)	86	83
All	All	741/754 (98%)	729 (98%)	12 (2%)	70	59

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	299	ARG
1	A	350	THR
1	A	352	ASP
1	A	380	ARG
1	A	481	ARG
1	A	527	ASN
1	A	547	ARG
1	A	569	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	645	LYS
1	B	350	THR
1	B	353	GLN
1	B	547	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	425	GLN
1	A	454	ASN
1	A	487	GLN
1	A	527	ASN
1	A	569	ASN
1	A	605	ASN
1	A	697	ASN
1	B	364	GLN
1	B	425	GLN
1	B	454	ASN
1	B	507	GLN
1	B	527	ASN
1	B	605	ASN
1	B	697	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 5 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	750	1	30,50,50	2.31	13 (43%)	24,82,82	2.54	9 (37%)
3	XFK	A	800	-	28,28,28	0.71	0	37,38,38	1.97	13 (35%)
3	XFK	A	805	6	28,28,28	0.64	0	37,38,38	1.81	11 (29%)
4	ACT	A	860	-	1,3,3	1.00	0	0,3,3	0.00	-
2	HEM	B	750	1	30,50,50	2.03	7 (23%)	24,82,82	2.39	9 (37%)
3	XFK	B	800	-	28,28,28	0.78	0	37,38,38	1.94	10 (27%)
3	XFK	B	805	6	28,28,28	0.61	0	37,38,38	1.66	9 (24%)
4	ACT	B	860	-	1,3,3	1.26	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/10/54/54	0/0/8/8
3	XFK	A	800	-	-	0/10/10/10	0/3/3/3
3	XFK	A	805	6	-	0/10/10/10	0/3/3/3
4	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/10/54/54	0/0/8/8
3	XFK	B	800	-	-	0/10/10/10	0/3/3/3
3	XFK	B	805	6	-	0/10/10/10	0/3/3/3
4	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3B-C4B	-6.70	1.45	1.51
2	B	750	HEM	C3B-C4B	-5.65	1.46	1.51
2	A	750	HEM	C3D-C4D	-5.40	1.44	1.51
2	B	750	HEM	C3D-C4D	-4.19	1.46	1.51
2	A	750	HEM	C2C-C1C	-3.96	1.45	1.52
2	B	750	HEM	C2C-C1C	-3.25	1.46	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C2B-C1B	-2.43	1.43	1.51
2	A	750	HEM	C2D-C1D	-2.34	1.44	1.51
2	A	750	HEM	FE-NB	2.00	2.08	1.97
2	A	750	HEM	C3B-CAB	2.03	1.55	1.51
2	A	750	HEM	FE-ND	2.04	2.08	1.97
2	A	750	HEM	C1C-NC	2.08	1.38	1.36
2	A	750	HEM	C3C-CAC	2.10	1.55	1.51
2	A	750	HEM	CMA-C3A	2.17	1.56	1.51
2	B	750	HEM	C3C-CAC	2.27	1.55	1.51
2	B	750	HEM	C3B-CAB	2.47	1.56	1.51
2	B	750	HEM	C1C-NC	2.63	1.39	1.36
2	A	750	HEM	CAA-C2A	2.65	1.56	1.52
2	A	750	HEM	FE-NC	3.27	2.08	1.95
2	B	750	HEM	FE-NC	3.48	2.09	1.95

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CBD-CAD-C3D	-4.22	101.28	113.55
3	A	800	XFK	C05-C06-N01	-3.92	118.53	122.96
2	B	750	HEM	CBD-CAD-C3D	-3.70	102.79	113.55
3	B	800	XFK	C05-C06-N01	-3.32	119.22	122.96
3	A	805	XFK	C04-C05-C06	-3.22	118.26	120.28
3	A	800	XFK	C26-C25-C24	-3.17	118.27	120.27
3	B	800	XFK	C09-C08-C06	-2.87	107.00	112.53
2	A	750	HEM	CMA-C3A-C4A	-2.62	124.03	128.36
3	A	800	XFK	C15-C14-C13	-2.52	117.98	121.25
3	A	805	XFK	C08-C06-C05	-2.49	117.64	121.13
3	A	805	XFK	C09-C13-C12	-2.42	117.74	121.85
3	B	800	XFK	C26-C25-C24	-2.32	118.81	120.27
2	B	750	HEM	C3B-C4B-NB	-2.30	107.24	111.63
3	B	805	XFK	C08-C06-C05	-2.26	117.95	121.13
3	B	805	XFK	C04-C05-C06	-2.15	118.93	120.28
2	B	750	HEM	CMA-C3A-C4A	-2.11	124.87	128.36
3	B	800	XFK	C15-C14-C13	-2.10	118.53	121.25
3	A	805	XFK	C17-C15-C16	-2.08	118.32	121.85
3	B	805	XFK	C15-C14-C13	-2.05	118.59	121.25
3	A	805	XFK	C15-C14-C13	-2.05	118.60	121.25
3	A	800	XFK	C17-C15-C16	-2.04	118.39	121.85
3	A	800	XFK	C09-C13-C12	-2.03	118.40	121.85
3	A	800	XFK	C14-C13-C12	2.05	118.81	116.57
3	A	800	XFK	C27-C24-N23	2.10	119.85	116.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	805	XFK	C14-C13-C12	2.14	118.91	116.57
3	A	800	XFK	C12-N11-C16	2.21	120.83	117.50
3	B	800	XFK	C12-N11-C16	2.36	121.06	117.50
3	B	805	XFK	N02-C02-N01	2.42	120.92	116.50
2	B	750	HEM	C3B-C4B-CHC	2.43	126.59	123.16
3	B	800	XFK	C08-C06-N01	2.49	119.36	115.69
3	A	805	XFK	C14-C15-C16	2.51	119.31	116.57
3	B	800	XFK	C22-N23-C24	2.58	120.93	118.12
3	A	805	XFK	N02-C02-N01	2.58	121.21	116.50
2	A	750	HEM	C2D-C3D-C4D	2.59	105.89	101.50
2	A	750	HEM	CMD-C2D-C3D	2.60	125.86	114.35
2	B	750	HEM	CMD-C2D-C3D	2.62	125.92	114.35
3	B	805	XFK	C27-C24-N23	2.72	120.80	116.59
3	B	805	XFK	C14-C15-C16	2.85	119.68	116.57
3	A	800	XFK	C08-C06-N01	2.87	119.93	115.69
3	A	805	XFK	C22-N23-C24	2.96	121.35	118.12
3	B	805	XFK	C02-N01-C06	2.98	120.35	118.23
3	B	800	XFK	N22-C22-N23	2.98	121.94	116.50
3	A	800	XFK	N22-C22-N23	3.05	122.07	116.50
3	B	805	XFK	C22-N23-C24	3.14	121.54	118.12
3	A	800	XFK	C14-C15-C16	3.28	120.15	116.57
3	A	800	XFK	C22-N23-C24	3.29	121.70	118.12
3	B	805	XFK	C08-C06-N01	3.40	120.70	115.69
3	B	800	XFK	C14-C15-C16	3.47	120.36	116.57
2	A	750	HEM	C3B-C4B-CHC	3.65	128.30	123.16
3	A	805	XFK	C08-C06-N01	3.70	121.16	115.69
2	B	750	HEM	CMC-C2C-C3C	4.31	127.29	116.53
2	A	750	HEM	CAD-C3D-C4D	4.43	128.09	112.47
2	A	750	HEM	CAD-C3D-C2D	4.44	125.99	113.22
2	B	750	HEM	CMB-C2B-C3B	4.51	127.79	116.53
2	B	750	HEM	CAD-C3D-C4D	4.53	128.46	112.47
2	A	750	HEM	CMB-C2B-C3B	4.60	128.01	116.53
3	A	805	XFK	C02-N01-C06	4.63	121.52	118.23
2	B	750	HEM	CAD-C3D-C2D	4.73	126.81	113.22
2	A	750	HEM	CMC-C2C-C3C	4.92	128.80	116.53
3	A	800	XFK	C02-N01-C06	5.53	122.16	118.23
3	B	800	XFK	C02-N01-C06	5.90	122.42	118.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	3	0
3	A	800	XFK	2	0
2	B	750	HEM	4	0
3	B	800	XFK	2	0
3	B	805	XFK	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	0.61	43 (10%) 8 6	17, 33, 60, 76	0
1	B	411/422 (97%)	0.22	19 (4%) 36 30	16, 26, 48, 66	0
All	All	818/844 (96%)	0.41	62 (7%) 17 13	16, 30, 57, 76	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	7.0
1	B	348	VAL	6.7
1	A	300	PHE	6.4
1	A	488	PRO	6.3
1	A	715	VAL	5.7
1	B	350	THR	5.6
1	A	355	PHE	5.5
1	A	350	THR	5.0
1	A	351	LYS	5.0
1	A	352	ASP	4.7
1	A	712	ASN	4.4
1	A	322	LEU	4.0
1	A	713	THR	3.9
1	B	619	ARG	3.8
1	A	299	ARG	3.8
1	A	716	TRP	3.7
1	A	392	SER	3.6
1	B	352	ASP	3.5
1	A	390	SER	3.3
1	B	718	GLY	3.2
1	A	567	VAL	3.1
1	A	321	THR	3.1
1	A	619	ARG	3.1
1	A	491	SER	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	490	GLY	2.9
1	A	489	ASP	2.9
1	A	680	VAL	2.9
1	A	389	GLU	2.9
1	B	321	THR	2.9
1	A	338	PRO	2.9
1	A	551	PHE	2.8
1	A	385	ASN	2.8
1	A	386	LYS	2.8
1	A	507	GLN	2.8
1	B	715	VAL	2.7
1	B	322	LEU	2.7
1	A	677	VAL	2.7
1	A	486	LYS	2.7
1	A	714	HIS	2.6
1	A	552	ASP	2.6
1	B	338	PRO	2.6
1	B	667	ARG	2.6
1	B	691	PHE	2.5
1	B	567	VAL	2.5
1	B	680	VAL	2.4
1	A	371	ARG	2.4
1	A	711	TRP	2.4
1	A	469	LYS	2.4
1	B	620	LYS	2.3
1	A	667	ARG	2.3
1	A	511	LYS	2.3
1	A	503	GLU	2.2
1	B	479	LEU	2.2
1	A	678	TRP	2.2
1	B	616	LEU	2.2
1	B	328	GLU	2.1
1	A	676	TRP	2.1
1	A	595	VAL	2.1
1	A	710	PRO	2.1
1	A	588	TYR	2.1
1	B	299	ARG	2.0
1	A	679	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ACT	A	860	4/4	0.95	0.12	1.04	40,40,40,42	0
3	XFK	A	800	26/26	0.93	0.20	0.91	19,26,33,34	0
3	XFK	B	800	26/26	0.95	0.17	0.81	17,22,27,29	0
2	HEM	A	750	43/43	0.98	0.18	0.80	17,20,28,33	0
2	HEM	B	750	43/43	0.98	0.15	0.78	14,17,24,28	0
4	ACT	B	860	4/4	0.95	0.09	0.39	28,32,32,33	0
3	XFK	A	805	26/26	0.88	0.18	0.26	23,30,44,45	0
3	XFK	B	805	26/26	0.93	0.14	-0.22	21,27,36,39	0
5	CL	A	910	1/1	0.98	0.15	-0.50	27,27,27,27	0
5	CL	B	910	1/1	0.98	0.13	-0.79	25,25,25,25	0
6	ZN	A	900	1/1	0.99	0.07	-1.08	23,23,23,23	0
6	ZN	A	719	1/1	0.99	0.13	-	20,20,20,20	1
6	ZN	A	901	1/1	0.99	0.11	-	22,22,22,22	1

6.5 Other polymers [i](#)

There are no such residues in this entry.