



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:03 PM GMT

PDB ID : 1NAK
Title : IGG1 FAB FRAGMENT (83.1) COMPLEX WITH 16-RESIDUE PEPTIDE
(RESIDUES 304-321 OF HIV-1 GP120 (MN ISOLATE))
Authors : Stanfield, R.L.; Ghiara, J.B.; Saphire, E.O.; Profy, A.T.; Wilson, I.A.
Deposited on : 2002-11-27
Resolution : 2.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2636 (2.60-2.56)
Clashscore	102246	3003 (2.60-2.56)
Ramachandran outliers	100387	2956 (2.60-2.56)
Sidechain outliers	100360	2956 (2.60-2.56)
RSRZ outliers	91569	2642 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	219	<div> <div>23%</div> <div>60%</div> <div>36%</div> <div>• •</div> </div>
1	M	219	<div> <div>32%</div> <div>58%</div> <div>36%</div> <div>5%</div> <div>•</div> </div>
2	H	214	<div> <div>17%</div> <div>64%</div> <div>33%</div> <div>•</div> </div>
2	I	214	<div> <div>24%</div> <div>63%</div> <div>35%</div> <div>•</div> </div>
3	P	16	<div> <div>38%</div> <div>31%</div> <div>19%</div> <div>13%</div> <div>38%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	16	 A horizontal bar chart showing the quality of chain Q. The bar is divided into four segments: red (31%), green (44%), yellow (19%), and grey (38%). The segments are stacked horizontally, with red on the left, followed by green, yellow, and grey on the right. The percentages are labeled above or below each segment.

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 83.1 - light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	217	Total	C	N	O	S	0	0	0
			1683	1053	286	338	6			
1	M	217	Total	C	N	O	S	0	0	0
			1683	1053	286	338	6			

- Molecule 2 is a protein called Fab 83.1 - heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	214	Total	C	N	O	S	40	0	0
			1625	1030	263	326	6			
2	I	214	Total	C	N	O	S	40	0	0
			1625	1030	263	326	6			

- Molecule 3 is a protein called Peptide MP1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	P	10	Total	C	N	O	0	0	0
			77	48	19	10			
3	Q	10	Total	C	N	O	0	0	0
			77	48	19	10			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	22	Total	O	0	0
			22	22		
4	I	19	Total	O	0	0
			19	19		
4	L	32	Total	O	0	0
			32	32		
4	M	34	Total	O	0	0
			34	34		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab 83.1 - light chain




- Molecule 1: Fab 83.1 - light chain




- Molecule 2: Fab 83.1 - heavy chain







• Molecule 2: Fab 83.1 - heavy chain



• Molecule 3: Peptide MP1



• Molecule 3: Peptide MP1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.51Å 122.60Å 69.46Å 90.00° 108.54° 90.00°	Depositor
Resolution (Å)	48.08 – 2.57 48.08 – 2.57	Depositor EDS
% Data completeness (in resolution range)	97.1 (48.08-2.57) 96.9 (48.08-2.57)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.58Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.288 , 0.326 0.290 , 0.329	Depositor DCC
R_{free} test set	1469 reflections (4.94%)	DCC
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	1 of 29751 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	6877	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.65 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.2623e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.36	0/1724	0.64	0/2338
1	M	0.36	0/1724	0.64	0/2338
2	H	0.38	0/1668	0.78	3/2282 (0.1%)
2	I	0.37	0/1668	0.78	4/2282 (0.2%)
3	P	0.82	0/78	1.66	2/102 (2.0%)
3	Q	0.64	0/78	1.39	1/102 (1.0%)
All	All	0.38	0/6940	0.74	10/9444 (0.1%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	313	ARG	N-CA-C	10.92	140.49	111.00
2	I	133	GLN	N-CA-C	9.81	137.48	111.00
2	H	133	GLN	N-CA-C	9.56	136.82	111.00
2	I	130	ALA	N-CA-C	-8.59	87.82	111.00
2	H	130	ALA	N-CA-C	-8.43	88.25	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1683	0	1612	68	0
1	M	1683	0	1612	74	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	1625	0	1594	74	0
2	I	1625	0	1594	66	0
3	P	77	0	85	7	0
3	Q	77	0	85	4	0
4	H	22	0	0	0	0
4	I	19	0	0	0	0
4	L	32	0	0	1	0
4	M	34	0	0	2	0
All	All	6877	0	6582	271	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 271 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:200:ARG:HG3	2:I:200:ARG:HH11	1.21	1.02
1:M:21:ILE:HD12	1:M:102:THR:HG21	1.46	0.95
1:L:50:LYS:HB2	1:L:53:ASN:HD22	1.34	0.93
2:H:200:ARG:HH11	2:H:200:ARG:CG	1.83	0.90
1:M:50:LYS:HB2	1:M:53:ASN:HD22	1.32	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	215/219 (98%)	200 (93%)	13 (6%)	2 (1%)	21	41
1	M	215/219 (98%)	200 (93%)	9 (4%)	6 (3%)	6	9
2	H	212/214 (99%)	192 (91%)	17 (8%)	3 (1%)	14	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	212/214 (99%)	192 (91%)	15 (7%)	5 (2%)	7	13
3	P	8/16 (50%)	4 (50%)	3 (38%)	1 (12%)	0	0
3	Q	8/16 (50%)	5 (62%)	3 (38%)	0	100	100
All	All	870/898 (97%)	793 (91%)	60 (7%)	17 (2%)	9	17

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	134	THR
3	P	313	ARG
1	M	68	GLY
2	I	134	THR
2	I	135	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	194/196 (99%)	184 (95%)	10 (5%)	29	53
1	M	194/196 (99%)	185 (95%)	9 (5%)	33	59
2	H	190/190 (100%)	184 (97%)	6 (3%)	46	73
2	I	190/190 (100%)	185 (97%)	5 (3%)	54	78
3	P	7/13 (54%)	6 (86%)	1 (14%)	4	7
3	Q	7/13 (54%)	7 (100%)	0	100	100
All	All	782/798 (98%)	751 (96%)	31 (4%)	38	65

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	187	LEU
1	M	1	ASP
2	I	183	ASP
2	H	200	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	M	4	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	16	GLN
2	H	162	ASN
1	M	53	ASN
1	L	190	ASN
1	M	190	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	217/219 (99%)	1.43	51 (23%) 1 0	11, 28, 44, 51	2 (0%)
1	M	217/219 (99%)	1.61	69 (31%) 1 0	9, 29, 45, 61	2 (0%)
2	H	208/214 (97%)	1.39	37 (17%) 2 1	12, 26, 43, 59	4 (1%)
2	I	208/214 (97%)	1.55	52 (25%) 1 0	13, 29, 47, 67	4 (1%)
3	P	10/16 (62%)	2.56	6 (60%) 0 0	33, 41, 45, 52	0
3	Q	10/16 (62%)	2.34	5 (50%) 0 0	26, 36, 44, 50	0
All	All	870/898 (96%)	1.52	220 (25%) 1 0	9, 28, 45, 67	12 (1%)

The worst 5 of 220 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	1	GLU	7.1
2	I	4	LEU	6.7
2	I	199	PRO	6.6
2	H	108	THR	6.4
2	H	229	ASP	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.