



wwPDB NMR Structure Validation Summary Report ⓘ

Jun 20, 2016 – 11:02 PM EDT

PDB ID : 2NBQ
Title : NMR Structure of the C-Terminal Domain of human APOBEC3B
Authors : Byeon, I.L.; Byeon, C.; Gronenborn, A.M.
Deposited on : 2016-03-09

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027790
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027790


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 64%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	205	<div> <div style="width: 28%; background-color: green;"></div> <div style="width: 44%; background-color: yellow;"></div> <div style="width: 8%; background-color: orange;"></div> <div style="width: 16%; background-color: cyan;"></div> <div style="width: 4%; background-color: grey;"></div> </div> <div> <div style="width: 28%;"></div> <div style="width: 44%;"></div> <div style="width: 8%;"></div> <div style="width: 16%;"></div> <div style="width: 4%;"></div> </div>

2 Ensemble composition and analysis

This entry contains 30 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:191-A:202, A:214-A:240, A:252-A:377 (165)	0.45	11

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 1 single-model cluster was found.

Cluster number	Models
1	6, 7, 8, 15, 16, 20, 23, 24, 25, 27
2	1, 4, 5, 11, 12, 17, 19, 22, 28, 30
3	14, 18, 21, 26
4	3, 9, 10
5	13, 29
Single-model clusters	2

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3208 atoms, of which 1563 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DNA dC->dU-editing enzyme APOBEC-3B.

Mol	Chain	Residues	Atoms						Trace
1	A	197	Total	C	H	N	O	S	0
			3207	1055	1563	282	294	13	

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	186	MET	-	EXPRESSION TAG	UNP Q9UH17
A	383	LEU	-	EXPRESSION TAG	UNP Q9UH17
A	384	GLU	-	EXPRESSION TAG	UNP Q9UH17
A	385	HIS	-	EXPRESSION TAG	UNP Q9UH17
A	386	HIS	-	EXPRESSION TAG	UNP Q9UH17
A	387	HIS	-	EXPRESSION TAG	UNP Q9UH17
A	388	HIS	-	EXPRESSION TAG	UNP Q9UH17
A	389	HIS	-	EXPRESSION TAG	UNP Q9UH17
A	390	HIS	-	EXPRESSION TAG	UNP Q9UH17

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3B



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 11. Colouring as in section 4.1 above.

- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3B



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 256 calculated structures, 30 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
XPLOR-NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2nbq_cs.cif
Number of chemical shift lists	2
Total number of shifts	2228
Number of shifts mapped to atoms	2228
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	64%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1382	1298	1298	87±10
All	All	41490	38940	38940	2604

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

5 of 763 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:335:ILE:HD11	1:A:369:LEU:HD23	0.85	1.46	16	4
1:A:327:ARG:CZ	1:A:377:LEU:HD11	0.81	2.06	13	2
1:A:312:ILE:HD12	1:A:312:ILE:H	0.80	1.36	3	1
1:A:307:ILE:HD12	1:A:307:ILE:N	0.79	1.92	11	4
1:A:238:LEU:N	1:A:238:LEU:HD23	0.79	1.91	6	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/205 (80%)	131±4 (80±2%)	25±3 (15±2%)	9±2 (6±1%)	4	23
All	All	4950/6150 (80%)	3936 (80%)	738 (15%)	276 (6%)	4	23

5 of 34 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	287	TRP	28
1	A	359	TRP	27
1	A	360	ASP	26
1	A	192	LEU	22
1	A	281	TRP	20

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/182 (80%)	104±4 (71±3%)	42±4 (29±3%)	2	19
All	All	4380/5460 (80%)	3129 (71%)	1251 (29%)	2	19

5 of 118 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	302	HIS	30
1	A	303	VAL	30
1	A	275	VAL	30
1	A	334	SER	30
1	A	337	THR	30

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 64% for the well-defined parts and 61% for the entire structure.

7.1 Chemical shift list 1

File name: 2nbq_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1840
Number of shifts mapped to atoms	1840
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	11

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	187	-0.41 ± 0.07	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	171	0.56 ± 0.18	Should be applied
$^{13}\text{C}'$	0	—	—
^{15}N	180	-0.12 ± 0.21	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 64%, i.e. 1362 atoms were assigned a chemical shift out of a possible 2129. 0 out of 27 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	613/811 (76%)	304/323 (94%)	156/330 (47%)	153/158 (97%)
Sidechain	632/1030 (61%)	387/607 (64%)	236/373 (63%)	9/50 (18%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	117/288 (41%)	111/152 (73%)	0/126 (0%)	6/10 (60%)
Overall	1362/2129 (64%)	802/1082 (74%)	392/829 (47%)	168/218 (77%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	351	ARG	HD3	0.32	4.36 – 1.86	-11.2
1	A	277	TRP	HE1	5.50	12.85 – 7.35	-8.4
1	A	255	GLU	H	12.41	11.34 – 5.34	6.8
1	A	223	LEU	HD23	-0.84	2.14 – -0.66	-5.6
1	A	223	LEU	HD21	-0.84	2.14 – -0.66	-5.6
1	A	223	LEU	HD22	-0.84	2.14 – -0.66	-5.6
1	A	351	ARG	HD2	1.85	4.27 – 1.97	-5.5
1	A	277	TRP	NE1	118.98	139.19 – 119.59	-5.3
1	A	347	THR	HG21	-0.03	2.29 – -0.01	-5.1
1	A	347	THR	HG22	-0.03	2.29 – -0.01	-5.1
1	A	347	THR	HG23	-0.03	2.29 – -0.01	-5.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: 2nbq_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1_dup*

7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	388
Number of shifts mapped to atoms	388
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	—
$^{13}\text{C}_\beta$	0	—	—
$^{13}\text{C}'$	0	—	—
^{15}N	170	0.25 ± 0.55	None needed (< 0.5 ppm)

7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 15%, i.e. 329 atoms were assigned a chemical shift out of a possible 2129. 0 out of 27 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	291/811 (36%)	145/323 (45%)	0/330 (0%)	146/158 (92%)
Sidechain	26/1030 (3%)	17/607 (3%)	0/373 (0%)	9/50 (18%)
Aromatic	12/288 (4%)	6/152 (4%)	0/126 (0%)	6/10 (60%)
Overall	329/2129 (15%)	168/1082 (16%)	0/829 (0%)	161/218 (74%)

7.2.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, <i>ppm</i>	Expected range, <i>ppm</i>	Z-score
1	A	277	TRP	HE1	5.52	12.85 – 7.35	-8.3
1	A	255	GLU	H	12.56	11.34 – 5.34	7.0
1	A	277	TRP	NE1	119.18	139.19 – 119.59	-5.2

7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

