



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:00 PM GMT

PDB ID : 4NGN
Title : Crystal Structure of Glutamate Carboxypeptidase II in a complex with urea-based inhibitor
Authors : Tykvart, J.; Pachl, P.
Deposited on : 2013-11-02
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1953 (1.66-1.62)
Clashscore	102246	2091 (1.66-1.62)
Ramachandran outliers	100387	2052 (1.66-1.62)
Sidechain outliers	100360	2052 (1.66-1.62)
RSRZ outliers	91569	1955 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	739	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MAN	A	814	-	-	-	X

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 6305 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	696	Total	C	N	O	S	0	9	0
			5547	3570	925	1034	18			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	ARG	-	EXPRESSION TAG	UNP Q04609
A	13	SER	-	EXPRESSION TAG	UNP Q04609
A	14	GLY	-	EXPRESSION TAG	UNP Q04609
A	15	LEU	-	EXPRESSION TAG	UNP Q04609
A	16	ASN	-	EXPRESSION TAG	UNP Q04609
A	17	ASP	-	EXPRESSION TAG	UNP Q04609
A	18	ILE	-	EXPRESSION TAG	UNP Q04609
A	19	PHE	-	EXPRESSION TAG	UNP Q04609
A	20	GLU	-	EXPRESSION TAG	UNP Q04609
A	21	ALA	-	EXPRESSION TAG	UNP Q04609
A	22	GLN	-	EXPRESSION TAG	UNP Q04609
A	23	LYS	-	EXPRESSION TAG	UNP Q04609
A	24	ILE	-	EXPRESSION TAG	UNP Q04609
A	25	GLU	-	EXPRESSION TAG	UNP Q04609
A	26	TRP	-	EXPRESSION TAG	UNP Q04609
A	27	HIS	-	EXPRESSION TAG	UNP Q04609
A	28	GLU	-	EXPRESSION TAG	UNP Q04609
A	29	GLY	-	EXPRESSION TAG	UNP Q04609
A	30	SER	-	EXPRESSION TAG	UNP Q04609
A	31	GLY	-	EXPRESSION TAG	UNP Q04609
A	32	SER	-	EXPRESSION TAG	UNP Q04609
A	33	GLY	-	EXPRESSION TAG	UNP Q04609
A	34	SER	-	EXPRESSION TAG	UNP Q04609
A	35	GLU	-	EXPRESSION TAG	UNP Q04609
A	36	ASN	-	EXPRESSION TAG	UNP Q04609
A	37	LEU	-	EXPRESSION TAG	UNP Q04609
A	38	TYR	-	EXPRESSION TAG	UNP Q04609

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	PHE	-	EXPRESSION TAG	UNP Q04609
A	40	GLN	-	EXPRESSION TAG	UNP Q04609
A	41	GLY	-	EXPRESSION TAG	UNP Q04609
A	42	ARG	-	EXPRESSION TAG	UNP Q04609
A	43	SER	-	EXPRESSION TAG	UNP Q04609

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 5 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Zn	0	0
			2	2		


- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Ca	0	0
			1	1		

- Molecule 9 is N-([(1S)-1-CARBOXY-5-({[2-({4,44-DIOXO-48-[(3AS,4S,6AR)-2-OXOH EXAHYDRO-1H-THIENO[3,4-D]IMIDAZOL-4-YL]-7,10,13,16,19,22,25,28,31,34,37,40-DODECAOXA-3,43-DIAZAOCTATETRACONT-1-YL}OXY)PHENYL]ACETYL}AMINO)PENTYL]CARBAMOYL}-L-GLUTAMIC ACID (three-letter code: J96) (formula: C₅₉H₉₉N₇O₂₄S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
9	A	1	24	13	3	8	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	541	Total	O	0	3
			543	543		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutamate carboxypeptidase 2



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.34Å 130.02Å 158.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.25 – 1.64 45.25 – 1.64	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.25-1.64) 99.8 (45.25-1.64)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 1.64Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.185 , 0.213 0.179 , 0.210	Depositor DCC
R_{free} test set	6387 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	20.5	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 127795 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6305	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: J96, ZN, BMA, NAG, CL, CA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	4/5730 (0.1%)	0.99	11/7772 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	437	GLU	CD-OE2	-6.06	1.19	1.25
1	A	434	TRP	CE3-CZ3	5.88	1.48	1.38
1	A	525	PHE	CG-CD1	5.60	1.47	1.38
1	A	557	GLU	CD-OE2	-5.25	1.19	1.25

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	673	ARG	NE-CZ-NH2	-8.49	116.05	120.30
1	A	370	ARG	NE-CZ-NH1	7.91	124.26	120.30
1	A	673	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	A	688	ARG	NE-CZ-NH1	-6.29	117.16	120.30
1	A	309	MET	CG-SD-CE	-6.08	90.48	100.20
1	A	387	ASP	CB-CG-OD1	-5.86	113.03	118.30
1	A	63	ASP	CB-CG-OD1	5.44	123.20	118.30
1	A	465	ASP	CB-CG-OD1	5.44	123.20	118.30
1	A	188	LEU	CA-CB-CG	5.42	127.78	115.30
1	A	370	ARG	NE-CZ-NH2	-5.04	117.78	120.30
1	A	684	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5547	0	5374	42	0
2	A	56	0	50	0	0
3	A	42	0	39	1	0
4	A	39	0	34	0	0
5	A	50	0	43	0	0
6	A	2	0	0	0	0
7	A	1	0	0	0	0
8	A	1	0	0	0	0
9	A	24	0	17	0	0
10	A	543	0	0	3	1
All	All	6305	0	5557	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:659:ILE:O	1:A:663[B]:MET:HG3	1.78	0.83
1:A:58:MET:CE	1:A:586:GLU:HG2	2.09	0.81
1:A:506:PHE:CB	1:A:509:MET:HG3	2.12	0.79
1:A:58:MET:HE1	1:A:586:GLU:HG2	1.65	0.79
1:A:122:LYS:HD2	1:A:122:LYS:N	2.08	0.66
1:A:122:LYS:CD	1:A:122:LYS:H	2.10	0.65
1:A:188:LEU:HD11	1:A:329:VAL:HG11	1.80	0.63
1:A:122:LYS:HD2	1:A:122:LYS:H	1.66	0.61
1:A:132:ASN:HD21	1:A:136:ASN:HD22	1.52	0.58
1:A:506:PHE:O	1:A:507:SER:HB3	2.03	0.58
1:A:215:LYS:HE2	10:A:1133:HOH:O	2.03	0.58
1:A:155:SER:O	1:A:156:ASP:HB2	2.06	0.56
1:A:177:VAL:HG12	1:A:188:LEU:HD21	1.91	0.53
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.43	0.53
1:A:659:ILE:HG22	1:A:663[B]:MET:CE	2.39	0.52
1:A:188:LEU:HD11	1:A:329:VAL:CG1	2.40	0.52
1:A:246:TRP:CD1	3:A:807:NAG:H83	2.46	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:551:LEU:HD22	1:A:556:TYR:HB2	1.94	0.48
1:A:188:LEU:CD1	1:A:329:VAL:HG11	2.45	0.47
1:A:663[A]:MET:HB2	1:A:663[A]:MET:HE2	1.68	0.47
1:A:58:MET:HE2	1:A:586:GLU:HG2	1.96	0.46
1:A:258:ILE:HD13	1:A:294:VAL:HB	1.98	0.46
1:A:517:SER:HB2	1:A:694:PRO:HG3	1.99	0.45
1:A:103:PHE:O	1:A:403:GLY:HA3	2.18	0.44
1:A:586:GLU:OE1	1:A:586:GLU:HA	2.18	0.44
1:A:507:SER:HA	10:A:1145:HOH:O	2.18	0.44
1:A:656:SER:O	1:A:658:PRO:HD3	2.18	0.43
1:A:169:MET:HA	1:A:344:MET:O	2.17	0.43
1:A:547:SER:O	1:A:548:GLY:C	2.54	0.43
1:A:148:PRO:HG2	1:A:151:TYR:CD1	2.53	0.43
1:A:497:TRP:CE2	1:A:510:PRO:HB2	2.54	0.42
1:A:310:GLY:O	1:A:328:ASN:HB3	2.20	0.42
1:A:503:SER:HB3	1:A:509:MET:O	2.19	0.41
1:A:657:ASN:HA	1:A:658:PRO:HD2	1.81	0.41
1:A:449:TYR:O	1:A:532:SER:HA	2.20	0.41
1:A:215:LYS:HE3	10:A:1209:HOH:O	2.20	0.41
1:A:659:ILE:HG22	1:A:663[B]:MET:HE2	2.02	0.41
1:A:718:LYS:HD3	1:A:718:LYS:HA	1.76	0.41
1:A:387:ASP:HA	1:A:388:PRO:HA	1.80	0.41
1:A:593:LEU:HA	1:A:594:PRO:HD3	1.99	0.41
1:A:551:LEU:HA	1:A:551:LEU:HD23	1.81	0.40
1:A:726:GLY:HA2	1:A:729[B]:LYS:HE3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:A:1352:HOH:O	10:A:1352:HOH:O[2_555]	2.15	0.05

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	703/739 (95%)	679 (97%)	22 (3%)	2 (0%)	46	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	338	SER
1	A	382	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	590/629 (94%)	579 (98%)	11 (2%)	65	38

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	LYS
1	A	188	LEU
1	A	189	GLU
1	A	303	GLN
1	A	339	THR
1	A	388	PRO
1	A	503	SER
1	A	519	ASN
1	A	537	TYR
1	A	600	TYR
1	A	673	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	136	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	620	GLN
1	A	651	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

11 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	801	1,2	14,14,15	0.49	0	15,19,21	0.79	0
2	NAG	A	802	2	14,14,15	0.57	0	15,19,21	1.02	1 (6%)
2	NAG	A	804	1,2	14,14,15	0.75	0	15,19,21	0.97	0
2	NAG	A	805	2	14,14,15	0.35	0	15,19,21	1.02	0
4	NAG	A	808	1,4	14,14,15	0.74	0	15,19,21	1.26	1 (6%)
4	NAG	A	809	4	14,14,15	0.64	0	15,19,21	1.85	3 (20%)
4	BMA	A	810	4	11,11,12	0.28	0	14,15,17	1.01	0
5	NAG	A	811	1,5	14,14,15	0.70	0	15,19,21	1.14	1 (6%)
5	NAG	A	812	5	14,14,15	0.89	1 (7%)	15,19,21	1.36	3 (20%)
5	BMA	A	813	5	11,11,12	0.69	0	14,15,17	1.30	1 (7%)
5	MAN	A	814	5	11,11,12	0.95	1 (9%)	14,15,17	1.90	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	801	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	802	2	-	0/6/23/26	0/1/1/1
2	NAG	A	804	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	805	2	-	0/6/23/26	0/1/1/1
4	NAG	A	808	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	809	4	-	0/6/23/26	0/1/1/1
4	BMA	A	810	4	-	0/2/19/22	0/1/1/1
5	NAG	A	811	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	812	5	-	0/6/23/26	0/1/1/1
5	BMA	A	813	5	-	0/2/19/22	0/1/1/1
5	MAN	A	814	5	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	812	NAG	O5-C1	-3.09	1.38	1.43
5	A	814	MAN	C2-C3	2.45	1.55	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	809	NAG	C2-N2-C7	-5.26	116.29	123.04
5	A	813	BMA	O3-C3-C2	-4.00	102.77	110.00
5	A	814	MAN	O2-C2-C1	-3.32	102.54	109.21
5	A	814	MAN	C1-C2-C3	-2.93	106.07	109.54
4	A	809	NAG	C3-C4-C5	-2.78	105.35	110.20
2	A	802	NAG	C3-C4-C5	-2.60	105.67	110.20
5	A	811	NAG	C6-C5-C4	-2.51	106.82	113.02
5	A	812	NAG	O7-C7-C8	-2.25	117.93	122.06
4	A	809	NAG	C1-O5-C5	-2.18	109.48	112.25
5	A	812	NAG	C8-C7-N2	2.56	121.01	116.11
5	A	814	MAN	O4-C4-C3	2.57	116.12	110.34
4	A	808	NAG	C8-C7-N2	2.63	121.13	116.11
5	A	814	MAN	O2-C2-C3	2.92	115.98	110.12
5	A	812	NAG	C1-O5-C5	2.93	115.97	112.25
5	A	814	MAN	C1-O5-C5	3.07	116.14	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	803	1	14,14,15	0.65	0	15,19,21	1.74	5 (33%)
3	NAG	A	806	1	14,14,15	0.52	0	15,19,21	1.73	3 (20%)
3	NAG	A	807	1	14,14,15	0.71	0	15,19,21	1.93	4 (26%)
9	J96	A	819	6	14,23,93	0.77	0	12,28,110	1.53	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	803	1	-	0/6/23/26	0/1/1/1
3	NAG	A	806	1	-	0/6/23/26	0/1/1/1
3	NAG	A	807	1	-	0/6/23/26	0/1/1/1
9	J96	A	819	6	-	0/18/28/110	0/0/0/3

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	807	NAG	O3-C3-C4	-2.98	103.62	110.34
3	A	803	NAG	O3-C3-C2	-2.85	103.46	109.11
9	A	819	J96	C5-C4-C3	-2.68	107.55	112.99
3	A	807	NAG	O7-C7-C8	-2.49	117.50	122.06
9	A	819	J96	C3-N1-C2	-2.41	115.97	120.90
3	A	803	NAG	O3-C3-C4	-2.36	105.03	110.34
3	A	806	NAG	O3-C3-C2	-2.27	104.62	109.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	807	NAG	C2-N2-C7	-2.07	120.38	123.04
3	A	803	NAG	O6-C6-C5	2.04	118.08	111.33
3	A	803	NAG	C4-C3-C2	2.59	115.25	111.23
3	A	806	NAG	O5-C5-C6	2.60	112.98	107.35
3	A	803	NAG	O5-C5-C6	3.78	115.54	107.35
3	A	806	NAG	C1-O5-C5	4.16	117.52	112.25
3	A	807	NAG	C1-O5-C5	4.42	117.86	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	807	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	696/739 (94%)	-0.07	20 (2%) 55 53	12, 21, 41, 67	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	541	TRP	6.3
1	A	543	THR	5.3
1	A	506	PHE	3.9
1	A	335	GLY	3.5
1	A	507	SER	3.5
1	A	336	ASN	3.1
1	A	153	ASN	3.1
1	A	155	SER	3.1
1	A	542	GLU	3.0
1	A	154	VAL	2.8
1	A	134	ASP	2.7
1	A	334	THR	2.7
1	A	339	THR	2.6
1	A	136	ASN	2.6
1	A	487	GLY	2.6
1	A	152	GLU	2.5
1	A	135	GLY	2.5
1	A	122	LYS	2.2
1	A	719	VAL	2.2
1	A	337	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MAN	A	814	11/12	0.66	0.23	4.79	47,54,59,62	0
5	NAG	A	811	14/15	0.94	0.07	1.87	19,25,34,46	0
4	NAG	A	808	14/15	0.93	0.08	1.33	25,27,30,37	0
2	NAG	A	802	14/15	0.75	0.28	-	33,42,50,60	0
2	NAG	A	801	14/15	0.94	0.12	-	26,35,41,45	0
5	BMA	A	813	11/12	0.89	0.10	-	34,36,40,41	0
2	NAG	A	805	14/15	0.87	0.24	-	46,50,61,68	0
5	NAG	A	812	14/15	0.91	0.14	-	32,35,45,55	0
2	NAG	A	804	14/15	0.88	0.12	-	31,36,41,47	0
4	NAG	A	809	14/15	0.93	0.11	-	31,39,50,50	0
4	BMA	A	810	11/12	0.69	0.25	-	53,58,67,69	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	803	14/15	0.77	0.23	1.31	39,47,57,64	0
9	J96	A	819	24/91	0.97	0.14	1.12	13,16,51,55	0
3	NAG	A	807	14/15	0.88	0.10	0.11	26,38,43,44	0
8	CA	A	818	1/1	1.00	0.08	-0.01	13,13,13,13	0
6	ZN	A	816	1/1	1.00	0.08	-1.44	15,15,15,15	0
6	ZN	A	815	1/1	1.00	0.08	-2.03	14,14,14,14	0
7	CL	A	817	1/1	0.99	0.07	-2.58	15,15,15,15	0
3	NAG	A	806	14/15	0.85	0.20	-	49,56,61,64	0

6.5 Other polymers ⓘ

There are no such residues in this entry.