



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:55 AM GMT

PDB ID : 3NHQ
Title : The dark Pfr structure of the photosensory core module of *P. aeruginosa* Bacteriophytochrome
Authors : Yang, X.; Ren, Z.; Kuk, J.; Moffat, K.
Deposited on : 2010-06-14
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4549 (2.58-2.50)
Clashscore	102246	5292 (2.58-2.50)
Ramachandran outliers	100387	5194 (2.58-2.50)
Sidechain outliers	100360	5196 (2.58-2.50)
RSRZ outliers	91569	4561 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	
1	B	505	
1	C	505	
1	D	505	
1	E	505	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	505	 <p>11% 64% 25% 5% 7%</p>
1	G	505	 <p>2% 65% 26% 5%</p>
1	H	505	 <p>15% 64% 28% 5%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 30788 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriophytochrome.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	479	3790	2386	682	703	19	0	0	0
1	B	470	3709	2332	665	693	19	0	0	0
1	C	482	3812	2398	688	707	19	0	0	0
1	D	480	3801	2391	686	705	19	0	0	0
1	E	479	3790	2386	683	702	19	0	0	0
1	F	471	3717	2338	666	694	19	0	0	0
1	G	482	3816	2401	688	708	19	0	1	0
1	H	481	3808	2396	687	706	19	0	0	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	498	LEU	-	EXPRESSION TAG	UNP Q9HWR3
A	499	GLU	-	EXPRESSION TAG	UNP Q9HWR3
A	500	HIS	-	EXPRESSION TAG	UNP Q9HWR3
A	501	HIS	-	EXPRESSION TAG	UNP Q9HWR3
A	502	HIS	-	EXPRESSION TAG	UNP Q9HWR3
A	503	HIS	-	EXPRESSION TAG	UNP Q9HWR3
A	504	HIS	-	EXPRESSION TAG	UNP Q9HWR3
A	505	HIS	-	EXPRESSION TAG	UNP Q9HWR3
B	498	LEU	-	EXPRESSION TAG	UNP Q9HWR3
B	499	GLU	-	EXPRESSION TAG	UNP Q9HWR3
B	500	HIS	-	EXPRESSION TAG	UNP Q9HWR3
B	501	HIS	-	EXPRESSION TAG	UNP Q9HWR3
B	502	HIS	-	EXPRESSION TAG	UNP Q9HWR3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	503	HIS	-	EXPRESSION TAG	UNP Q9HWR3
B	504	HIS	-	EXPRESSION TAG	UNP Q9HWR3
B	505	HIS	-	EXPRESSION TAG	UNP Q9HWR3
C	498	LEU	-	EXPRESSION TAG	UNP Q9HWR3
C	499	GLU	-	EXPRESSION TAG	UNP Q9HWR3
C	500	HIS	-	EXPRESSION TAG	UNP Q9HWR3
C	501	HIS	-	EXPRESSION TAG	UNP Q9HWR3
C	502	HIS	-	EXPRESSION TAG	UNP Q9HWR3
C	503	HIS	-	EXPRESSION TAG	UNP Q9HWR3
C	504	HIS	-	EXPRESSION TAG	UNP Q9HWR3
C	505	HIS	-	EXPRESSION TAG	UNP Q9HWR3
D	498	LEU	-	EXPRESSION TAG	UNP Q9HWR3
D	499	GLU	-	EXPRESSION TAG	UNP Q9HWR3
D	500	HIS	-	EXPRESSION TAG	UNP Q9HWR3
D	501	HIS	-	EXPRESSION TAG	UNP Q9HWR3
D	502	HIS	-	EXPRESSION TAG	UNP Q9HWR3
D	503	HIS	-	EXPRESSION TAG	UNP Q9HWR3
D	504	HIS	-	EXPRESSION TAG	UNP Q9HWR3
D	505	HIS	-	EXPRESSION TAG	UNP Q9HWR3
E	498	LEU	-	EXPRESSION TAG	UNP Q9HWR3
E	499	GLU	-	EXPRESSION TAG	UNP Q9HWR3
E	500	HIS	-	EXPRESSION TAG	UNP Q9HWR3
E	501	HIS	-	EXPRESSION TAG	UNP Q9HWR3
E	502	HIS	-	EXPRESSION TAG	UNP Q9HWR3
E	503	HIS	-	EXPRESSION TAG	UNP Q9HWR3
E	504	HIS	-	EXPRESSION TAG	UNP Q9HWR3
E	505	HIS	-	EXPRESSION TAG	UNP Q9HWR3
F	498	LEU	-	EXPRESSION TAG	UNP Q9HWR3
F	499	GLU	-	EXPRESSION TAG	UNP Q9HWR3
F	500	HIS	-	EXPRESSION TAG	UNP Q9HWR3
F	501	HIS	-	EXPRESSION TAG	UNP Q9HWR3
F	502	HIS	-	EXPRESSION TAG	UNP Q9HWR3
F	503	HIS	-	EXPRESSION TAG	UNP Q9HWR3
F	504	HIS	-	EXPRESSION TAG	UNP Q9HWR3
F	505	HIS	-	EXPRESSION TAG	UNP Q9HWR3
G	498	LEU	-	EXPRESSION TAG	UNP Q9HWR3
G	499	GLU	-	EXPRESSION TAG	UNP Q9HWR3
G	500	HIS	-	EXPRESSION TAG	UNP Q9HWR3
G	501	HIS	-	EXPRESSION TAG	UNP Q9HWR3
G	502	HIS	-	EXPRESSION TAG	UNP Q9HWR3
G	503	HIS	-	EXPRESSION TAG	UNP Q9HWR3
G	504	HIS	-	EXPRESSION TAG	UNP Q9HWR3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	505	HIS	-	EXPRESSION TAG	UNP Q9HWR3
H	498	LEU	-	EXPRESSION TAG	UNP Q9HWR3
H	499	GLU	-	EXPRESSION TAG	UNP Q9HWR3
H	500	HIS	-	EXPRESSION TAG	UNP Q9HWR3
H	501	HIS	-	EXPRESSION TAG	UNP Q9HWR3
H	502	HIS	-	EXPRESSION TAG	UNP Q9HWR3
H	503	HIS	-	EXPRESSION TAG	UNP Q9HWR3
H	504	HIS	-	EXPRESSION TAG	UNP Q9HWR3
H	505	HIS	-	EXPRESSION TAG	UNP Q9HWR3

- Molecule 2 is BILIVERDINE IX ALPHA (three-letter code: BLA) (formula: C₃₃H₃₄N₄O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	N	O	0	0
			43	33	4	6		
2	B	1	Total	C	N	O	0	0
			43	33	4	6		
2	C	1	Total	C	N	O	0	0
			43	33	4	6		
2	D	1	Total	C	N	O	0	0
			43	33	4	6		
2	E	1	Total	C	N	O	0	0
			43	33	4	6		
2	F	1	Total	C	N	O	0	0
			43	33	4	6		
2	G	1	Total	C	N	O	0	0
			43	33	4	6		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	H	1	43	33	4	6	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	61	Total	O	0	0
			61	61		
3	B	16	Total	O	0	0
			16	16		
3	C	44	Total	O	0	0
			44	44		
3	D	14	Total	O	0	0
			14	14		
3	E	15	Total	O	0	0
			15	15		
3	F	7	Total	O	0	0
			7	7		
3	G	37	Total	O	0	0
			37	37		
3	H	7	Total	O	0	0
			7	7		




• Molecule 1: Bacteriophytochrome



• Molecule 1: Bacteriophytochrome



- Molecule 1: Bacteriophytochrome



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	154.38Å 162.98Å 436.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.48 – 2.55 49.49 – 2.53	Depositor EDS
% Data completeness (in resolution range)	82.9 (48.48-2.55) 81.3 (49.49-2.53)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 2.51Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, R_{free}	0.215 , 0.258 0.216 , 0.256	Depositor DCC
R_{free} test set	7397 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	56.2	Xtrriage
Anisotropy	0.538	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 56.8	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	1 of 148657 reflections (0.001%)	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	30788	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3874	0.48	0/5258
1	B	0.24	0/3789	0.45	0/5144
1	C	0.25	0/3897	0.46	0/5289
1	D	0.24	0/3884	0.45	0/5269
1	E	0.23	0/3874	0.45	0/5258
1	F	0.23	0/3797	0.45	0/5155
1	G	0.25	0/3904	0.47	0/5299
1	H	0.24	0/3893	0.44	0/5284
All	All	0.24	0/30912	0.46	0/41956

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	407	ASP	Peptide
1	A	441	PRO	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	92	GLU	Peptide
1	C	62	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3790	0	3734	104	0
1	B	3709	0	3641	119	0
1	C	3812	0	3750	108	0
1	D	3801	0	3739	122	0
1	E	3790	0	3736	124	0
1	F	3717	0	3652	125	0
1	G	3816	0	3757	116	0
1	H	3808	0	3747	115	0
2	A	43	0	30	4	0
2	B	43	0	31	8	0
2	C	43	0	31	14	0
2	D	43	0	31	5	0
2	E	43	0	30	7	0
2	F	43	0	31	9	0
2	G	43	0	30	5	0
2	H	43	0	31	6	0
3	A	61	0	0	4	0
3	B	16	0	0	1	0
3	C	44	0	0	4	0
3	D	14	0	0	6	0
3	E	15	0	0	4	0
3	F	7	0	0	1	0
3	G	37	0	0	0	0
3	H	7	0	0	0	0
All	All	30788	0	30001	925	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 925 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:70:ARG:HG3	1:F:70:ARG:HH11	1.00	1.10
1:H:444:LEU:HD23	1:H:444:LEU:H	1.26	0.96
1:F:70:ARG:HG3	1:F:70:ARG:NH1	1.76	0.93
1:G:84:ASN:HD22	1:G:85:SER:H	1.02	0.92
1:D:112:ILE:HD13	1:D:112:ILE:H	1.36	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	475/505 (94%)	442 (93%)	30 (6%)	3 (1%)	30	48
1	B	464/505 (92%)	436 (94%)	28 (6%)	0	100	100
1	C	478/505 (95%)	455 (95%)	20 (4%)	3 (1%)	30	48
1	D	474/505 (94%)	451 (95%)	23 (5%)	0	100	100
1	E	475/505 (94%)	452 (95%)	22 (5%)	1 (0%)	52	73
1	F	465/505 (92%)	436 (94%)	27 (6%)	2 (0%)	39	60
1	G	479/505 (95%)	460 (96%)	18 (4%)	1 (0%)	52	73
1	H	477/505 (94%)	444 (93%)	28 (6%)	5 (1%)	19	33
All	All	3787/4040 (94%)	3576 (94%)	196 (5%)	15 (0%)	39	60

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	LEU
1	H	444	LEU
1	H	445	LEU
1	E	443	LYS
1	A	441	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	408/431 (95%)	377 (92%)	31 (8%)	16	29
1	B	400/431 (93%)	365 (91%)	35 (9%)	12	22
1	C	410/431 (95%)	376 (92%)	34 (8%)	14	25
1	D	409/431 (95%)	384 (94%)	25 (6%)	23	40
1	E	408/431 (95%)	369 (90%)	39 (10%)	10	18
1	F	401/431 (93%)	366 (91%)	35 (9%)	13	22
1	G	411/431 (95%)	380 (92%)	31 (8%)	17	30
1	H	410/431 (95%)	370 (90%)	40 (10%)	10	17
All	All	3257/3448 (94%)	2987 (92%)	270 (8%)	14	25

5 of 270 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	444	LEU
1	E	364	LEU
1	H	230	ASN
1	E	17	ILE
1	E	178	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 87 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	308	GLN
1	E	394	ASN
1	H	198	GLN
1	D	496	HIS
1	E	137	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BLA	A	900	1	35,46,46	2.80	14 (40%)	43,67,67	1.28	4 (9%)
2	BLA	B	900	1	35,46,46	2.89	14 (40%)	43,67,67	1.46	5 (11%)
2	BLA	C	900	1	35,46,46	2.81	16 (45%)	43,67,67	1.26	3 (6%)
2	BLA	D	900	1	35,46,46	2.86	14 (40%)	43,67,67	1.38	3 (6%)
2	BLA	E	900	1	35,46,46	2.91	14 (40%)	43,67,67	1.44	5 (11%)
2	BLA	F	900	1	35,46,46	2.80	13 (37%)	43,67,67	1.28	4 (9%)
2	BLA	G	900	1	35,46,46	2.86	13 (37%)	43,67,67	1.37	6 (13%)
2	BLA	H	900	1	35,46,46	2.83	13 (37%)	43,67,67	1.34	4 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BLA	A	900	1	-	1/22/74/74	0/4/4/4
2	BLA	B	900	1	-	1/22/74/74	0/4/4/4
2	BLA	C	900	1	-	1/22/74/74	0/4/4/4
2	BLA	D	900	1	-	1/22/74/74	0/4/4/4
2	BLA	E	900	1	-	1/22/74/74	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BLA	F	900	1	-	1/22/74/74	0/4/4/4
2	BLA	G	900	1	-	1/22/74/74	0/4/4/4
2	BLA	H	900	1	-	1/22/74/74	0/4/4/4

The worst 5 of 111 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	900	BLA	C4D-C3D	-3.67	1.39	1.45
2	C	900	BLA	C4D-C3D	-3.58	1.39	1.45
2	G	900	BLA	C4D-C3D	-3.52	1.39	1.45
2	D	900	BLA	C4D-C3D	-3.51	1.39	1.45
2	H	900	BLA	C4D-C3D	-3.50	1.39	1.45

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	900	BLA	C4C-CHD-C1D	-5.48	113.88	128.06
2	E	900	BLA	C4C-CHD-C1D	-5.33	114.28	128.06
2	D	900	BLA	C4C-CHD-C1D	-4.81	115.61	128.06
2	H	900	BLA	C4C-CHD-C1D	-4.40	116.67	128.06
2	G	900	BLA	C4C-CHD-C1D	-4.27	117.02	128.06

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	900	BLA	C4A-CHB-C1B-NB
2	B	900	BLA	C4A-CHB-C1B-NB
2	D	900	BLA	C4A-CHB-C1B-NB
2	H	900	BLA	C4A-CHB-C1B-NB
2	E	900	BLA	C4A-CHB-C1B-NB

There are no ring outliers.

8 monomers are involved in 58 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	900	BLA	4	0
2	B	900	BLA	8	0
2	C	900	BLA	14	0
2	D	900	BLA	5	0
2	E	900	BLA	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	900	BLA	9	0
2	G	900	BLA	5	0
2	H	900	BLA	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	479/505 (94%)	0.17	11 (2%) 64 69	37, 58, 109, 160	0
1	B	470/505 (93%)	0.74	54 (11%) 6 7	48, 88, 136, 183	0
1	C	482/505 (95%)	0.25	13 (2%) 58 63	39, 61, 102, 147	0
1	D	480/505 (95%)	0.71	52 (10%) 8 8	55, 92, 144, 191	0
1	E	479/505 (94%)	0.58	40 (8%) 14 15	49, 92, 134, 191	0
1	F	471/505 (93%)	0.66	56 (11%) 6 6	41, 81, 144, 171	0
1	G	482/505 (95%)	0.11	8 (1%) 73 77	37, 63, 105, 165	1 (0%)
1	H	481/505 (95%)	0.92	74 (15%) 3 3	55, 95, 158, 207	0
All	All	3824/4040 (94%)	0.52	308 (8%) 15 16	37, 79, 138, 207	1 (0%)

The worst 5 of 308 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	59	LEU	9.6
1	H	51	PHE	9.2
1	H	442	GLU	8.6
1	B	453	ARG	8.3
1	H	30	VAL	8.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	BLA	E	900	43/43	0.93	0.30	1.82	72,92,104,113	0
2	BLA	A	900	43/43	0.96	0.17	0.81	34,49,64,77	0
2	BLA	G	900	43/43	0.95	0.18	0.55	37,58,68,89	0
2	BLA	C	900	43/43	0.96	0.18	0.54	34,54,68,85	0
2	BLA	H	900	43/43	0.92	0.21	0.52	64,83,93,105	0
2	BLA	B	900	43/43	0.94	0.21	0.20	60,76,95,107	0
2	BLA	F	900	43/43	0.96	0.17	0.13	39,60,70,84	0
2	BLA	D	900	43/43	0.95	0.16	-0.41	62,81,95,101	0

6.5 Other polymers [i](#)

There are no such residues in this entry.