



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:02 AM GMT

PDB ID : 3NNG
Title : Crystal structure of the F5/8 type C domain of Q5LFR2_BACFN protein from *Bacteroides fragilis*. Northeast Structural Genomics Consortium Target BfR258E
Authors : Vorobiev, S.; Su, M.; Dimaio, F.; Baker, D.; Seetharaman, J.; Janjua, J.; Xiao, R.; Ciccocanti, C.; Foote, E.L.; Lee, D.; Everett, J.K.; Nair, R.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2010-06-23
Resolution : 2.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5130 (2.20-2.16)
Clashscore	102246	5965 (2.20-2.16)
Ramachandran outliers	100387	5863 (2.20-2.16)
Sidechain outliers	100360	5864 (2.20-2.16)
RSRZ outliers	91569	5142 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	 73% 17% • 9%
1	B	168	 69% 18% • 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2529 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	0	0	0
			1216	776	205	232	3			
1	B	151	Total	C	N	O	S	0	0	0
			1202	767	202	230	3			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	178	MET	-	expression tag	UNP Q5LFR2
A	335	GLU	VAL	engineered	UNP Q5LFR2
A	338	LEU	-	expression tag	UNP Q5LFR2
A	339	GLU	-	expression tag	UNP Q5LFR2
A	340	HIS	-	expression tag	UNP Q5LFR2
A	341	HIS	-	expression tag	UNP Q5LFR2
A	342	HIS	-	expression tag	UNP Q5LFR2
A	343	HIS	-	expression tag	UNP Q5LFR2
A	344	HIS	-	expression tag	UNP Q5LFR2
A	345	HIS	-	expression tag	UNP Q5LFR2
B	178	MET	-	expression tag	UNP Q5LFR2
B	335	GLU	VAL	engineered	UNP Q5LFR2
B	338	LEU	-	expression tag	UNP Q5LFR2
B	339	GLU	-	expression tag	UNP Q5LFR2
B	340	HIS	-	expression tag	UNP Q5LFR2
B	341	HIS	-	expression tag	UNP Q5LFR2
B	342	HIS	-	expression tag	UNP Q5LFR2
B	343	HIS	-	expression tag	UNP Q5LFR2
B	344	HIS	-	expression tag	UNP Q5LFR2
B	345	HIS	-	expression tag	UNP Q5LFR2

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Ca 1	0	0
2	A	1	Total 1	Ca 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	64	Total 64	O 64	0	0
3	B	45	Total 45	O 45	0	0

- Molecule 1: uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.21Å 62.46Å 72.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.23 – 2.18 31.23 – 2.18	Depositor EDS
% Data completeness (in resolution range)	99.6 (31.23-2.18) 99.6 (31.23-2.18)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.32 (at 2.18Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6 _289)	Depositor
R, R_{free}	0.164 , 0.224 0.166 , 0.224	Depositor DCC
R_{free} test set	693 reflections (5.21%)	DCC
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.483	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 13991 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2529	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1252	0.54	0/1711
1	B	0.35	0/1238	0.51	0/1692
All	All	0.38	0/2490	0.53	0/3403

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1216	0	1148	20	0
1	B	1202	0	1134	34	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	64	0	0	1	0
3	B	45	0	0	2	0
All	All	2529	0	2282	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:SER:HB2	1:A:337:GLN:HG2	1.38	1.05
1:B:293:ARG:HH12	1:B:296:LYS:HD3	1.44	0.81
1:A:242:ASN:HD22	1:A:316:ARG:HE	1.30	0.79
1:B:294:ASN:ND2	1:B:296:LYS:HE2	1.97	0.79
1:A:192:SER:HB2	1:A:337:GLN:CG	2.16	0.71
1:B:296:LYS:O	1:B:299:ASP:HB2	1.92	0.69
1:B:269:ARG:HG3	1:B:322:PRO:HD3	1.75	0.69
1:B:190:ASN:H	1:B:190:ASN:HD22	1.42	0.68
1:B:242:ASN:HD22	1:B:316:ARG:HE	1.42	0.66
1:B:296:LYS:HD2	1:B:296:LYS:N	2.10	0.66
1:A:187:LYS:HB2	1:A:302:TYR:CZ	2.31	0.66
1:B:196:ARG:HD2	1:B:220:ILE:HG22	1.80	0.64
1:A:242:ASN:ND2	1:A:316:ARG:HE	1.94	0.64
1:A:253:GLY:HA2	1:A:305:ILE:HG12	1.81	0.63
1:B:242:ASN:ND2	1:B:316:ARG:HE	1.97	0.62
1:B:242:ASN:HD22	1:B:316:ARG:HB2	1.66	0.61
1:B:293:ARG:NH1	1:B:296:LYS:HD3	2.18	0.57
1:B:190:ASN:N	1:B:190:ASN:HD22	2.02	0.56
1:B:259:GLN:H	1:B:267:ASN:ND2	2.03	0.56
1:B:321:THR:HA	1:B:322:PRO:C	2.26	0.56
1:B:286:TYR:CD1	1:B:287:PRO:HD2	2.42	0.55
1:B:294:ASN:HD21	1:B:296:LYS:HE2	1.69	0.54
1:A:274:LYS:HE3	1:A:285:THR:OG1	2.07	0.54
1:B:204:SER:HB3	1:B:240:TRP:CD1	2.43	0.53
1:B:250:THR:HA	1:B:310:PRO:HA	1.91	0.52
1:A:313:LYS:HE2	3:B:578:HOH:O	2.09	0.52
1:B:293:ARG:HH11	1:B:296:LYS:HE3	1.74	0.52
1:A:259:GLN:HG3	1:A:267:ASN:HD22	1.76	0.49
1:B:308:PRO:HG2	1:B:336:LYS:HE2	1.95	0.48
1:A:321:THR:HA	1:A:322:PRO:C	2.32	0.48
1:A:272:GLU:HG3	3:A:513:HOH:O	2.15	0.47
1:A:273:ILE:HD12	1:A:315:PHE:CZ	2.50	0.47
1:A:242:ASN:HD22	1:A:316:ARG:HB2	1.80	0.47
1:B:275:VAL:HG22	1:B:284:VAL:HG22	1.96	0.46
1:B:293:ARG:NH1	1:B:296:LYS:HE3	2.30	0.46
1:B:196:ARG:HG2	1:B:199:TRP:CZ3	2.50	0.45
1:B:245:LEU:N	1:B:245:LEU:HD12	2.31	0.45
1:B:229:PHE:CD1	1:B:259:GLN:HG3	2.51	0.45
1:B:252:THR:HG22	1:B:334:TYR:O	2.16	0.45
1:B:196:ARG:HD2	1:B:220:ILE:CG2	2.46	0.45
1:B:259:GLN:OE1	1:B:261:ALA:N	2.51	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:PRO:HB2	1:A:191:TRP:CD1	2.53	0.44
1:B:229:PHE:HD1	1:B:259:GLN:HG3	1.83	0.44
1:B:275:VAL:CG2	1:B:284:VAL:HG22	2.47	0.43
1:B:317:ILE:O	1:B:317:ILE:HG23	2.18	0.43
1:B:255:SER:HB3	1:B:334:TYR:HE1	1.84	0.42
1:A:277:LYS:HB2	1:A:280:GLU:HG3	2.01	0.42
1:A:313:LYS:CE	3:B:578:HOH:O	2.67	0.41
1:B:293:ARG:NH1	1:B:296:LYS:CD	2.81	0.41
1:A:206:VAL:HG11	1:A:211:ASP:HA	2.02	0.41
1:B:253:GLY:HA2	1:B:305:ILE:HG12	2.03	0.41
1:A:204:SER:HB3	1:A:240:TRP:CD1	2.56	0.41
1:A:188:PRO:HG2	1:A:191:TRP:CE2	2.56	0.40
1:A:338:LEU:HD23	1:A:338:LEU:C	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/168 (90%)	148 (98%)	3 (2%)	0	100	100
1	B	149/168 (89%)	145 (97%)	4 (3%)	0	100	100
All	All	300/336 (89%)	293 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/143 (89%)	123 (97%)	4 (3%)	47	56
1	B	126/143 (88%)	121 (96%)	5 (4%)	38	44
All	All	253/286 (88%)	244 (96%)	9 (4%)	42	49

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	268	LEU
1	A	269	ARG
1	A	280	GLU
1	A	296	LYS
1	B	190	ASN
1	B	259	GLN
1	B	268	LEU
1	B	294	ASN
1	B	296	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	242	ASN
1	A	267	ASN
1	B	190	ASN
1	B	242	ASN
1	B	267	ASN
1	B	294	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	153/168 (91%)	-0.47	0 100 100	13, 22, 42, 61	0
1	B	151/168 (89%)	-0.13	4 (2%) 59 61	16, 31, 54, 70	0
All	All	304/336 (90%)	-0.30	4 (1%) 79 80	13, 26, 51, 70	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	189	GLY	3.7
1	B	190	ASN	2.4
1	B	191	TRP	2.3
1	B	337	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CA	B	401	1/1	0.98	0.06	-1.75	37,37,37,37	0
2	CA	A	402	1/1	0.98	0.04	-4.02	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.