



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 04:03 PM BST

PDB ID : 1NVL
Title : RDC-refined NMR structure of bovine Acyl-coenzyme A Binding Protein, ACBP, in complex with palmitoyl-coenzyme A
Authors : Lerche, M.H.; Kragelund, B.B.; Redfield, C.; Poulsen, F.M.
Deposited on : 2003-02-04

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	86	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
2	A	COA	87	14	-

2 Ensemble composition and analysis

This entry contains 20 models. Model 12 is the overall representative, medoid model (most similar to other models). The authors have identified model 20 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:86 (84)	0.47	12

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 19
2	4, 17, 20
3	16, 18
Single-model clusters	1


3 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1518 atoms, of which 755 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Acyl-CoA-binding protein.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	86	1391	445	693	114	136	3	0

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).



Mol	Chain	Residues	Atoms						
			Total	C	H	N	O	P	S
2	A	1	79	21	31	7	16	3	1

- Molecule 3 is PALMITIC ACID (three-letter code: PLM) (formula: $C_{16}H_{32}O_2$).



Mol	Chain	Residues	Atoms			
			Total	C	H	O
3	A	1	48	16	31	1

5 Refinement protocol and experimental data overview

The models were refined using the following method: *SIMULATED ANNEALING FOLLOWED BY RESTRAINED MOLECULAR DYNAMICS IN FULL CHARMM FORCE FIELD*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	modified 3.8

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: COA, PLM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.45±0.02	1±1/697 (0.1±0.1%)	1.75±0.04	15±3/930 (1.6±0.3%)
All	All	1.45	17/13940 (0.1%)	1.75	294/18600 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.9±0.2
All	All	0	19

5 of 6 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	43	ARG	CZ-NH1	-5.84	1.25	1.33	19	3
1	A	43	ARG	CZ-NH2	-5.50	1.25	1.33	10	1
1	A	67	GLU	CB-CG	5.50	1.62	1.52	10	4
1	A	55	TRP	CG-CD2	-5.34	1.34	1.43	12	5
1	A	23	GLU	CB-CG	5.27	1.62	1.52	14	1

5 of 61 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	43	ARG	NE-CZ-NH2	-11.62	114.49	120.30	7	14
1	A	55	TRP	CD1-NE1-CE2	10.21	118.19	109.00	3	20
1	A	38	ASP	CB-CG-OD2	-9.95	109.34	118.30	3	9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	58	TRP	CD1-NE1-CE2	9.93	117.94	109.00	16	20
1	A	55	TRP	NE1-CE2-CZ2	9.22	140.54	130.40	2	18

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	43	ARG	Sidechain	19

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	683	678	678	29±5
2	A	48	31	31	7±2
3	A	17	31	31	1±1
All	All	14960	14800	14799	612

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

5 of 353 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:A:87:COA:C6P	2:A:87:COA:C5P	1.02	2.36	7	1
2:A:87:COA:C6P	2:A:87:COA:C7P	0.88	2.52	7	1
2:A:87:COA:C7P	2:A:87:COA:C5P	0.81	2.58	7	1
1:A:33:GLN:NE2	1:A:58:TRP:CD1	0.75	2.55	15	2
2:A:87:COA:C5B	2:A:87:COA:H143	0.71	2.15	14	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	83/86 (97%)	62±3 (74±4%)	16±3 (19±4%)	5±2 (6±2%)	3	20
All	All	1660/1720 (97%)	1233 (74%)	321 (19%)	106 (6%)	3	20

5 of 21 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	3	ALA	13
1	A	44	PRO	12
1	A	16	LYS	10
1	A	18	LYS	9
1	A	48	ASP	8

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	70/72 (97%)	62±2 (88±3%)	8±2 (12±3%)	11	54
All	All	1400/1440 (97%)	1239 (88%)	161 (12%)	11	54

5 of 39 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	12	VAL	18
1	A	75	ASP	11
1	A	52	LYS	10
1	A	78	GLU	10
1	A	18	LYS	9

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	COA	A	87	3	41,50,50	1.44±1.62	0±0 (0±1%)
3	PLM	A	88	2	16,16,17	0.79±0.08	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	COA	A	87	3	49,75,75	1.74±0.83	0±1 (0±1%)
3	PLM	A	88	2	15,15,17	0.82±0.10	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	87	3	1±0,1,13,13	0±0,44,64,64	0±0,3,3,3
3	PLM	A	88	2	-	0±0,13,14,15	0±0,0,0,0

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	87	COA	C6P-C5P	43.38	2.36	1.51	7	1
2	A	87	COA	C7P-C6P	31.96	2.52	1.51	7	1

5 of 6 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	A	87	COA	C7P-C6P-C5P	28.31	63.67	112.22	7	2
2	A	87	COA	O5P-C5P-C6P	16.70	92.98	121.97	7	1
2	A	87	COA	C6P-C5P-N4P	14.18	141.13	116.46	7	1
2	A	87	COA	C2P-C3P-N4P	9.37	93.89	112.44	1	1
2	A	87	COA	CDP-CBP-CCP	5.97	116.24	108.50	16	2

All unique chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
2	A	87	COA	CAP	14

All unique torsion outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
2	A	87	COA	CAP-C9P-N8P-C7P	2

There are no ring outliers.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided