



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:47 AM GMT

PDB ID : 2O8G
Title : Rat pp1c gamma complexed with mouse inhibitor-2
Authors : Hurley, T.D.
Deposited on : 2006-12-12
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	 % 74% 15% • 10%
1	B	329	 2% 74% 15% • 10%
2	I	206	 14% 20% 7% • 71%
2	J	206	 11% 17% 10% • 72%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5987 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein phosphatase PP1-gamma catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	295	2380	1527	399	436	18	0	0	0
1	B	295	2380	1527	399	436	18	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	INITIATING METHIONINE	UNP P63088
A	-4	HIS	-	EXPRESSION TAG	UNP P63088
A	-3	HIS	-	EXPRESSION TAG	UNP P63088
A	-2	HIS	-	EXPRESSION TAG	UNP P63088
A	-1	HIS	-	EXPRESSION TAG	UNP P63088
A	0	HIS	-	EXPRESSION TAG	UNP P63088
A	1	HIS	-	EXPRESSION TAG	UNP P63088
B	-5	MET	-	INITIATING METHIONINE	UNP P63088
B	-4	HIS	-	EXPRESSION TAG	UNP P63088
B	-3	HIS	-	EXPRESSION TAG	UNP P63088
B	-2	HIS	-	EXPRESSION TAG	UNP P63088
B	-1	HIS	-	EXPRESSION TAG	UNP P63088
B	0	HIS	-	EXPRESSION TAG	UNP P63088
B	1	HIS	-	EXPRESSION TAG	UNP P63088

- Molecule 2 is a protein called Protein phosphatase inhibitor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	I	59	495	309	93	91	2	0	0	0
2	J	58	487	305	92	88	2	0	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mn 1	0	0
3	A	1	Total 1	Mn 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	111	Total 111	O 111	0	0
4	B	99	Total 99	O 99	0	0
4	I	18	Total 18	O 18	0	0
4	J	15	Total 15	O 15	0	0

LEU
GLN
HIS
LYS
SER
GLN
SER
SER

• Molecule 2: Protein phosphatase inhibitor 2



MET ALA ALA MET ASP THR ASP SER HIS ARG PRO ILE K12 M17 LYS THR SER ALA ALA SER PRO PRO VAL PRO VAL ASP SER ALA GLN GLN PRO ARG PRO ILE VAL GLU LEU SER LYS K44 S45 K48 D49 E50 M51 N52 I53 L54 A55 T56 TYR HIS PRO ALA ASP LYS ASP TYR

GLY LEU MET ASN ILE ASP GLU PRO ASN THR PRO TYR HIS ASN MET ILE GLY ASP ASP ASP TYR SER ASP SER GLU ASN GLN VAL MET THR ASP MET THR PRO ASP ILE LEU ALA LYS LEU ALA ALA GLU GLY SER VAL GLU PRO LYS TYR ARG THR ARG GLN GLU SER ASP GLY

GLU GLU ASP ASP ASP S130 S131 P132 E133 E134 R135 E136 K137 K138 R139 E142 M143 K144 L153 K156 K164 D165 L166 H167 D168 ASP ASP GLU ASP GLU VAL MET THR ALA ASP GLY ASP MET MET ASN VAL GLU SER SER SER SER GLN GLY THR THR THR SER ASP HIS LEU

GLN
HIS
LYS
SER
GLN
SER
SER

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.27Å 103.23Å 149.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.80 – 2.50 48.81 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (48.80-2.50) 99.3 (48.81-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.194 , 0.235 0.190 , 0.230	Depositor DCC
R_{free} test set	2622 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	42.8	Xtrriage
Anisotropy	0.342	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.8	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 52097 reflections	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5987	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/2434	0.77	7/3286 (0.2%)
1	B	0.47	0/2434	0.77	7/3286 (0.2%)
2	I	0.46	0/499	0.71	2/658 (0.3%)
2	J	0.48	0/491	0.69	1/647 (0.2%)
All	All	0.48	0/5858	0.76	17/7877 (0.2%)

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	286	ASP	CB-CG-OD2	7.49	125.04	118.30
1	A	92	ASP	CB-CG-OD2	7.12	124.71	118.30
1	B	286	ASP	CB-CG-OD2	6.95	124.56	118.30
1	B	92	ASP	CB-CG-OD2	6.64	124.28	118.30
2	J	168	ASP	CB-CG-OD2	6.25	123.93	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2380	0	2351	16	0
1	B	2380	0	2351	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	I	495	0	507	2	0
2	J	487	0	503	5	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	111	0	0	4	0
4	B	99	0	0	4	0
4	I	18	0	0	2	0
4	J	15	0	0	2	0
All	All	5987	0	5712	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:143:MET:HG2	4:I:222:HOH:O	2.00	0.61
2:J:132:PRO:HA	4:J:221:HOH:O	1.99	0.61
1:B:131:ASN:HB3	1:B:149:TRP:HE1	1.70	0.56
1:A:94:VAL:O	1:A:95:ASP:HB2	2.06	0.55
1:A:131:ASN:HB3	1:A:149:TRP:HE1	1.74	0.52

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/329 (89%)	279 (95%)	14 (5%)	0	100	100
1	B	293/329 (89%)	277 (94%)	16 (6%)	0	100	100
2	I	53/206 (26%)	53 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	J	52/206 (25%)	51 (98%)	1 (2%)	0	100	100
All	All	691/1070 (65%)	660 (96%)	31 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/291 (90%)	241 (92%)	20 (8%)	16	30
1	B	261/291 (90%)	244 (94%)	17 (6%)	21	39
2	I	54/181 (30%)	39 (72%)	15 (28%)	0	0
2	J	53/181 (29%)	37 (70%)	16 (30%)	0	0
All	All	629/944 (67%)	561 (89%)	68 (11%)	8	15

5 of 68 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	164	LYS
1	B	26	LYS
2	J	153	LEU
2	I	165	ASP
2	I	168	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
2	I	140	GLN
1	B	294	GLN
1	B	181	GLN
1	A	294	GLN
1	B	99	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/329 (89%)	-0.27	4 (1%) 78 80	31, 40, 59, 75	0
1	B	295/329 (89%)	-0.13	5 (1%) 73 76	32, 43, 63, 78	0
2	I	59/206 (28%)	2.20	28 (47%) 0 0	35, 69, 103, 103	0
2	J	58/206 (28%)	1.82	22 (37%) 0 0	34, 72, 100, 100	0
All	All	707/1070 (66%)	0.17	59 (8%) 14 15	31, 43, 90, 103	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	54	LEU	11.2
2	I	51	MET	8.6
2	I	56	THR	8.2
2	J	54	LEU	7.7
2	I	167	HIS	7.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MN	B	324	1/1	1.00	0.05	-10.52	52,52,52,52	0
3	MN	A	324	1/1	0.99	0.04	-10.98	51,51,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.