



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 06:20 PM GMT

PDB ID : 1O94
Title : Ternary complex between trimethylamine dehydrogenase and electron transferring flavoprotein
Authors : Leys, D.; Basran, J.; Talfournier, F.; Sutcliffe, M.J.; Scrutton, N.S.
Deposited on : 2002-12-11
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	729	 90% 9% •
1	B	729	 87% 11% •
2	C	264	 68% 16% • • 12%
2	E	264	 73% 12% 5% 11%
3	D	320	 48% 10% • 41%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	320	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SF4	B	1732	-	-	X	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 19925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRIMETHYLAMINE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	729	Total	C	N	O	S	0	0	0
			5692	3589	996	1079	28			
1	B	729	Total	C	N	O	S	0	0	0
			5676	3583	994	1071	28			


- Molecule 2 is a protein called ELECTRON TRANSFER FLAVOPROTEIN BETA-SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	233	Total	C	N	O	S	0	0	0
			1749	1097	301	341	10			
2	E	236	Total	C	N	O	S	0	0	0
			1751	1102	299	340	10			

- Molecule 3 is a protein called ELECTRON TRANSFER FLAVOPROTEIN ALPHA-SUBUNIT.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	189	Total	C	N	O	0	0	0
			1354	857	230	267			
3	F	189	Total	C	N	O	0	0	0
			1376	870	232	274			

- Molecule 4 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
4	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		


- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
5	B	1	27	10	5	10	2	0	0

- Molecule 6 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
6	A	1	8	4	4	0	0
6	B	1	8	4	4	0	0

- Molecule 7 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
7	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	718	Total	O	0	0
			718	718		
8	B	831	Total	O	0	0
			831	831		
8	C	178	Total	O	0	0
			178	178		
8	D	73	Total	O	0	0
			73	73		
8	E	198	Total	O	0	0
			198	198		
8	F	151	Total	O	0	0
			151	151		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TRIMETHYLAMINE DEHYDROGENASE



- Molecule 1: TRIMETHYLAMINE DEHYDROGENASE



- Molecule 2: ELECTRON TRANSFER FLAVOPROTEIN BETA-SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.93Å 211.54Å 125.97Å 90.00° 100.03° 90.00°	Depositor
Resolution (Å)	19.88 – 2.00 19.90 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.88-2.00) 94.8 (19.90-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.1.08	Depositor
R, R_{free}	0.172 , 0.212 0.175 , 0.213	Depositor DCC
R_{free} test set	9811 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	28.7	Xtrriage
Anisotropy	0.550	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.2	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	0 of 194762 reflections	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	19925	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, AMP, SF4, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.04	1/5834 (0.0%)	1.02	29/7918 (0.4%)
1	B	1.11	8/5818 (0.1%)	1.03	31/7902 (0.4%)
2	C	1.06	2/1772 (0.1%)	1.12	18/2402 (0.7%)
2	E	0.97	1/1775 (0.1%)	1.21	18/2408 (0.7%)
3	D	0.72	0/1378	0.90	4/1884 (0.2%)
3	F	0.87	0/1400	0.97	6/1913 (0.3%)
All	All	1.03	12/17977 (0.1%)	1.04	106/24427 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
2	C	0	1
3	D	0	1
3	F	0	1
All	All	0	4

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	54	ASP	C-N	21.72	1.83	1.34
1	B	601	ARG	CG-CD	8.79	1.74	1.51
2	E	110	LYS	CE-NZ	8.01	1.69	1.49
2	C	110	LYS	CE-NZ	7.27	1.67	1.49
1	A	251	VAL	CB-CG2	-6.27	1.39	1.52

The worst 5 of 106 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	162	ARG	NE-CZ-NH2	-16.61	112.00	120.30
2	E	162	ARG	NE-CZ-NH1	14.90	127.75	120.30
1	A	157	ARG	NE-CZ-NH1	-14.87	112.86	120.30
2	E	86	ARG	NE-CZ-NH2	-13.81	113.39	120.30
2	C	54	ASP	C-N-CA	13.74	156.06	121.70

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	167	TYR	Peptide
2	C	156	ASN	Peptide
3	D	109	LYS	Peptide
3	F	110	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5692	0	5444	43	1
1	B	5676	0	5423	60	1
2	C	1749	0	1722	48	1
2	E	1751	0	1716	45	1
3	D	1354	0	1311	28	0
3	F	1376	0	1358	18	0
4	A	31	0	18	0	0
4	B	31	0	18	1	0
5	A	27	0	12	0	0
5	B	27	0	12	0	0
6	A	8	0	0	0	0
6	B	8	0	0	2	0
7	C	23	0	12	2	0
7	E	23	0	12	1	0
8	A	718	0	0	22	0
8	B	831	0	0	31	0
8	C	178	0	0	9	0
8	D	73	0	0	2	0
8	E	198	0	0	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	F	151	0	0	3	0
All	All	19925	0	17058	225	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 225 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:110:LYS:CE	2:E:110:LYS:NZ	1.69	1.55
1:B:253:MET:CE	1:B:253:MET:SD	2.04	1.45
1:B:229:TYR:CD1	8:B:2325:HOH:O	1.75	1.35
2:C:54:ASP:C	2:C:56:ASP:N	1.84	1.30
1:A:11:GLU:HG2	8:A:2015:HOH:O	1.31	1.26

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:GLU:OE2	2:E:156:ASN:OD1 4_545	1.97	0.23
1:B:11:GLU:OE2	2:C:156:ASN:OD1 4_556	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	727/729 (100%)	702 (97%)	25 (3%)	0	100 100
1	B	727/729 (100%)	700 (96%)	27 (4%)	0	100 100
2	C	229/264 (87%)	222 (97%)	4 (2%)	3 (1%)	15 7
2	E	234/264 (89%)	230 (98%)	4 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	D	187/320 (58%)	179 (96%)	7 (4%)	1 (0%)	34 26
3	F	187/320 (58%)	184 (98%)	3 (2%)	0	100 100
All	All	2291/2626 (87%)	2217 (97%)	70 (3%)	4 (0%)	52 48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	56	ASP
2	C	157	LYS
3	D	31	SER
2	C	145	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	584/602 (97%)	574 (98%)	10 (2%)	68 71
1	B	578/602 (96%)	565 (98%)	13 (2%)	60 62
2	C	180/216 (83%)	171 (95%)	9 (5%)	30 24
2	E	177/216 (82%)	171 (97%)	6 (3%)	44 41
3	D	143/258 (55%)	134 (94%)	9 (6%)	22 16
3	F	150/258 (58%)	141 (94%)	9 (6%)	24 17
All	All	1812/2152 (84%)	1756 (97%)	56 (3%)	47 46

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	56	ASP
2	C	231	ARG
3	F	66	SER
2	C	89	ASP
2	C	156	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	199	GLN
3	D	137	GLN
3	F	124	GLN
3	D	44	GLN
3	D	173	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FMN	A	1730	1	32,33,33	1.65	5 (15%)	34,50,50	2.59	10 (29%)
5	ADP	A	1731	-	24,29,29	1.17	1 (4%)	23,45,45	2.35	3 (13%)
6	SF4	A	1732	1	0,12,12	0.00	-	0,24,24	0.00	-
4	FMN	B	1730	1	32,33,33	1.59	6 (18%)	34,50,50	2.78	7 (20%)
5	ADP	B	1731	-	24,29,29	1.22	3 (12%)	23,45,45	2.39	4 (17%)
6	SF4	B	1732	1	0,12,12	0.00	-	0,24,24	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	AMP	C	1236	-	22,25,25	1.35	3 (13%)	22,38,38	3.56	8 (36%)
7	AMP	E	1237	-	22,25,25	1.75	5 (22%)	22,38,38	3.36	8 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FMN	A	1730	1	-	0/18/18/18	0/3/3/3
5	ADP	A	1731	-	-	0/12/32/32	0/3/3/3
6	SF4	A	1732	1	-	0/0/48/48	0/6/5/5
4	FMN	B	1730	1	-	0/18/18/18	0/3/3/3
5	ADP	B	1731	-	-	0/12/32/32	0/3/3/3
6	SF4	B	1732	1	-	0/0/48/48	0/6/5/5
7	AMP	C	1236	-	-	0/6/26/26	0/3/3/3
7	AMP	E	1237	-	-	0/6/26/26	0/3/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1730	FMN	C6-C5A	-3.31	1.36	1.41
4	B	1730	FMN	C6-C5A	-3.23	1.36	1.41
7	E	1237	AMP	C2'-C1'	-2.87	1.49	1.53
7	E	1237	AMP	O4'-C4'	-2.37	1.39	1.45
7	E	1237	AMP	O5'-C5'	-2.32	1.35	1.44

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	1236	AMP	N3-C2-N1	-13.46	118.29	128.87
7	E	1237	AMP	N3-C2-N1	-12.62	118.96	128.87
5	A	1731	ADP	N3-C2-N1	-10.09	120.94	128.87
5	B	1731	ADP	N3-C2-N1	-10.08	120.95	128.87
4	B	1730	FMN	C4A-C4-N3	-6.29	115.30	123.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1730	FMN	1	0
6	B	1732	SF4	2	0
7	C	1236	AMP	2	0
7	E	1237	AMP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	729/729 (100%)	-0.50	0 100 100	21, 28, 40, 47	0
1	B	729/729 (100%)	-0.58	3 (0%) 93 93	18, 26, 38, 53	0
2	C	233/264 (88%)	-0.22	3 (1%) 79 80	25, 35, 56, 66	0
2	E	236/264 (89%)	-0.20	5 (2%) 67 67	25, 33, 50, 58	0
3	D	189/320 (59%)	0.52	22 (11%) 6 7	32, 50, 66, 76	0
3	F	189/320 (59%)	-0.15	6 (3%) 51 52	27, 36, 52, 63	0
All	All	2305/2626 (87%)	-0.35	39 (1%) 73 73	18, 30, 53, 76	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	177	PRO	6.2
3	D	125	GLY	5.1
3	D	31	SER	4.9
3	F	32	GLY	4.6
3	D	32	GLY	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	SF4	B	1732	8/8	0.87	0.14	2.39	28,29,43,49	0
4	FMN	B	1730	31/31	0.95	0.11	1.55	19,23,26,28	0
4	FMN	A	1730	31/31	0.96	0.09	0.38	18,23,25,27	0
7	AMP	E	1237	23/23	0.97	0.09	-0.33	23,28,33,35	0
7	AMP	C	1236	23/23	0.97	0.08	-0.54	26,30,37,38	0
5	ADP	B	1731	27/27	0.99	0.06	-0.72	20,24,28,29	0
5	ADP	A	1731	27/27	0.98	0.07	-1.24	23,28,30,31	0
6	SF4	A	1732	8/8	0.96	0.07	-1.78	29,31,32,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.