



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:20 PM GMT

PDB ID : 1OKC  
Title : structure of mitochondrial ADP/ATP carrier in complex with carboxyatractyloside  
Authors : Pebay-Peyroula, E.; Dahout-Gonzalez, C.; Kahn, R.; Trezguet, V.; Lauquin, G.J.-M.; Brandolin, G.  
Deposited on : 2003-07-21  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CXT	A	401	X	-	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CDL	A	800	-	-	-	X
3	CDL	A	801	-	-	-	X
3	CDL	A	802	-	-	-	X
4	LDM	A	903	-	-	-	X
4	LDM	A	904	-	-	-	X
5	PC1	A	980	-	-	-	X
5	PC1	A	981	-	-	-	X
5	PC1	A	982	-	-	-	X

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 2686 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP, ATP CARRIER PROTEIN HEART ISOFORM T1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	0	0
			2254	1460	392	391	11			

- Molecule 2 is CARBOXYATRACTYLOSIDE (three-letter code: CXT) (formula:  $C_{31}H_{46}O_{18}S_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	0
			51	31	18	2		

- Molecule 3 is CARDIOLIPIN (three-letter code: CDL) (formula:  $C_{81}H_{156}O_{17}P_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			57	38	17	2		
3	A	1	Total	C	O	P	0	0
			49	30	17	2		
3	A	1	Total	C	O		0	0
			36	32	4			

- Molecule 4 is 3-LAURLAMIDO-N,N'-DIMETHYLPROPYLAMINOXYDE (three-letter code: LDM) (formula:  $C_{17}H_{36}N_2O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			18	14	2	2		
4	A	1	Total	C	N	O	0	0
			21	17	2	2		

- Molecule 5 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PC1) (formula:  $C_{44}H_{88}NO_8P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			28	18	1	8	1		
5	A	1	Total	C	N	O	P	0	0
			41	31	1	8	1		
5	A	1	Total	C	N	O	P	0	0
			27	17	1	8	1		
5	A	1	Total	C	O	P		0	0
			21	12	8	1			


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	83	Total	O	0	0
			83	83		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ADP, ATP CARRIER PROTEIN HEART ISOFORM T1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.44Å 83.46Å 49.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.99 – 2.20 24.74 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (14.99-2.20) 99.3 (24.74-2.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.81 (at 2.19Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.220 , 0.266 0.217 , 0.264	Depositor DCC
$R_{free}$ test set	894 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.2	Xtriage
Anisotropy	0.235	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 59.6	EDS
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 18595 reflections (0.011%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2686	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CDL, PC1, CXT, LDM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/2306	0.54	0/3113

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2254	0	2258	64	0
2	A	51	0	41	1	0
3	A	142	0	153	8	0
4	A	39	0	63	0	0
5	A	117	0	132	4	0
6	A	83	0	0	0	0
All	All	2686	0	2647	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:LEU:O	1:A:102:VAL:HG12	1.69	0.93
1:A:60:ILE:HG13	1:A:61:PRO:HD3	1.54	0.89
1:A:30:ARG:O	1:A:34:LEU:HD23	1.79	0.83
1:A:30:ARG:HH22	1:A:64:GLN:HE22	1.29	0.79
1:A:198:LYS:HA	1:A:202:PRO:HG2	1.67	0.75
3:A:802:CDL:H332	3:A:802:CDL:H151	1.71	0.73
1:A:200:MET:O	1:A:202:PRO:HD3	1.90	0.72
1:A:5:LEU:HG	1:A:9:LYS:HE3	1.73	0.71
1:A:200:MET:C	1:A:202:PRO:HD3	2.13	0.67
1:A:60:ILE:HG13	1:A:61:PRO:CD	2.25	0.64
1:A:274:TRP:HB2	3:A:801:CDL:HB61	1.78	0.63
1:A:73:ASN:HD22	1:A:76:ASN:HD22	1.47	0.63
1:A:154:THR:H	1:A:158:ASN:HD22	1.46	0.62
1:A:198:LYS:HA	1:A:202:PRO:CG	2.28	0.61
1:A:67:LEU:HD13	5:A:982:PC1:H31	1.82	0.61
1:A:203:ASP:O	1:A:205:LYS:N	2.35	0.59
1:A:47:GLU:N	1:A:47:GLU:OE1	2.36	0.59
1:A:195:ASP:O	1:A:198:LYS:HG2	2.06	0.56
1:A:45:SER:OG	1:A:48:LYS:HG3	2.07	0.55
1:A:8:LEU:O	1:A:12:LEU:HG	2.08	0.54
1:A:154:THR:H	1:A:158:ASN:ND2	2.04	0.54
1:A:32:LYS:NZ	1:A:36:GLN:HE22	2.06	0.54
1:A:59:ARG:O	1:A:63:GLU:HB2	2.08	0.53
1:A:203:ASP:O	1:A:204:PRO:C	2.47	0.53
1:A:231:ASP:O	1:A:235:ARG:HG2	2.09	0.53
1:A:12:LEU:O	1:A:16:VAL:HG23	2.10	0.52
1:A:177:ASN:OD1	3:A:802:CDL:H311	2.09	0.52
1:A:30:ARG:NH2	1:A:64:GLN:HE22	2.04	0.51
1:A:215:ILE:C	1:A:215:ILE:HD13	2.30	0.51
1:A:73:ASN:ND2	1:A:76:ASN:HD22	2.10	0.50
1:A:147:GLY:O	1:A:151:ARG:HG2	2.12	0.50
1:A:127:LEU:HG	2:A:401:CXT:H11	1.94	0.49
1:A:53:ILE:HG13	3:A:801:CDL:HB21	1.95	0.48
1:A:211:VAL:O	1:A:215:ILE:HG22	2.14	0.47
1:A:250:TYR:CD1	1:A:256:CYS:HA	2.50	0.47
1:A:200:MET:O	1:A:201:LEU:CB	2.62	0.47
1:A:30:ARG:HH22	1:A:64:GLN:NE2	2.06	0.46
1:A:211:VAL:HA	1:A:214:MET:CE	2.46	0.46
1:A:200:MET:O	1:A:201:LEU:HB3	2.16	0.45
1:A:74:LEU:HD22	1:A:78:ILE:CD1	2.47	0.45
1:A:80:TYR:C	1:A:80:TYR:CD1	2.89	0.45
1:A:80:TYR:CZ	1:A:84:GLN:HG3	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:GLY:O	1:A:56:CYS:HB2	2.17	0.45
1:A:113:ALA:HB2	5:A:981:PC1:H362	1.98	0.45
1:A:244:LYS:O	1:A:247:ASP:HB2	2.16	0.45
1:A:34:LEU:HD21	1:A:141:ALA:O	2.15	0.45
1:A:250:TYR:HE1	1:A:259:LYS:HB3	1.82	0.44
1:A:56:CYS:O	1:A:60:ILE:HG23	2.17	0.44
1:A:154:THR:HB	5:A:980:PC1:H132	2.00	0.44
1:A:36:GLN:NE2	1:A:235:ARG:HG3	2.33	0.44
1:A:46:ALA:HB3	1:A:47:GLU:OE1	2.18	0.44
1:A:225:LEU:HB3	3:A:801:CDL:H562	1.99	0.44
3:A:801:CDL:O1	5:A:983:PC1:H12	2.18	0.44
1:A:26:ALA:N	1:A:27:PRO:CD	2.81	0.43
1:A:181:GLN:HG2	3:A:802:CDL:H402	2.00	0.43
1:A:147:GLY:N	1:A:150:GLN:OE1	2.51	0.43
1:A:60:ILE:HG12	1:A:69:PHE:CE1	2.54	0.43
1:A:61:PRO:HG3	1:A:69:PHE:CE2	2.54	0.42
1:A:198:LYS:CA	1:A:202:PRO:HG2	2.44	0.42
1:A:67:LEU:HA	1:A:67:LEU:HD12	1.90	0.42
1:A:211:VAL:HA	1:A:214:MET:HE2	2.02	0.42
1:A:153:PHE:CD1	1:A:159:CYS:HA	2.53	0.42
1:A:288:VAL:HG23	1:A:289:LEU:N	2.35	0.42
1:A:77:VAL:HG11	3:A:800:CDL:H311	2.01	0.41
1:A:149:ALA:HB3	1:A:150:GLN:HE21	1.86	0.41
1:A:200:MET:C	1:A:202:PRO:CD	2.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	290/297 (98%)	281 (97%)	4 (1%)	5 (2%)	<b>11</b> <b>7</b>

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	201	LEU
1	A	202	PRO
1	A	204	PRO
1	A	205	LYS
1	A	210	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/234 (95%)	210 (94%)	13 (6%)	25	28

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	47	GLU
1	A	60	ILE
1	A	74	LEU
1	A	137	ARG
1	A	194	TYR
1	A	203	ASP
1	A	204	PRO
1	A	208	HIS
1	A	215	ILE
1	A	233	VAL
1	A	247	ASP
1	A	291	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	A	64	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	73	ASN
1	A	158	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	CXT	A	401	-	49,55,55	2.98	20 (40%)	66,89,89	2.95	19 (28%)
3	CDL	A	800	-	56,56,99	0.65	0	58,68,111	1.05	4 (6%)
3	CDL	A	801	-	48,48,99	0.82	2 (4%)	50,60,111	1.01	5 (10%)
3	CDL	A	802	-	35,35,99	0.75	2 (5%)	37,37,111	1.02	2 (5%)
4	LDM	A	903	-	17,17,20	4.21	3 (17%)	19,20,23	2.45	3 (15%)
4	LDM	A	904	-	20,20,20	3.84	3 (15%)	22,23,23	2.31	3 (13%)
5	PC1	A	980	-	27,27,53	2.12	2 (7%)	31,35,61	1.15	3 (9%)
5	PC1	A	981	-	40,40,53	1.73	2 (5%)	44,48,61	0.90	1 (2%)
5	PC1	A	982	-	26,26,53	1.85	2 (7%)	30,34,61	0.93	1 (3%)
5	PC1	A	983	-	20,20,53	2.18	3 (15%)	24,25,61	1.20	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CXT	A	401	-	1/1/20/20	0/24/120/120	0/3/5/5
3	CDL	A	800	-	-	0/67/67/110	0/0/0/0
3	CDL	A	801	-	-	0/58/58/110	0/0/0/0
3	CDL	A	802	-	-	0/36/36/110	0/0/0/0
4	LDM	A	903	-	-	0/16/16/19	0/0/0/0
4	LDM	A	904	-	-	0/19/19/19	0/0/0/0
5	PC1	A	980	-	-	0/31/31/57	0/0/0/0
5	PC1	A	981	-	-	0/44/44/57	0/0/0/0
5	PC1	A	982	-	-	0/29/29/57	0/0/0/0
5	PC1	A	983	-	-	0/21/21/57	0/0/0/0

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	903	LDM	O1-N1	-16.53	1.23	1.39
4	A	904	LDM	O1-N1	-16.35	1.24	1.39
2	A	401	CXT	O21-C35	-4.90	1.32	1.42
2	A	401	CXT	O2-C1	-3.81	1.31	1.41
4	A	904	LDM	CM2-N1	-3.27	1.44	1.49
4	A	903	LDM	CM2-N1	-3.26	1.44	1.49
2	A	401	CXT	O13-C7	-3.03	1.13	1.22
2	A	401	CXT	O4-C3	-2.79	1.40	1.46
4	A	903	LDM	CM1-N1	-2.47	1.45	1.49
4	A	904	LDM	CM1-N1	-2.20	1.46	1.49
2	A	401	CXT	O8-S1	-2.02	1.39	1.50
3	A	802	CDL	CA6-CA4	2.04	1.55	1.50
3	A	801	CDL	CA3-CA4	2.08	1.56	1.50
3	A	802	CDL	OA6-CA4	2.15	1.51	1.47
2	A	401	CXT	C21-C22	2.23	1.56	1.51
5	A	983	PC1	P-O13	2.37	1.64	1.55
2	A	401	CXT	C23-C22	2.59	1.56	1.51
2	A	401	CXT	O1-C1	2.65	1.48	1.41
2	A	401	CXT	C27-C26	2.75	1.59	1.53
2	A	401	CXT	C21-C30	2.88	1.59	1.54
3	A	801	CDL	CB3-CB4	2.91	1.58	1.50
5	A	982	PC1	O31-C31	3.49	1.51	1.33
5	A	983	PC1	O31-C31	3.71	1.52	1.33
2	A	401	CXT	O2-C22	4.12	1.50	1.44
2	A	401	CXT	C39-C30	4.60	1.62	1.54

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	CXT	C26-C25	4.66	1.61	1.53
2	A	401	CXT	C24-C38	5.23	1.60	1.50
2	A	401	CXT	C24-C37	5.26	1.60	1.50
2	A	401	CXT	C40-C36	5.74	1.44	1.32
2	A	401	CXT	C28-C29	5.90	1.68	1.55
2	A	401	CXT	O3-C7	6.03	1.52	1.34
2	A	401	CXT	C31-C29	6.07	1.64	1.53
5	A	980	PC1	O31-C31	6.14	1.51	1.33
5	A	981	PC1	O31-C31	6.38	1.52	1.33
2	A	401	CXT	C30-C25	7.45	1.68	1.56
5	A	982	PC1	O21-C21	7.47	1.56	1.34
5	A	981	PC1	O21-C21	7.56	1.57	1.34
5	A	983	PC1	O21-C21	7.89	1.58	1.34
5	A	980	PC1	O21-C21	7.96	1.58	1.34

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	903	LDM	CM2-N1-CM1	-8.11	99.68	108.83
4	A	904	LDM	CM2-N1-CM1	-7.89	99.93	108.83
2	A	401	CXT	O2-C22-C23	-7.17	86.64	107.83
2	A	401	CXT	O3-C7-C8	-6.42	99.10	111.54
5	A	980	PC1	C3-C2-C1	-4.56	101.41	112.07
2	A	401	CXT	C26-C25-C30	-4.55	105.91	110.94
5	A	983	PC1	C3-C2-C1	-3.92	102.89	112.07
5	A	981	PC1	C3-C2-C1	-3.59	103.66	112.07
2	A	401	CXT	C39-C30-C21	-3.47	103.38	108.87
2	A	401	CXT	O5-S2-O9	-3.45	95.67	106.86
5	A	982	PC1	C3-C2-C1	-2.74	105.66	112.07
3	A	800	CDL	CB6-CB4-CB3	-2.60	105.98	112.07
3	A	800	CDL	CA6-CA4-CA3	-2.55	106.12	112.07
2	A	401	CXT	O2-C1-O1	-2.52	104.30	110.68
2	A	401	CXT	O4-S1-O7	-2.47	98.84	106.86
5	A	980	PC1	C3-O31-C31	-2.26	110.52	116.85
2	A	401	CXT	O3-C2-C3	-2.19	103.87	108.23
3	A	801	CDL	O1-C1-CA2	-2.05	101.54	109.35
3	A	801	CDL	OB6-CB5-C51	2.01	115.91	111.53
2	A	401	CXT	C32-C33-C36	2.03	114.00	110.55
3	A	800	CDL	OB6-CB5-C51	2.06	116.01	111.53
3	A	801	CDL	CB4-OB6-CB5	2.08	122.88	117.89
2	A	401	CXT	O13-C7-C8	2.12	129.57	124.69
3	A	801	CDL	C52-C51-CB5	2.20	122.25	113.59

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	980	PC1	O21-C21-C22	2.29	116.50	111.53
2	A	401	CXT	O2-C1-C2	2.31	113.81	109.05
3	A	802	CDL	OA6-CA4-CA6	2.36	111.24	105.90
2	A	401	CXT	O3-C2-C1	2.48	114.19	108.48
2	A	401	CXT	C39-C30-C25	2.51	117.64	112.95
4	A	903	LDM	O1-N1-C1	2.82	113.45	110.27
2	A	401	CXT	C1-O1-C5	2.83	119.25	113.75
2	A	401	CXT	O3-C7-O13	2.85	131.33	123.67
4	A	904	LDM	O1-N1-C1	2.91	113.55	110.27
3	A	800	CDL	OB8-CB6-CB4	3.15	117.18	108.69
3	A	801	CDL	OB8-CB6-CB4	3.43	117.93	108.69
2	A	401	CXT	C1-C2-C3	3.43	117.97	110.75
3	A	802	CDL	OA6-CA4-CA3	4.12	117.18	107.88
2	A	401	CXT	O12-C6-C5	4.15	125.03	111.33
4	A	903	LDM	O1-N1-CM2	5.33	116.18	109.05
4	A	904	LDM	O1-N1-CM2	5.34	116.19	109.05
2	A	401	CXT	C3-O4-S1	7.44	132.95	118.77
2	A	401	CXT	O2-C22-C21	15.12	152.52	107.83

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	401	CXT	C1

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	CXT	1	0
3	A	800	CDL	1	0
3	A	801	CDL	4	0
3	A	802	CDL	3	0
5	A	980	PC1	1	0
5	A	981	PC1	1	0
5	A	982	PC1	1	0
5	A	983	PC1	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/297 (98%)	0.04	21 (7%) 18 18	20, 31, 76, 108	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	201	LEU	6.2
1	A	209	ILE	5.8
1	A	204	PRO	5.7
1	A	210	ILE	5.5
1	A	200	MET	5.4
1	A	288	VAL	4.0
1	A	208	HIS	4.0
1	A	291	ASP	3.8
1	A	203	ASP	3.8
1	A	202	PRO	3.8
1	A	2	ASP	3.7
1	A	194	TYR	3.3
1	A	211	VAL	3.2
1	A	205	LYS	3.2
1	A	206	ASN	2.9
1	A	290	TYR	2.8
1	A	207	VAL	2.8
1	A	7	PHE	2.7
1	A	214	MET	2.6
1	A	292	GLU	2.6
1	A	3	GLN	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	LDM	A	904	21/21	0.57	0.38	11.55	96,98,99,100	0
5	PC1	A	982	27/54	0.85	0.22	8.95	54,59,67,68	0
3	CDL	A	802	36/100	0.57	0.27	5.37	62,84,88,89	0
5	PC1	A	980	28/54	0.80	0.31	4.81	99,99,99,99	0
5	PC1	A	981	41/54	0.79	0.33	3.88	85,89,91,91	0
3	CDL	A	800	57/100	0.86	0.24	3.57	34,56,74,75	0
4	LDM	A	903	18/21	0.67	0.29	3.52	75,80,82,82	0
3	CDL	A	801	49/100	0.83	0.23	3.47	34,50,71,71	0
2	CXT	A	401	51/51	0.93	0.14	0.41	21,33,45,46	0
5	PC1	A	983	21/54	0.47	0.35	-	108,115,121,121	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.