



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:53 PM GMT

PDB ID : 4OKI  
Title : X-ray structure of the nucleotide-binding subdomain of the enoylreductase domain of PpsC from Mycobacterium tuberculosis  
Authors : Faille, A.; Mourey, L.; Pedelacq, J.D.  
Deposited on : 2014-01-22  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	<div> <div>2%</div> <div>72% 12% 16%</div> </div>
1	B	216	<div> <div>4%</div> <div>73% 9% 17%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NA	B	901	-	-	-	X
2	NA	B	904	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phthiocerol synthesis polyketide synthase type I PpsC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	3	0
			1369	865	242	257	5			
1	B	179	Total	C	N	O	S	0	4	0
			1348	855	236	253	4			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	636	MET	-	EXPRESSION TAG	UNP P96202
A	637	GLY	-	EXPRESSION TAG	UNP P96202
A	638	SER	-	EXPRESSION TAG	UNP P96202
A	639	SER	-	EXPRESSION TAG	UNP P96202
A	640	HIS	-	EXPRESSION TAG	UNP P96202
A	641	HIS	-	EXPRESSION TAG	UNP P96202
A	642	HIS	-	EXPRESSION TAG	UNP P96202
A	643	HIS	-	EXPRESSION TAG	UNP P96202
A	644	HIS	-	EXPRESSION TAG	UNP P96202
A	645	HIS	-	EXPRESSION TAG	UNP P96202
A	646	SER	-	EXPRESSION TAG	UNP P96202
A	647	SER	-	EXPRESSION TAG	UNP P96202
A	648	GLY	-	EXPRESSION TAG	UNP P96202
A	649	LEU	-	EXPRESSION TAG	UNP P96202
A	650	VAL	-	EXPRESSION TAG	UNP P96202
A	651	PRO	-	EXPRESSION TAG	UNP P96202
A	652	ARG	-	EXPRESSION TAG	UNP P96202
A	653	GLY	-	EXPRESSION TAG	UNP P96202
A	654	SER	-	EXPRESSION TAG	UNP P96202
A	655	HIS	-	EXPRESSION TAG	UNP P96202
A	656	MET	-	EXPRESSION TAG	UNP P96202
A	850	GLY	-	EXPRESSION TAG	UNP P96202
A	851	SER	-	EXPRESSION TAG	UNP P96202
B	636	MET	-	EXPRESSION TAG	UNP P96202
B	637	GLY	-	EXPRESSION TAG	UNP P96202

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	638	SER	-	EXPRESSION TAG	UNP P96202
B	639	SER	-	EXPRESSION TAG	UNP P96202
B	640	HIS	-	EXPRESSION TAG	UNP P96202
B	641	HIS	-	EXPRESSION TAG	UNP P96202
B	642	HIS	-	EXPRESSION TAG	UNP P96202
B	643	HIS	-	EXPRESSION TAG	UNP P96202
B	644	HIS	-	EXPRESSION TAG	UNP P96202
B	645	HIS	-	EXPRESSION TAG	UNP P96202
B	646	SER	-	EXPRESSION TAG	UNP P96202
B	647	SER	-	EXPRESSION TAG	UNP P96202
B	648	GLY	-	EXPRESSION TAG	UNP P96202
B	649	LEU	-	EXPRESSION TAG	UNP P96202
B	650	VAL	-	EXPRESSION TAG	UNP P96202
B	651	PRO	-	EXPRESSION TAG	UNP P96202
B	652	ARG	-	EXPRESSION TAG	UNP P96202
B	653	GLY	-	EXPRESSION TAG	UNP P96202
B	654	SER	-	EXPRESSION TAG	UNP P96202
B	655	HIS	-	EXPRESSION TAG	UNP P96202
B	656	MET	-	EXPRESSION TAG	UNP P96202
B	850	GLY	-	EXPRESSION TAG	UNP P96202
B	851	SER	-	EXPRESSION TAG	UNP P96202

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	5	Total Na 5 5	0	0
2	A	2	Total Na 2 2	0	0

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	158	Total	O	0	0
			158	158		
4	B	143	Total	O	0	0
			143	143		



- Molecule 1: Phthiocerol synthesis polyketide synthase type I PpsC



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.88Å 78.74Å 88.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.00 – 1.50 42.00 – 1.50	Depositor EDS
% Data completeness (in resolution range)	90.3 (42.00-1.50) 90.4 (42.00-1.50)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.22 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.182 , 0.218 0.191 , 0.224	Depositor DCC
$R_{free}$ test set	3445 reflections (5.92%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.2	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 61343 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3033	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.31	3/1393 (0.2%)	1.26	3/1881 (0.2%)
1	B	1.18	0/1378	1.14	2/1865 (0.1%)
All	All	1.25	3/2771 (0.1%)	1.20	5/3746 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	762	TYR	CE1-CZ	5.50	1.45	1.38
1	A	763	GLY	N-CA	5.39	1.54	1.46
1	A	693	SER	CB-OG	-5.26	1.35	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	749	VAL	CG1-CB-CG2	5.94	120.41	110.90
1	A	818	ASP	CB-CG-OD2	-5.80	113.08	118.30
1	B	686	LEU	CB-CG-CD1	-5.79	101.17	111.00
1	A	732	ARG	NE-CZ-NH1	5.77	123.19	120.30
1	A	762	TYR	CG-CD1-CE1	5.24	125.49	121.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1369	0	1399	16	0
1	B	1348	0	1382	13	0
2	A	2	0	0	0	0
2	B	5	0	0	0	0
3	A	8	0	14	0	0
4	A	158	0	0	5	0
4	B	143	0	0	2	0
All	All	3033	0	2795	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:698:VAL:HG23	1:B:766[A]:VAL:HG13	1.36	1.03
1:A:777:GLN:HG2	4:A:1051:HOH:O	1.73	0.87
1:A:749:VAL:CG1	1:A:778:ARG:HG3	2.14	0.78
1:B:698:VAL:CG2	1:B:766[A]:VAL:HG13	2.20	0.70
1:A:749:VAL:HG11	1:A:778:ARG:HG3	1.74	0.69
1:B:671:GLU:N	4:B:1109:HOH:O	2.30	0.64
1:A:749:VAL:HG13	1:A:778:ARG:HG3	1.79	0.63
1:B:698:VAL:HG23	1:B:766[A]:VAL:CG1	2.22	0.62
1:A:831:ARG:HG2	4:A:1147:HOH:O	1.99	0.61
1:B:730:ALA:O	1:B:734:MET:HG3	2.04	0.58
1:A:832:GLN:HG3	4:A:1127:HOH:O	2.04	0.57
1:A:705:GLY:HA3	4:A:1092:HOH:O	2.06	0.54
1:A:730:ALA:O	1:A:734:MET:HG3	2.08	0.53
1:B:803:LEU:HD22	1:B:804:GLY:O	2.10	0.51
1:A:670:ASN:N	4:A:1042:HOH:O	2.45	0.50
1:B:701:HIS:HD2	1:B:725:THR:OG1	1.95	0.49
1:B:843[A]:ASP:OD1	1:B:845:LYS:HG3	2.13	0.49
1:A:695:GLY:O	1:A:721[B]:ARG:CZ	2.63	0.47
1:A:701:HIS:HD2	1:A:725:THR:OG1	1.97	0.47
1:B:692:LEU:HD13	1:B:766[A]:VAL:HG11	1.97	0.46
1:A:794:LYS:H	1:A:794:LYS:HG2	1.53	0.45
1:B:716:LYS:HD2	1:B:849:LEU:HD11	2.00	0.44
1:A:749:VAL:HG21	1:A:774:GLU:HB3	2.00	0.43
1:B:747:ARG:HD2	4:B:1023:HOH:O	2.19	0.43
1:A:822:ASN:HD21	1:B:811:SER:CB	2.32	0.41
1:A:805:LEU:HD21	1:B:803:LEU:HB2	2.03	0.40
1:A:723:TYR:CD1	1:A:723:TYR:N	2.90	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/216 (84%)	181 (100%)	1 (0%)	0	100	100
1	B	181/216 (84%)	179 (99%)	2 (1%)	0	100	100
All	All	363/432 (84%)	360 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/168 (83%)	136 (97%)	4 (3%)	50	16
1	B	139/168 (83%)	133 (96%)	6 (4%)	35	7
All	All	279/336 (83%)	269 (96%)	10 (4%)	41	10

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	794	LYS
1	A	803	LEU
1	A	820	ASP
1	A	840	HIS
1	B	733	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	749	VAL
1	B	753	ASP
1	B	796	ASP
1	B	803	LEU
1	B	832	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	701	HIS
1	A	781	GLN
1	A	822	ASN
1	A	826	GLN
1	B	701	HIS
1	B	822	ASN
1	B	832	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 7 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MPD	A	902	-	6,7,7	0.31	0	6,10,10	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	902	-	-	0/5/5/5	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	181/216 (83%)	-0.10	4 (2%) 65 68	15, 23, 41, 50	0
1	B	179/216 (82%)	0.09	8 (4%) 37 38	16, 25, 53, 71	0
All	All	360/432 (83%)	-0.01	12 (3%) 50 52	15, 25, 48, 71	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	734	MET	5.1
1	B	797	VAL	4.3
1	B	672	ALA	4.2
1	B	730	ALA	3.0
1	A	730	ALA	2.7
1	A	766	VAL	2.7
1	B	673	ALA	2.6
1	B	671	GLU	2.6
1	A	768	LEU	2.5
1	B	733	GLU	2.4
1	A	699	LEU	2.4
1	B	798	TYR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NA	B	904	1/1	0.93	0.14	5.92	47,47,47,47	0
2	NA	B	901	1/1	0.95	0.17	5.53	40,40,40,40	0
2	NA	A	901	1/1	0.92	0.10	1.50	45,45,45,45	0
3	MPD	A	902	8/8	0.94	0.09	1.01	40,49,53,53	0
2	NA	A	903	1/1	0.96	0.15	0.45	46,46,46,46	0
2	NA	B	902	1/1	0.96	0.12	0.38	43,43,43,43	0
2	NA	B	903	1/1	0.96	0.07	-3.05	41,41,41,41	0
2	NA	B	905	1/1	0.97	0.03	-	36,36,36,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.