



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:18 AM GMT

PDB ID : 3ONY
Title : Crystal Structure of P Domain from Norwalk Virus Strain Vietnam 026 in complex with Fucose
Authors : Hansman, G.S.; Biertumpfel, C.; Chen, L.; Georgiev, I.; McLellan, J.S.; Katayama, K.; Kwong, P.D.
Deposited on : 2010-08-30
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	<div> <div>7%</div> <div>92%</div> <div>6%</div> </div>
1	B	318	<div> <div>12%</div> <div>92%</div> <div>5%</div> </div>
1	C	318	<div> <div>6%</div> <div>92%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	3	-	-	-	X
2	EDO	B	9	-	-	-	X
3	FUC	A	1	-	-	-	X
3	FUC	B	2	-	-	-	X
3	FUC	C	3	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14964 atoms, of which 7015 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	310	Total	C	H	N	O	S	0	0	0
			4709	1524	2322	402	452	9			
1	B	309	Total	C	H	N	O	S	0	0	0
			4698	1521	2317	401	450	9			
1	C	309	Total	C	H	N	O	S	0	0	0
			4697	1521	2316	401	450	9			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	221	GLY	-	EXPRESSION TAG	UNP Q5F4T5
A	222	PRO	-	EXPRESSION TAG	UNP Q5F4T5
A	223	GLY	-	EXPRESSION TAG	UNP Q5F4T5
A	224	SER	-	EXPRESSION TAG	UNP Q5F4T5
B	221	GLY	-	EXPRESSION TAG	UNP Q5F4T5
B	222	PRO	-	EXPRESSION TAG	UNP Q5F4T5
B	223	GLY	-	EXPRESSION TAG	UNP Q5F4T5
B	224	SER	-	EXPRESSION TAG	UNP Q5F4T5
C	221	GLY	-	EXPRESSION TAG	UNP Q5F4T5
C	222	PRO	-	EXPRESSION TAG	UNP Q5F4T5
C	223	GLY	-	EXPRESSION TAG	UNP Q5F4T5
C	224	SER	-	EXPRESSION TAG	UNP Q5F4T5

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	C	1	Total	C	H	O	0	0
			10	2	6	2		
2	C	1	Total	C	H	O	0	0
			10	2	6	2		
2	C	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 3 is SUGAR (ALPHA-L-FUCOSE) (three-letter code: FUC) (formula: C₆H₁₂O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		
3	C	1	Total	C	O	0	0
			11	6	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	268	Total	O	0	0
			268	268		
4	B	174	Total	O	0	0
			174	174		
4	C	285	Total	O	0	0
			285	285		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Capsid protein



• Molecule 1: Capsid protein



• Molecule 1: Capsid protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	80.53 Å 115.91 Å 267.80 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.67 – 1.85 25.67 – 1.85	Depositor EDS
% Data completeness (in resolution range)	93.7 (25.67-1.85) 93.1 (25.67-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 1.85 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.164 , 0.189 0.157 , 0.182	Depositor DCC
R_{free} test set	5039 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	23.9	Xtriage
Anisotropy	0.510	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 105628 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	14964	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.36	0/2456	0.51	0/3366
1	B	0.32	0/2450	0.50	0/3358
1	C	0.35	0/2450	0.52	0/3358
All	All	0.34	0/7356	0.51	0/10082

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2387	2322	2317	12	0
1	B	2381	2317	2312	11	0
1	C	2381	2316	2312	9	0
2	A	12	18	18	0	0
2	B	16	24	24	1	0
2	C	12	18	18	0	0
3	A	11	0	12	0	0
3	B	11	0	12	1	0
3	C	11	0	12	0	0
4	A	268	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	174	0	0	3	0
4	C	285	0	0	3	2
All	All	7949	7015	7037	32	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:315:PHE:O	1:C:317:PRO:HD3	2.10	0.51
1:B:347:PRO:HA	1:B:351:ASN:ND2	2.26	0.51
1:A:224:SER:HB2	4:A:719:HOH:O	2.10	0.51
1:C:538:MET:C	4:C:660:HOH:O	2.51	0.49
1:B:293:GLN:HB2	1:B:303:TRP:CZ3	2.47	0.49
1:B:510:ASN:HB3	4:B:655:HOH:O	2.13	0.48
1:B:477:LEU:HD12	1:B:528:TRP:CD1	2.48	0.48
1:A:471:GLN:HG3	2:B:5:EDO:H21	1.95	0.48
1:C:304:ASN:OD1	1:C:375:ASN:OD1	2.30	0.48
1:B:524:ARG:NH1	4:B:604:HOH:O	2.46	0.47
1:C:487:ASN:OD1	1:C:489:GLU:HB3	2.15	0.47
1:B:524:ARG:NH2	4:B:680:HOH:O	2.48	0.47
1:C:288:GLY:HA3	1:C:306:THR:O	2.14	0.47
1:A:487:ASN:OD1	1:A:489:GLU:HB3	2.15	0.46
1:B:323:ALA:HB1	1:B:324:PRO:HD2	1.98	0.46
1:C:492:ARG:HD3	4:C:637:HOH:O	2.15	0.46
1:A:484:ARG:NH2	1:A:524:ARG:HD2	2.32	0.45
1:A:494:LEU:HD11	1:A:518:PRO:HD2	1.99	0.44
1:A:484:ARG:HH22	1:A:524:ARG:HH11	1.65	0.43
1:A:294:VAL:HG21	1:A:377:GLN:NE2	2.32	0.43
1:A:259:VAL:HG13	1:A:411:TRP:CZ3	2.54	0.43
1:A:393:LYS:HE2	4:A:182:HOH:O	2.19	0.43
1:C:287:ARG:HG2	4:C:617:HOH:O	2.18	0.42
1:C:323:ALA:HB1	1:C:324:PRO:HD2	2.01	0.42
1:B:293:GLN:HG2	1:B:294:VAL:N	2.35	0.42
1:A:494:LEU:CD1	1:A:518:PRO:HD2	2.50	0.42
1:B:323:ALA:HB1	1:B:324:PRO:CD	2.50	0.41
1:B:484:ARG:HE	1:B:526:ASP:HB2	1.85	0.41
1:C:225:LYS:HA	1:C:226:PRO:HD3	1.91	0.41
1:A:452:TYR:CE1	3:B:2:FUC:H61	2.56	0.40
1:A:242:PHE:CD2	1:A:243:PRO:HD2	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:462:PRO:HG2	1:B:465:TRP:CD1	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:697:HOH:O	4:C:697:HOH:O[3_654]	1.84	0.36
4:C:658:HOH:O	4:C:662:HOH:O[3_654]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/318 (96%)	298 (97%)	7 (2%)	1 (0%)	46	29
1	B	305/318 (96%)	297 (97%)	8 (3%)	0	100	100
1	C	305/318 (96%)	294 (96%)	11 (4%)	0	100	100
All	All	916/954 (96%)	889 (97%)	26 (3%)	1 (0%)	56	39

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	529	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/271 (98%)	264 (100%)	1 (0%)	93	92
1	B	264/271 (97%)	263 (100%)	1 (0%)	93	92
1	C	264/271 (97%)	264 (100%)	0	100	100
All	All	793/813 (98%)	791 (100%)	2 (0%)	94	94

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	356	ARG
1	B	356	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	377	GLN
1	A	405	ASN
1	B	351	ASN
1	C	375	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	FUC	A	1	-	11,11,11	0.59	0	16,16,16	1.05	1 (6%)
2	EDO	A	10	-	3,3,3	0.48	0	2,2,2	0.37	0
2	EDO	A	3	-	3,3,3	0.49	0	2,2,2	0.44	0
2	EDO	A	6	-	3,3,3	0.52	0	2,2,2	0.37	0
3	FUC	B	2	-	11,11,11	0.60	0	16,16,16	1.05	1 (6%)
2	EDO	B	4	-	3,3,3	0.39	0	2,2,2	0.51	0
2	EDO	B	5	-	3,3,3	0.45	0	2,2,2	0.47	0
2	EDO	B	7	-	3,3,3	0.51	0	2,2,2	0.38	0
2	EDO	B	9	-	3,3,3	0.53	0	2,2,2	0.26	0
2	EDO	C	1	-	3,3,3	0.47	0	2,2,2	0.47	0
2	EDO	C	2	-	3,3,3	0.45	0	2,2,2	0.39	0
3	FUC	C	3	-	11,11,11	0.58	0	16,16,16	1.05	1 (6%)
2	EDO	C	8	-	3,3,3	0.54	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FUC	A	1	-	-	0/0/20/20	0/1/1/1
2	EDO	A	10	-	-	0/1/1/1	0/0/0/0
2	EDO	A	3	-	-	0/1/1/1	0/0/0/0
2	EDO	A	6	-	-	0/1/1/1	0/0/0/0
3	FUC	B	2	-	-	0/0/20/20	0/1/1/1
2	EDO	B	4	-	-	0/1/1/1	0/0/0/0
2	EDO	B	5	-	-	0/1/1/1	0/0/0/0
2	EDO	B	7	-	-	0/1/1/1	0/0/0/0
2	EDO	B	9	-	-	0/1/1/1	0/0/0/0
2	EDO	C	1	-	-	0/1/1/1	0/0/0/0
2	EDO	C	2	-	-	0/1/1/1	0/0/0/0
3	FUC	C	3	-	-	0/0/20/20	0/1/1/1
2	EDO	C	8	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	C	3	FUC	O2-C2-C1	2.24	114.74	109.82
3	B	2	FUC	O2-C2-C1	2.24	114.75	109.82
3	A	1	FUC	O2-C2-C1	2.24	114.76	109.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	2	FUC	1	0
2	B	5	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	310/318 (97%)	0.04	23 (7%) 17 17	16, 32, 71, 138	0
1	B	309/318 (97%)	0.39	37 (11%) 6 6	18, 42, 82, 139	0
1	C	309/318 (97%)	0.03	20 (6%) 22 21	14, 32, 67, 146	0
All	All	928/954 (97%)	0.15	80 (8%) 13 12	14, 35, 76, 146	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	346	VAL	7.9
1	B	532	PHE	6.7
1	C	347	PRO	6.6
1	A	224	SER	5.7
1	B	531	GLN	5.5
1	A	294	VAL	5.4
1	A	346	VAL	5.3
1	B	346	VAL	5.1
1	C	348	GLY	4.9
1	B	492	ARG	4.7
1	B	294	VAL	4.7
1	A	350	GLY	4.6
1	A	492	ARG	4.4
1	B	490	THR	4.4
1	A	347	PRO	4.3
1	C	345	THR	4.2
1	B	433	VAL	4.2
1	C	350	GLY	4.2
1	B	347	PRO	4.1
1	B	435	PRO	4.1
1	B	345	THR	4.1
1	B	434	PHE	4.0
1	B	383	THR	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	348	GLY	4.0
1	C	351	ASN	4.0
1	C	404	ALA	3.8
1	A	383	THR	3.8
1	A	490	THR	3.7
1	B	534	THR	3.7
1	B	225	LYS	3.6
1	A	348	GLY	3.5
1	A	345	THR	3.5
1	B	505	LEU	3.4
1	B	350	GLY	3.3
1	A	351	ASN	3.3
1	C	492	ARG	3.3
1	B	510	ASN	3.2
1	B	349	GLU	3.1
1	C	383	THR	3.0
1	A	384	GLN	3.0
1	A	404	ALA	3.0
1	B	384	GLN	3.0
1	C	405	ASN	3.0
1	B	405	ASN	3.0
1	B	499	LEU	2.9
1	C	300	GLY	2.9
1	C	499	LEU	2.9
1	C	489	GLU	2.8
1	C	435	PRO	2.8
1	A	499	LEU	2.8
1	B	404	ALA	2.7
1	B	477	LEU	2.7
1	C	349	GLU	2.6
1	C	294	VAL	2.6
1	C	384	GLN	2.6
1	A	489	GLU	2.6
1	B	256	SER	2.6
1	B	460	LEU	2.5
1	A	344	ASN	2.5
1	A	505	LEU	2.5
1	B	465	TRP	2.4
1	B	488	PRO	2.4
1	B	300	GLY	2.4
1	C	490	THR	2.4
1	C	505	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	292	GLN	2.3
1	A	300	GLY	2.3
1	B	489	GLU	2.3
1	A	481	ALA	2.3
1	B	351	ASN	2.2
1	B	474	ALA	2.1
1	A	349	GLU	2.1
1	B	491	GLY	2.1
1	B	292	GLN	2.1
1	B	344	ASN	2.1
1	B	440	LEU	2.1
1	A	352	LEU	2.0
1	B	318	THR	2.0
1	A	405	ASN	2.0
1	C	256	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	FUC	B	2	11/11	0.84	0.18	5.94	43,46,52,63	0
3	FUC	A	1	11/11	0.85	0.16	5.51	42,47,59,60	0
3	FUC	C	3	11/11	0.79	0.18	5.47	40,45,53,62	0
2	EDO	A	3	4/4	0.97	0.15	2.42	23,35,43,43	0
2	EDO	B	9	4/4	0.94	0.13	2.30	29,34,40,40	0
2	EDO	A	10	4/4	0.92	0.13	1.34	27,42,54,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	EDO	C	1	4/4	0.94	0.15	1.33	26,38,46,46	0
2	EDO	C	2	4/4	0.93	0.13	1.14	22,39,47,47	0
2	EDO	C	8	4/4	0.97	0.11	1.11	22,32,38,38	0
2	EDO	B	4	4/4	0.92	0.12	0.92	40,48,55,66	0
2	EDO	B	5	4/4	0.93	0.12	0.70	35,46,57,57	0
2	EDO	A	6	4/4	0.91	0.10	0.48	33,40,48,48	0
2	EDO	B	7	4/4	0.98	0.07	-0.81	25,31,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.