



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:20 AM GMT

PDB ID : 3OOQ
Title : CRYSTAL STRUCTURE OF amidohydrolase from *Thermotoga maritima* MSB8
Authors : Malashkevich, V.N.; Toro, R.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2010-08-31
Resolution : 2.06 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry. We welcome your comments at validation@mail.wwpdb.org. A user guide is available at <http://wwpdb.org/validation/2016/XrayValidationReportHelp> with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1799 (2.08-2.04)
Clashscore	102246	1910 (2.08-2.04)
Ramachandran outliers	100387	1893 (2.08-2.04)
Sidechain outliers	100360	1893 (2.08-2.04)
RSRZ outliers	91569	1802 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	 2% 86% 9% . .
1	B	396	 3% 88% 8% . .
1	C	396	 % 87% 8% . .
1	D	396	 3% 88% 7% . .
1	E	396	 2% 86% 10% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	396	 3% 87% 9% ..
1	G	396	 2% 85% 10% ..
1	H	396	 3% 85% 10% ..
1	I	396	 4% 86% 10% ..
1	J	396	 3% 86% 10% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	C	500	-	-	-	X
2	GOL	E	500	-	-	-	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 31584 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called amidohydrolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	384	2985	1914	502	556	2	11	0	1	0
1	B	383	2970	1905	501	551	2	11	0	0	0
1	C	384	2996	1921	503	559	2	11	0	3	0
1	D	384	2979	1910	502	554	2	11	0	0	0
1	E	383	2970	1905	501	551	2	11	0	0	0
1	F	385	2986	1915	503	555	2	11	0	0	0
1	G	383	2974	1908	501	552	2	11	0	1	0
1	H	383	2976	1909	502	552	2	11	0	1	0
1	I	383	2976	1909	501	553	2	11	0	1	0
1	J	383	2970	1905	501	551	2	11	0	0	0

There are 110 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP Q9X247
A	2	SER	-	expression tag	UNP Q9X247
A	3	LEU	-	expression tag	UNP Q9X247
A	389	GLU	-	expression tag	UNP Q9X247
A	390	GLY	-	expression tag	UNP Q9X247
A	391	HIS	-	expression tag	UNP Q9X247
A	392	HIS	-	expression tag	UNP Q9X247
A	393	HIS	-	expression tag	UNP Q9X247
A	394	HIS	-	expression tag	UNP Q9X247

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	395	HIS	-	expression tag	UNP Q9X247
A	396	HIS	-	expression tag	UNP Q9X247
B	1	MSE	-	expression tag	UNP Q9X247
B	2	SER	-	expression tag	UNP Q9X247
B	3	LEU	-	expression tag	UNP Q9X247
B	389	GLU	-	expression tag	UNP Q9X247
B	390	GLY	-	expression tag	UNP Q9X247
B	391	HIS	-	expression tag	UNP Q9X247
B	392	HIS	-	expression tag	UNP Q9X247
B	393	HIS	-	expression tag	UNP Q9X247
B	394	HIS	-	expression tag	UNP Q9X247
B	395	HIS	-	expression tag	UNP Q9X247
B	396	HIS	-	expression tag	UNP Q9X247
C	1	MSE	-	expression tag	UNP Q9X247
C	2	SER	-	expression tag	UNP Q9X247
C	3	LEU	-	expression tag	UNP Q9X247
C	389	GLU	-	expression tag	UNP Q9X247
C	390	GLY	-	expression tag	UNP Q9X247
C	391	HIS	-	expression tag	UNP Q9X247
C	392	HIS	-	expression tag	UNP Q9X247
C	393	HIS	-	expression tag	UNP Q9X247
C	394	HIS	-	expression tag	UNP Q9X247
C	395	HIS	-	expression tag	UNP Q9X247
C	396	HIS	-	expression tag	UNP Q9X247
D	1	MSE	-	expression tag	UNP Q9X247
D	2	SER	-	expression tag	UNP Q9X247
D	3	LEU	-	expression tag	UNP Q9X247
D	389	GLU	-	expression tag	UNP Q9X247
D	390	GLY	-	expression tag	UNP Q9X247
D	391	HIS	-	expression tag	UNP Q9X247
D	392	HIS	-	expression tag	UNP Q9X247
D	393	HIS	-	expression tag	UNP Q9X247
D	394	HIS	-	expression tag	UNP Q9X247
D	395	HIS	-	expression tag	UNP Q9X247
D	396	HIS	-	expression tag	UNP Q9X247
E	1	MSE	-	expression tag	UNP Q9X247
E	2	SER	-	expression tag	UNP Q9X247
E	3	LEU	-	expression tag	UNP Q9X247
E	389	GLU	-	expression tag	UNP Q9X247
E	390	GLY	-	expression tag	UNP Q9X247
E	391	HIS	-	expression tag	UNP Q9X247
E	392	HIS	-	expression tag	UNP Q9X247

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	393	HIS	-	expression tag	UNP Q9X247
E	394	HIS	-	expression tag	UNP Q9X247
E	395	HIS	-	expression tag	UNP Q9X247
E	396	HIS	-	expression tag	UNP Q9X247
F	1	MSE	-	expression tag	UNP Q9X247
F	2	SER	-	expression tag	UNP Q9X247
F	3	LEU	-	expression tag	UNP Q9X247
F	389	GLU	-	expression tag	UNP Q9X247
F	390	GLY	-	expression tag	UNP Q9X247
F	391	HIS	-	expression tag	UNP Q9X247
F	392	HIS	-	expression tag	UNP Q9X247
F	393	HIS	-	expression tag	UNP Q9X247
F	394	HIS	-	expression tag	UNP Q9X247
F	395	HIS	-	expression tag	UNP Q9X247
F	396	HIS	-	expression tag	UNP Q9X247
G	1	MSE	-	expression tag	UNP Q9X247
G	2	SER	-	expression tag	UNP Q9X247
G	3	LEU	-	expression tag	UNP Q9X247
G	389	GLU	-	expression tag	UNP Q9X247
G	390	GLY	-	expression tag	UNP Q9X247
G	391	HIS	-	expression tag	UNP Q9X247
G	392	HIS	-	expression tag	UNP Q9X247
G	393	HIS	-	expression tag	UNP Q9X247
G	394	HIS	-	expression tag	UNP Q9X247
G	395	HIS	-	expression tag	UNP Q9X247
G	396	HIS	-	expression tag	UNP Q9X247
H	1	MSE	-	expression tag	UNP Q9X247
H	2	SER	-	expression tag	UNP Q9X247
H	3	LEU	-	expression tag	UNP Q9X247
H	389	GLU	-	expression tag	UNP Q9X247
H	390	GLY	-	expression tag	UNP Q9X247
H	391	HIS	-	expression tag	UNP Q9X247
H	392	HIS	-	expression tag	UNP Q9X247
H	393	HIS	-	expression tag	UNP Q9X247
H	394	HIS	-	expression tag	UNP Q9X247
H	395	HIS	-	expression tag	UNP Q9X247
H	396	HIS	-	expression tag	UNP Q9X247
I	1	MSE	-	expression tag	UNP Q9X247
I	2	SER	-	expression tag	UNP Q9X247
I	3	LEU	-	expression tag	UNP Q9X247
I	389	GLU	-	expression tag	UNP Q9X247
I	390	GLY	-	expression tag	UNP Q9X247

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	391	HIS	-	expression tag	UNP Q9X247
I	392	HIS	-	expression tag	UNP Q9X247
I	393	HIS	-	expression tag	UNP Q9X247
I	394	HIS	-	expression tag	UNP Q9X247
I	395	HIS	-	expression tag	UNP Q9X247
I	396	HIS	-	expression tag	UNP Q9X247
J	1	MSE	-	expression tag	UNP Q9X247
J	2	SER	-	expression tag	UNP Q9X247
J	3	LEU	-	expression tag	UNP Q9X247
J	389	GLU	-	expression tag	UNP Q9X247
J	390	GLY	-	expression tag	UNP Q9X247
J	391	HIS	-	expression tag	UNP Q9X247
J	392	HIS	-	expression tag	UNP Q9X247
J	393	HIS	-	expression tag	UNP Q9X247
J	394	HIS	-	expression tag	UNP Q9X247
J	395	HIS	-	expression tag	UNP Q9X247
J	396	HIS	-	expression tag	UNP Q9X247

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	F	1	Total	C	O	0	0
			6	3	3		
2	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	229	Total	O	0	0
			229	229		
3	B	179	Total	O	0	0
			179	179		
3	C	213	Total	O	0	0
			213	213		
3	D	188	Total	O	0	0
			188	188		
3	E	167	Total	O	0	0
			167	167		
3	F	199	Total	O	0	0
			199	199		
3	G	170	Total	O	0	0
			170	170		
3	H	149	Total	O	0	0
			149	149		
3	I	136	Total	O	0	0
			136	136		
3	J	142	Total	O	0	0
			142	142		





● Molecule 1: amidohydrolase





● Molecule 1: amidohydrolase




● Molecule 1: amidohydrolase



● Molecule 1: amidohydrolase



● Molecule 1: amidohydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	152.45Å 148.30Å 195.02Å 90.00° 105.31° 90.00°	Depositor
Resolution (Å)	19.92 – 2.06 19.92 – 2.06	Depositor EDS
% Data completeness (in resolution range)	98.4 (19.92-2.06) 98.4 (19.92-2.06)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.06Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.167 , 0.213 0.173 , 0.218	Depositor DCC
R_{free} test set	12737 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	25.8	Xtrriage
Anisotropy	0.018	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.5	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	1 of 252990 reflections (0.000%)	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	31584	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/3034	0.68	0/4078
1	B	0.53	0/3016	0.64	0/4054
1	C	0.58	0/3051	0.67	1/4101 (0.0%)
1	D	0.56	0/3025	0.63	0/4066
1	E	0.56	0/3016	0.65	0/4054
1	F	0.57	0/3032	0.64	0/4076
1	G	0.54	0/3023	0.65	0/4064
1	H	0.53	0/3025	0.65	0/4066
1	I	0.53	0/3025	0.62	0/4066
1	J	0.52	0/3016	0.61	0/4054
All	All	0.55	0/30263	0.65	1/40679 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	207	ASP	CB-CG-OD1	5.11	122.90	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	200	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2985	0	3061	30	0
1	B	2970	0	3049	22	0
1	C	2996	0	3073	26	0
1	D	2979	0	3055	22	0
1	E	2970	0	3049	24	0
1	F	2986	0	3064	20	0
1	G	2974	0	3056	39	0
1	H	2976	0	3057	24	0
1	I	2976	0	3055	26	0
1	J	2970	0	3049	25	0
2	A	6	0	8	0	0
2	C	6	0	8	2	0
2	E	6	0	8	0	0
2	F	6	0	8	0	0
2	H	6	0	8	0	0
3	A	229	0	0	6	0
3	B	179	0	0	5	0
3	C	213	0	0	6	0
3	D	188	0	0	3	0
3	E	167	0	0	5	0
3	F	199	0	0	3	0
3	G	170	0	0	3	0
3	H	149	0	0	1	0
3	I	136	0	0	4	0
3	J	142	0	0	2	0
All	All	31584	0	30608	238	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 238 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:195:LEU:O	1:B:199:GLU:HG2	1.68	0.94
1:I:195:LEU:O	1:I:199:GLU:HG2	1.68	0.92
1:F:135:ARG:HD2	1:F:141:GLU:O	1.71	0.89
3:E:426:HOH:O	1:H:205:GLU:HG2	1.71	0.89
1:A:388:ARG:O	1:A:389:GLU:HB2	1.76	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/396 (97%)	373 (97%)	10 (3%)	0	100	100
1	B	381/396 (96%)	369 (97%)	12 (3%)	0	100	100
1	C	385/396 (97%)	372 (97%)	13 (3%)	0	100	100
1	D	382/396 (96%)	370 (97%)	12 (3%)	0	100	100
1	E	381/396 (96%)	367 (96%)	14 (4%)	0	100	100
1	F	383/396 (97%)	369 (96%)	14 (4%)	0	100	100
1	G	382/396 (96%)	372 (97%)	10 (3%)	0	100	100
1	H	382/396 (96%)	370 (97%)	12 (3%)	0	100	100
1	I	382/396 (96%)	369 (97%)	13 (3%)	0	100	100
1	J	381/396 (96%)	370 (97%)	11 (3%)	0	100	100
All	All	3822/3960 (96%)	3701 (97%)	121 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	323/321 (101%)	309 (96%)	14 (4%)	35	27
1	B	321/321 (100%)	311 (97%)	10 (3%)	47	41
1	C	325/321 (101%)	314 (97%)	11 (3%)	44	37
1	D	322/321 (100%)	306 (95%)	16 (5%)	30	20
1	E	321/321 (100%)	310 (97%)	11 (3%)	44	37
1	F	323/321 (101%)	310 (96%)	13 (4%)	38	30
1	G	322/321 (100%)	309 (96%)	13 (4%)	38	30
1	H	322/321 (100%)	305 (95%)	17 (5%)	28	18
1	I	322/321 (100%)	313 (97%)	9 (3%)	51	45
1	J	321/321 (100%)	307 (96%)	14 (4%)	35	26
All	All	3222/3210 (100%)	3094 (96%)	128 (4%)	38	30

5 of 128 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	105	GLU
1	F	247	LEU
1	J	39	ASN
1	E	164	GLU
1	F	47	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	30	ASN
1	G	63	HIS
1	I	226	HIS
1	E	39	ASN
1	F	226	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	500	-	5,5,5	0.41	0	5,5,5	0.34	0
2	GOL	C	500	-	5,5,5	0.39	0	5,5,5	0.59	0
2	GOL	E	500	-	5,5,5	0.33	0	5,5,5	0.65	0
2	GOL	F	500	-	5,5,5	0.33	0	5,5,5	0.48	0
2	GOL	H	500	-	5,5,5	0.30	0	5,5,5	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	500	-	-	0/4/4/4	0/0/0/0
2	GOL	C	500	-	-	0/4/4/4	0/0/0/0
2	GOL	E	500	-	-	0/4/4/4	0/0/0/0
2	GOL	F	500	-	-	0/4/4/4	0/0/0/0
2	GOL	H	500	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	500	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	373/396 (94%)	-0.42	7 (1%) 70 73	15, 21, 40, 54	0
1	B	372/396 (93%)	-0.23	12 (3%) 51 56	16, 25, 53, 73	0
1	C	373/396 (94%)	-0.41	5 (1%) 79 81	16, 23, 37, 53	0
1	D	373/396 (94%)	-0.29	10 (2%) 58 63	17, 24, 47, 68	0
1	E	372/396 (93%)	-0.37	7 (1%) 70 73	17, 26, 42, 65	0
1	F	374/396 (94%)	-0.31	12 (3%) 51 56	16, 24, 46, 67	0
1	G	372/396 (93%)	-0.25	8 (2%) 65 69	17, 26, 48, 69	0
1	H	372/396 (93%)	-0.25	13 (3%) 48 53	17, 28, 52, 71	0
1	I	372/396 (93%)	-0.13	16 (4%) 39 43	18, 29, 56, 81	0
1	J	372/396 (93%)	-0.16	11 (2%) 54 59	19, 30, 55, 78	0
All	All	3725/3960 (94%)	-0.28	101 (2%) 58 63	15, 26, 49, 81	0

The worst 5 of 101 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	200	GLY	7.6
1	D	200	GLY	6.5
1	G	75	TYR	6.3
1	H	44	ASP	5.9
1	I	43	PRO	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	E	500	6/6	0.79	0.19	3.13	38,41,42,42	0
2	GOL	C	500	6/6	0.78	0.24	2.54	50,52,56,56	0
2	GOL	H	500	6/6	0.85	0.15	1.89	54,54,56,56	0
2	GOL	A	500	6/6	0.87	0.12	1.15	36,37,37,38	0
2	GOL	F	500	6/6	0.94	0.12	1.12	27,28,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.