



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:19 AM GMT

PDB ID : 3OP5
Title : Human vaccinia-related kinase 1
Authors : Allerston, C.K.; Uttarkar, S.; Savitsky, P.; Elkins, J.M.; Filippakopoulos, P.; Krojer, T.; Rellos, P.; Fedorov, O.; Eswaran, J.; Brenner, B.; Keates, T.; Das, S.; King, O.; Chalk, R.; Berridge, G.; von Delft, F.; Gileadi, O.; Arrow-smith, C.H.; Edwards, A.M.; Weigelt, J.; Bountra, C.; Knapp, S.; Structural Genomics Consortium (SGC)
Deposited on : 2010-08-31
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	<div> <div></div> <div>83% 5% 12%</div> </div>
1	B	364	<div> <div></div> <div>82% 5% 12%</div> </div>
1	C	364	<div> <div>10%</div> <div>80% • 15%</div> </div>
1	D	364	<div> <div>6%</div> <div>81% • 17%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	366	-	-	-	X
3	EDO	C	366	-	-	-	X
3	EDO	C	367	-	-	-	X
4	GOL	A	367	-	-	-	X
4	GOL	B	366	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10797 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase VRK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	320	Total	C	N	O	S	0	0	0
			2516	1611	427	465	13			
1	B	319	Total	C	N	O	S	0	1	0
			2517	1612	428	464	13			
1	C	308	Total	C	N	O	S	0	0	0
			2399	1542	406	439	12			
1	D	302	Total	C	N	O	S	0	0	0
			2333	1501	397	423	12			

There are 52 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	EXPRESSION TAG	UNP Q99986
A	2	MET	-	EXPRESSION TAG	UNP Q99986
A	34	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
A	35	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
A	36	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
A	212	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
A	214	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
A	215	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
A	292	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
A	293	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
A	295	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
A	359	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
A	360	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
B	1	SER	-	EXPRESSION TAG	UNP Q99986
B	2	MET	-	EXPRESSION TAG	UNP Q99986
B	34	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
B	35	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
B	36	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
B	212	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
B	214	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
B	215	ALA	GLU	ENGINEERED MUTATION	UNP Q99986

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	292	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
B	293	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
B	295	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
B	359	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
B	360	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
C	1	SER	-	EXPRESSION TAG	UNP Q99986
C	2	MET	-	EXPRESSION TAG	UNP Q99986
C	34	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
C	35	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
C	36	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
C	212	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
C	214	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
C	215	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
C	292	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
C	293	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
C	295	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
C	359	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
C	360	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
D	1	SER	-	EXPRESSION TAG	UNP Q99986
D	2	MET	-	EXPRESSION TAG	UNP Q99986
D	34	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
D	35	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
D	36	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
D	212	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
D	214	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
D	215	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
D	292	ALA	GLU	ENGINEERED MUTATION	UNP Q99986
D	293	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
D	295	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
D	359	ALA	LYS	ENGINEERED MUTATION	UNP Q99986
D	360	ALA	LYS	ENGINEERED MUTATION	UNP Q99986

- Molecule 2 is [4-({4-[(5-CYCLOPROPYL-1H-PYRAZOL-3-YL)AMINO]PYRIMIDIN-2-YL}AMINO)PHENYL]ACETONITRILE (three-letter code: REB) (formula: C₁₈H₁₇N₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			25	18	7		
2	B	1	Total	C	N	0	0
			25	18	7		
2	C	1	Total	C	N	0	0
			25	18	7		
2	D	1	Total	C	N	0	0
			25	18	7		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	230	Total O 230 230	0	0
5	B	243	Total O 243 243	0	0
5	C	222	Total O 222 222	0	0

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	205	Total 205	O 205	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase VRK1

Chain A: 





- Molecule 1: Serine/threonine-protein kinase VRK1

Chain B: 





- Molecule 1: Serine/threonine-protein kinase VRK1

Chain C: 



- Molecule 1: Serine/threonine-protein kinase VRK1

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.97Å 97.19Å 191.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.34 – 2.40 46.34 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.5 (46.34-2.40) 98.6 (46.34-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.39Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.178 , 0.213 0.189 , 0.226	Depositor DCC
R_{free} test set	2007 reflections (3.05%)	DCC
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.505	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 53.0	EDS
Estimated twinning fraction	0.019 for k,h,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 67764 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10797	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: REB, GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/2575	0.62	0/3487
1	B	0.44	0/2576	0.61	0/3491
1	C	0.43	0/2453	0.60	0/3322
1	D	0.42	0/2386	0.60	0/3234
All	All	0.43	0/9990	0.61	0/13534

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	133	ARG	Sidechain
1	B	44	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2516	0	2439	10	0
1	B	2517	0	2429	10	0
1	C	2399	0	2279	7	0
1	D	2333	0	2220	4	0
2	A	25	0	16	0	0
2	B	25	0	16	0	0
2	C	25	0	16	3	0
2	D	25	0	16	1	0
3	A	4	0	6	2	0
3	B	4	0	6	0	0
3	C	12	0	18	1	0
4	A	6	0	8	2	0
4	B	6	0	8	1	0
5	A	230	0	0	1	0
5	B	243	0	0	0	0
5	C	222	0	0	0	0
5	D	205	0	0	0	0
All	All	10797	0	9477	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 33 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:230:ILE:HB	3:C:366:EDO:H22	1.56	0.87
2:C:365:REB:H16A	2:C:365:REB:H22	1.69	0.72
1:D:43:ILE:HD12	2:D:365:REB:H15A	1.77	0.65
1:A:73:GLU:HB2	1:A:80:LEU:HD22	1.85	0.59
1:A:89:ARG:HH22	4:A:367:GOL:H2	1.67	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/364 (87%)	308 (98%)	8 (2%)	0	100	100
1	B	316/364 (87%)	308 (98%)	8 (2%)	0	100	100
1	C	298/364 (82%)	290 (97%)	8 (3%)	0	100	100
1	D	292/364 (80%)	283 (97%)	9 (3%)	0	100	100
All	All	1222/1456 (84%)	1189 (97%)	33 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/304 (84%)	252 (98%)	4 (2%)	70	86
1	B	255/304 (84%)	252 (99%)	3 (1%)	78	90
1	C	234/304 (77%)	230 (98%)	4 (2%)	68	85
1	D	226/304 (74%)	223 (99%)	3 (1%)	76	89
All	All	971/1216 (80%)	957 (99%)	14 (1%)	74	88

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	318	GLU
1	C	133	ARG
1	D	103	ARG
1	B	160	ARG
1	C	318	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	REB	A	365	-	27,28,28	2.43	8 (29%)	33,38,38	4.41	20 (60%)
3	EDO	A	366	-	3,3,3	0.51	0	2,2,2	0.36	0
4	GOL	A	367	-	5,5,5	0.45	0	5,5,5	0.45	0
2	REB	B	365	-	27,28,28	2.42	9 (33%)	33,38,38	4.00	18 (54%)
4	GOL	B	366	-	5,5,5	0.35	0	5,5,5	0.41	0
3	EDO	B	367	-	3,3,3	0.41	0	2,2,2	0.55	0
2	REB	C	365	-	27,28,28	2.16	8 (29%)	33,38,38	4.04	18 (54%)
3	EDO	C	366	-	3,3,3	0.48	0	2,2,2	0.65	0
3	EDO	C	367	-	3,3,3	0.45	0	2,2,2	0.33	0
3	EDO	C	368	-	3,3,3	0.47	0	2,2,2	0.39	0
2	REB	D	365	-	27,28,28	2.25	8 (29%)	33,38,38	4.85	19 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	REB	A	365	-	-	0/12/17/17	0/3/4/4
3	EDO	A	366	-	-	0/1/1/1	0/0/0/0
4	GOL	A	367	-	-	0/4/4/4	0/0/0/0
2	REB	B	365	-	-	0/12/17/17	0/3/4/4
4	GOL	B	366	-	-	0/4/4/4	0/0/0/0
3	EDO	B	367	-	-	0/1/1/1	0/0/0/0
2	REB	C	365	-	-	0/12/17/17	0/3/4/4
3	EDO	C	366	-	-	0/1/1/1	0/0/0/0
3	EDO	C	367	-	-	0/1/1/1	0/0/0/0
3	EDO	C	368	-	-	0/1/1/1	0/0/0/0
2	REB	D	365	-	-	0/12/17/17	0/3/4/4

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	365	REB	C09-N13	-4.32	1.28	1.34
2	B	365	REB	C09-N13	-4.09	1.28	1.34
2	B	365	REB	C11-N12	-4.00	1.28	1.33
2	A	365	REB	C11-N12	-3.96	1.28	1.33
2	C	365	REB	C11-N12	-3.50	1.29	1.33

The worst 5 of 75 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	365	REB	N04-C05-N06	-9.66	116.38	126.67
2	B	365	REB	N04-C05-N06	-8.48	117.64	126.67
2	D	365	REB	N04-C05-N06	-8.18	117.95	126.67
2	C	365	REB	N04-C05-N06	-7.71	118.46	126.67
2	C	365	REB	C24-C25-N26	-6.65	169.84	178.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	366	EDO	2	0
4	A	367	GOL	2	0
4	B	366	GOL	1	0
2	C	365	REB	3	0
3	C	366	EDO	1	0
2	D	365	REB	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/364 (87%)	-0.39	1 (0%) 94 94	22, 38, 63, 84	0
1	B	319/364 (87%)	-0.38	3 (0%) 85 85	24, 43, 69, 82	0
1	C	308/364 (84%)	0.04	38 (12%) 5 5	23, 42, 96, 121	0
1	D	302/364 (82%)	-0.10	22 (7%) 18 18	25, 46, 103, 131	0
All	All	1249/1456 (85%)	-0.21	64 (5%) 32 32	22, 42, 88, 131	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	36	ALA	5.4
1	C	126	TYR	5.1
1	C	34	ALA	4.9
1	C	292	ALA	4.7
1	D	29	ILE	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	A	366	4/4	0.95	0.26	8.29	39,43,43,50	0
4	GOL	B	366	6/6	0.88	0.21	6.16	85,85,86,87	0
3	EDO	C	367	4/4	0.91	0.20	4.80	52,54,55,61	0
4	GOL	A	367	6/6	0.82	0.20	4.71	72,73,74,74	0
3	EDO	C	366	4/4	0.96	0.18	3.55	30,32,36,42	0
2	REB	D	365	25/25	0.90	0.20	1.57	58,60,64,67	25
3	EDO	B	367	4/4	0.95	0.17	1.01	45,46,46,56	0
2	REB	B	365	25/25	0.94	0.16	-0.04	25,30,33,35	25
2	REB	C	365	25/25	0.95	0.15	-0.30	35,41,45,48	25
2	REB	A	365	25/25	0.97	0.12	-0.51	23,25,32,37	25
3	EDO	C	368	4/4	0.89	0.36	-	56,56,56,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.