



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:56 AM GMT

PDB ID : 2OTE
Title : Crystal structure of a monomeric cyan fluorescent protein in the photobleached state
Authors : Henderson, J.N.; Ai, H.; Campbell, R.E.; Remington, S.J.
Deposited on : 2007-02-07
Resolution : 1.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3129 (1.50-1.46)
Clashscore	102246	3380 (1.50-1.46)
Ramachandran outliers	100387	3310 (1.50-1.46)
Sidechain outliers	100360	3308 (1.50-1.46)
RSRZ outliers	91569	3133 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	<div> <div>2%</div> <div>86%</div> <div>12%</div> <div>.</div> </div>
1	B	214	<div> <div>%</div> <div>85%</div> <div>14%</div> <div>.</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4007 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GFP-like fluorescent chromoprotein cFP484.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	19	0
			1779	1164	282	328	5			
1	B	214	Total	C	N	O	S	0	14	0
			1774	1149	285	335	5			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ASN	HIS	ENGINEERED MUTATION	UNP Q9U6Y3
A	44	ILE	LEU	ENGINEERED MUTATION	UNP Q9U6Y3
A	62	THR	SER	ENGINEERED MUTATION	UNP Q9U6Y3
A	66	PIA	GLN	CHROMOPHORE	UNP Q9U6Y3
A	66	PIA	TYR	CHROMOPHORE	UNP Q9U6Y3
A	66	PIA	GLY	CHROMOPHORE	UNP Q9U6Y3
A	72	PHE	LEU	ENGINEERED MUTATION	UNP Q9U6Y3
A	80	PRO	ALA	ENGINEERED MUTATION	UNP Q9U6Y3
A	81	ASN	ASP	ENGINEERED MUTATION	UNP Q9U6Y3
A	123	HIS	ARG	ENGINEERED MUTATION	UNP Q9U6Y3
A	124	LEU	PHE	ENGINEERED MUTATION	UNP Q9U6Y3
A	125	LYS	ASP	ENGINEERED MUTATION	UNP Q9U6Y3
A	127	GLU	MET	ENGINEERED MUTATION	UNP Q9U6Y3
A	150	LEU	MET	ENGINEERED MUTATION	UNP Q9U6Y3
A	162	LYS	SER	ENGINEERED MUTATION	UNP Q9U6Y3
A	164	LYS	SER	ENGINEERED MUTATION	UNP Q9U6Y3
A	173	HIS	TYR	ENGINEERED MUTATION	UNP Q9U6Y3
A	175	VAL	CYS	ENGINEERED MUTATION	UNP Q9U6Y3
A	179	THR	SER	ENGINEERED MUTATION	UNP Q9U6Y3
A	182	ARG	LYS	ENGINEERED MUTATION	UNP Q9U6Y3
A	186	ALA	VAL	ENGINEERED MUTATION	UNP Q9U6Y3
A	213	VAL	LEU	ENGINEERED MUTATION	UNP Q9U6Y3
A	216	SER	ASN	ENGINEERED MUTATION	UNP Q9U6Y3
B	42	ASN	HIS	ENGINEERED MUTATION	UNP Q9U6Y3
B	44	ILE	LEU	ENGINEERED MUTATION	UNP Q9U6Y3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	62	THR	SER	ENGINEERED MUTATION	UNP Q9U6Y3
B	66	PIA	GLN	CHROMOPHORE	UNP Q9U6Y3
B	66	PIA	TYR	CHROMOPHORE	UNP Q9U6Y3
B	66	PIA	GLY	CHROMOPHORE	UNP Q9U6Y3
B	72	PHE	LEU	ENGINEERED MUTATION	UNP Q9U6Y3
B	80	PRO	ALA	ENGINEERED MUTATION	UNP Q9U6Y3
B	81	ASN	ASP	ENGINEERED MUTATION	UNP Q9U6Y3
B	123	HIS	ARG	ENGINEERED MUTATION	UNP Q9U6Y3
B	124	LEU	PHE	ENGINEERED MUTATION	UNP Q9U6Y3
B	125	LYS	ASP	ENGINEERED MUTATION	UNP Q9U6Y3
B	127	GLU	MET	ENGINEERED MUTATION	UNP Q9U6Y3
B	150	LEU	MET	ENGINEERED MUTATION	UNP Q9U6Y3
B	162	LYS	SER	ENGINEERED MUTATION	UNP Q9U6Y3
B	164	LYS	SER	ENGINEERED MUTATION	UNP Q9U6Y3
B	173	HIS	TYR	ENGINEERED MUTATION	UNP Q9U6Y3
B	175	VAL	CYS	ENGINEERED MUTATION	UNP Q9U6Y3
B	179	THR	SER	ENGINEERED MUTATION	UNP Q9U6Y3
B	182	ARG	LYS	ENGINEERED MUTATION	UNP Q9U6Y3
B	186	ALA	VAL	ENGINEERED MUTATION	UNP Q9U6Y3
B	213	VAL	LEU	ENGINEERED MUTATION	UNP Q9U6Y3
B	216	SER	ASN	ENGINEERED MUTATION	UNP Q9U6Y3

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	243	Total 243	O 243	0	13
3	B	207	Total 207	O 207	0	6


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GFP-like fluorescent chromoprotein cFP484



- Molecule 1: GFP-like fluorescent chromoprotein cFP484



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.49Å 66.31Å 97.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.47 31.03 – 1.47	Depositor EDS
% Data completeness (in resolution range)	94.7 (10.00-1.47) 85.7 (31.03-1.47)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 1.47Å)	Xtriage
Refinement program	SHELXL	Depositor
R, R_{free}	0.190 , 0.261 0.167 , 0.172	Depositor DCC
R_{free} test set	3426 reflections (5.83%)	DCC
Wilson B-factor (Å ²)	14.8	Xtriage
Anisotropy	0.274	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 75.3	EDS
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Outliers	0 of 68646 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4007	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.46 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.6744e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PIA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1847	1.23	5/2504 (0.2%)
1	B	0.51	0/1824	1.29	10/2468 (0.4%)
All	All	0.50	0/3671	1.26	15/4972 (0.3%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	182	ARG	CD-NE-CZ	19.47	150.86	123.60
1	B	182	ARG	NE-CZ-NH1	12.65	126.62	120.30
1	A	220	ARG	NE-CZ-NH2	-10.60	115.00	120.30
1	A	173	HIS	CA-CB-CG	7.67	126.63	113.60
1	B	70[A]	ARG	NE-CZ-NH2	7.07	123.83	120.30
1	B	101	ASP	CB-CG-OD1	7.02	124.62	118.30
1	A	118	PHE	CB-CG-CD2	-6.96	115.93	120.80
1	B	220	ARG	NE-CZ-NH1	6.55	123.57	120.30
1	B	153	ARG	CD-NE-CZ	6.32	132.45	123.60
1	B	153	ARG	NE-CZ-NH2	5.76	123.18	120.30
1	A	220	ARG	CD-NE-CZ	5.71	131.60	123.60
1	B	101	ASP	CB-CG-OD2	-5.68	113.19	118.30
1	B	160	ASP	CB-CG-OD2	5.68	123.41	118.30
1	B	182	ARG	NE-CZ-NH2	-5.63	117.48	120.30
1	A	93	TRP	CA-CB-CG	5.59	124.32	113.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1779	0	1704	23	0
1	B	1774	0	1688	16	0
2	A	4	0	3	1	0
3	A	243	0	0	2	0
3	B	207	0	0	6	0
All	All	4007	0	3395	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66[A]:PIA:HE2	1:B:181:TYR:OH	1.89	0.71
1:B:209:ASN:HB3	3:B:368:HOH:O	1.91	0.70
1:A:85:GLN:HE22	1:A:188:LYS:H	1.36	0.70
1:A:96:THR:HG23	1:A:104[A]:ILE:HD11	1.78	0.65
1:A:33:GLY:HA2	2:A:1:ACT:H1	1.83	0.61
1:B:79:ILE:HG12	1:B:221:TYR:CZ	2.36	0.60
1:A:104[A]:ILE:HD12	1:A:105:VAL:N	2.19	0.58
1:B:23:ASN:HD21	1:B:129:PHE:H	1.51	0.57
1:A:146[B]:SER:HB3	1:A:163:HIS:CE1	2.40	0.57
1:B:63[B]:ASN:H	1:B:63[B]:ASN:ND2	2.04	0.56
1:A:66[A]:PIA:HE1	1:A:181:TYR:OH	2.07	0.54
1:A:146[A]:SER:HB2	3:A:327:HOH:O	2.07	0.54
1:A:104[A]:ILE:HD12	1:A:105:VAL:H	1.73	0.53
1:A:63[B]:ASN:ND2	1:A:63[B]:ASN:H	2.04	0.52
1:B:146[B]:SER:HB3	1:B:163:HIS:CE1	2.45	0.52
1:B:11:ASP:OD1	1:B:34:GLU:OE2	2.29	0.51
1:A:105:VAL:HG12	1:A:107[B]:VAL:HG23	1.93	0.49
1:B:53:PRO:HD2	3:B:427:HOH:O	2.12	0.49
1:A:182[A]:ARG:HD2	3:A:299:HOH:O	2.13	0.48
1:B:149:ILE:O	1:B:159:GLY:HA2	2.13	0.48
1:A:101:ASP:O	1:A:102:LYS:HB2	2.14	0.48
1:B:218[B]:VAL:HG13	3:B:362:HOH:O	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:GLN:NE2	1:A:188:LYS:H	2.08	0.47
1:A:96:THR:HG23	1:A:104[A]:ILE:CD1	2.44	0.47
1:A:107[B]:VAL:HG22	1:A:124:LEU:HG	1.96	0.46
1:B:198:ARG:NE	3:B:384:HOH:O	2.49	0.46
1:B:141:LEU:HD21	1:B:168:GLU:HG2	1.97	0.46
1:B:6:GLY:N	3:B:390:HOH:O	2.50	0.45
1:B:66[A]:PIA:HE2	1:B:181:TYR:HH	1.82	0.44
1:A:141[B]:LEU:HD21	1:A:168:GLU:HG2	1.99	0.44
1:B:42:ASN:ND2	3:B:296:HOH:O	2.51	0.43
1:B:79:ILE:HG12	1:B:221:TYR:CE2	2.52	0.43
1:A:141[B]:LEU:HD12	1:A:166:LEU:HG	2.00	0.43
1:A:118:PHE:C	1:A:119[A]:ILE:HD13	2.41	0.41
1:A:141[A]:LEU:HD12	1:A:141[A]:LEU:HA	1.90	0.41
1:A:11[A]:ASP:OD2	1:A:34[A]:GLU:OE2	2.38	0.41
1:A:162:LYS:HG2	1:A:176:ASP:OD1	2.21	0.41
1:A:144:GLU:O	1:A:163:HIS:HE1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/214 (105%)	219 (98%)	5 (2%)	0	100	100
1	B	218/214 (102%)	212 (97%)	6 (3%)	0	100	100
All	All	442/428 (103%)	431 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/188 (98%)	175 (95%)	9 (5%)	31	4
1	B	187/188 (100%)	179 (96%)	8 (4%)	35	6
All	All	371/376 (99%)	354 (95%)	17 (5%)	41	5

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104[A]	ILE
1	A	104[B]	ILE
1	A	107[A]	VAL
1	A	107[B]	VAL
1	A	141[A]	LEU
1	A	141[B]	LEU
1	A	162	LYS
1	A	163	HIS
1	A	197[A]	HIS
1	B	23	ASN
1	B	32[A]	GLU
1	B	32[B]	GLU
1	B	104	ILE
1	B	107	VAL
1	B	163	HIS
1	B	200	GLU
1	B	215[A]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN
1	A	85	GLN
1	B	23	ASN
1	B	42	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PIA	A	66[A]	1	21,21,22	2.20	8 (38%)	25,29,31	2.92	9 (36%)
1	PIA	B	66[A]	1	21,21,22	2.36	8 (38%)	25,29,31	4.27	9 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PIA	A	66[A]	1	-	0/8/27/28	0/2/2/2
1	PIA	B	66[A]	1	-	0/8/27/28	0/2/2/2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66[A]	PIA	OH-CZ	-5.18	1.24	1.37
1	B	66[A]	PIA	CG2-CB2	-4.72	1.37	1.46
1	A	66[A]	PIA	OH-CZ	-4.68	1.25	1.37
1	A	66[A]	PIA	CG2-CB2	-4.48	1.37	1.46
1	A	66[A]	PIA	CE2-CD2	-2.70	1.33	1.38
1	A	66[A]	PIA	CA1-C1	-2.57	1.47	1.51
1	B	66[A]	PIA	CE2-CD2	-2.45	1.34	1.38
1	B	66[A]	PIA	CA1-C1	-2.33	1.47	1.51
1	B	66[A]	PIA	CD2-CG2	2.18	1.43	1.39
1	A	66[A]	PIA	CD2-CG2	2.41	1.44	1.39
1	B	66[A]	PIA	CE2-CZ	2.98	1.44	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66[A]	PIA	CE2-CZ	2.98	1.44	1.38
1	A	66[A]	PIA	CE1-CZ	3.27	1.45	1.38
1	B	66[A]	PIA	CE1-CZ	3.34	1.45	1.38
1	A	66[A]	PIA	C1-N3	3.45	1.43	1.37
1	B	66[A]	PIA	C1-N3	3.49	1.43	1.37

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66[A]	PIA	O2-C2-N3	-5.24	113.20	124.50
1	A	66[A]	PIA	CA3-N3-C1	-3.70	123.05	127.36
1	B	66[A]	PIA	CA3-N3-C1	-2.88	124.01	127.36
1	A	66[A]	PIA	O2-C2-N3	-2.80	118.45	124.50
1	B	66[A]	PIA	CE1-CD1-CG2	-2.68	117.94	121.29
1	A	66[A]	PIA	CE2-CZ-CE1	-2.66	116.00	119.79
1	B	66[A]	PIA	CB2-CA2-N2	-2.36	124.44	128.67
1	B	66[A]	PIA	CD1-CE1-CZ	2.15	122.35	119.87
1	A	66[A]	PIA	OH-CZ-CE1	2.26	126.44	120.05
1	A	66[A]	PIA	CG2-CB2-CA2	2.27	133.16	130.22
1	A	66[A]	PIA	CA2-C2-N3	2.35	104.58	103.40
1	B	66[A]	PIA	CB2-CA2-C2	2.83	126.50	122.36
1	A	66[A]	PIA	CD2-CE2-CZ	2.92	123.24	119.87
1	B	66[A]	PIA	CA3-N3-C2	3.74	130.09	123.99
1	A	66[A]	PIA	CA3-N3-C2	3.88	130.30	123.99
1	B	66[A]	PIA	CA2-C2-N3	5.04	105.92	103.40
1	A	66[A]	PIA	O2-C2-CA2	11.15	136.97	130.95
1	B	66[A]	PIA	O2-C2-CA2	18.39	140.88	130.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66[A]	PIA	1	0
1	B	66[A]	PIA	2	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	1	-	1,3,3	2.58	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	A	1	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	ACT	CH3-C	2.58	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/214 (99%)	-0.29	4 (1%) 70 72	13, 18, 27, 35	10 (4%)
1	B	213/214 (99%)	-0.27	3 (1%) 78 80	14, 18, 26, 31	10 (4%)
All	All	426/428 (99%)	-0.28	7 (1%) 74 78	13, 18, 26, 35	20 (4%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	221	TYR	2.8
1	B	199	ILE	2.8
1	B	213	VAL	2.4
1	B	201	ILE	2.0
1	A	213	VAL	2.0
1	A	6	GLY	2.0
1	A	104[A]	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	PIA	B	66[A]	20/21	0.95	0.13	-	13,18,30,38	8
1	PIA	A	66[A]	20/21	0.96	0.11	-	13,18,30,38	8

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ACT	A	1	4/4	0.90	0.10	1.33	29,34,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.