



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:25 PM GMT

PDB ID : 4OVI  
Title : Phenylacetone monooxygenase: oxidised enzyme in complex with APADP  
Authors : Martinoli, C.; Fraaije, M.W.; Mattevi, A.  
Deposited on : 2013-09-19  
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6965 (1.90-1.86)
Clashscore	102246	7778 (1.90-1.86)
Ramachandran outliers	100387	7691 (1.90-1.86)
Sidechain outliers	100360	7692 (1.90-1.86)
RSRZ outliers	91569	6979 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylacetone monooxygenase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	531	Total	C	N	O	S	0	2	0
			4234	2701	728	795	10			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is 3-ACETILPYRIDINE ADENINE DINUCLEOTIDE PHOSPHATE (three-letter code: N01) (formula:  $C_{22}H_{26}N_6O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	22	6	17	3		

- Molecule 4 is HEXAETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula:  $Cl$ ).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	214	Total	O	0	0
			214	214		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phenylacetone monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.24Å 107.24Å 107.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 1.87 37.90 – 1.87	Depositor EDS
% Data completeness (in resolution range)	100.0 (40.00-1.87) 100.0 (37.90-1.87)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.59 (at 1.87Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.221 , 0.277 0.220 , 0.276	Depositor DCC
$R_{free}$ test set	3032 reflections (5.40%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.0	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.2	EDS
Estimated twinning fraction	0.031 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 59179 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4570	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PGE, CL, ACT, N01, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.96	0/4353	0.99	6/5924 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	525	ARG	NE-CZ-NH1	8.48	124.54	120.30
1	A	89	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	A	66	ASP	CB-CG-OD1	6.19	123.87	118.30
1	A	166	ASP	CB-CG-OD2	-5.91	112.99	118.30
1	A	390	ASP	CB-CG-OD1	5.47	123.22	118.30
1	A	297	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4234	0	4083	57	0
2	A	53	0	31	4	0
3	A	48	0	23	3	0
4	A	10	0	14	3	1
5	A	1	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	4	0	3	1	0
7	A	6	0	8	0	0
8	A	214	0	0	12	0
All	All	4570	0	4162	61	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:313:ILE:O	1:A:316:THR:HG22	1.63	0.97
1:A:316:THR:HG23	1:A:317:VAL:HG23	1.62	0.82
1:A:297:ARG:NH2	8:A:1029:HOH:O	2.15	0.79
1:A:279:ARG:HB3	4:A:603:PGE:H42	1.64	0.77
1:A:517:TYR:CE2	1:A:524:TYR:HB2	2.20	0.76
1:A:506:ASN:HB2	8:A:1005:HOH:O	1.85	0.75
1:A:402:ARG:HD2	1:A:406:ASN:ND2	2.08	0.69
1:A:402:ARG:HD2	1:A:406:ASN:HD21	1.58	0.67
1:A:484:GLU:O	1:A:488:GLU:HB2	1.95	0.67
1:A:346:GLU:CB	8:A:987:HOH:O	2.44	0.66
1:A:335:THR:OG1	1:A:498:THR:HG21	1.94	0.66
1:A:12:GLN:HB3	8:A:1103:HOH:O	1.95	0.65
1:A:265:ALA:HB3	1:A:301:ALA:HB1	1.81	0.62
1:A:260:GLN:HG2	8:A:1002:HOH:O	2.01	0.60
1:A:17:VAL:O	1:A:143:ALA:HA	2.04	0.58
1:A:471:ARG:O	1:A:541:LEU:HA	2.06	0.54
1:A:161:PHE:HB2	1:A:162:PRO:HD2	1.88	0.54
2:A:601:FAD:C10	5:A:604:CL:CL	2.92	0.54
1:A:466:LYS:HA	8:A:1092:HOH:O	2.08	0.54
1:A:535:GLY:O	1:A:536:TYR:HB2	2.09	0.53
1:A:510:LYS:HG3	1:A:511:PRO:HD2	1.92	0.52
1:A:506:ASN:OD1	8:A:1020:HOH:O	2.19	0.50
1:A:279:ARG:HB3	4:A:603:PGE:H52	1.94	0.50
1:A:234:PHE:O	1:A:237:ASP:HB2	2.13	0.49
1:A:351:ASP:HB2	6:A:605:ACT:CH3	2.43	0.48
1:A:202:SER:HB2	1:A:203:PRO:HD3	1.94	0.48
1:A:224:PRO:HD3	1:A:281:TRP:CH2	2.48	0.48
1:A:231:ASP:OD1	1:A:231:ASP:C	2.51	0.48
1:A:198:GLY:HA2	1:A:201:VAL:HG12	1.95	0.48
1:A:177:TRP:CG	1:A:178:PRO:HD2	2.49	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:423:LEU:O	1:A:432:PHE:HA	2.14	0.47
1:A:316:THR:CG2	1:A:345:TYR:HB3	2.45	0.47
1:A:152:GLN:HE22	2:A:601:FAD:H5'2	1.79	0.47
1:A:316:THR:HG21	1:A:345:TYR:HB3	1.97	0.47
1:A:129:ASN:OD1	1:A:471:ARG:HD2	2.14	0.47
1:A:196:SER:HB3	3:A:602:N01:C4N	2.44	0.47
1:A:126:GLU:HB2	8:A:1113:HOH:O	2.14	0.47
2:A:601:FAD:N5	3:A:602:N01:CAA	2.78	0.46
1:A:466:LYS:N	8:A:1092:HOH:O	2.47	0.46
1:A:526:GLN:O	1:A:530:GLU:HG3	2.15	0.46
1:A:421:LEU:HA	1:A:482:TRP:CE2	2.51	0.46
1:A:466:LYS:CA	8:A:1092:HOH:O	2.64	0.46
1:A:325:ARG:HD2	1:A:357:ASP:HB2	1.99	0.45
1:A:399:ILE:HG22	1:A:401:ILE:HG13	1.98	0.45
1:A:332:PRO:HD2	1:A:498:THR:CG2	2.48	0.44
4:A:603:PGE:H5	8:A:911:HOH:O	2.16	0.44
1:A:23:GLY:O	1:A:28:GLY:HA3	2.17	0.44
1:A:294:ASP:N	1:A:294:ASP:OD1	2.51	0.43
1:A:389:PHE:O	1:A:506:ASN:ND2	2.50	0.43
1:A:412:LYS:NZ	1:A:479:GLU:OE1	2.49	0.43
1:A:164:LEU:HD13	1:A:366:ILE:HG13	2.01	0.43
1:A:166:ASP:O	1:A:368:PRO:HA	2.18	0.43
1:A:44:VAL:HB	1:A:113:ILE:HG12	2.01	0.42
1:A:144:ARG:HG2	1:A:145:TYR:CE2	2.54	0.42
1:A:424:SER:HB3	8:A:1022:HOH:O	2.20	0.42
2:A:601:FAD:HM73	3:A:602:N01:C5N	2.50	0.41
1:A:177:TRP:HA	1:A:178:PRO:HD3	1.97	0.41
1:A:199:ILE:HG12	1:A:344:TYR:CZ	2.56	0.41
1:A:319:ASP:C	1:A:319:ASP:OD1	2.59	0.41
1:A:341:GLU:HB2	1:A:344:TYR:HB3	2.04	0.40
1:A:65[B]:CYS:HB3	1:A:96:ILE:HD11	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:603:PGE:O1	4:A:603:PGE:O1[5_555]	1.71	0.49

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	531/542 (98%)	510 (96%)	19 (4%)	2 (0%)	39	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	400	ASP
1	A	536	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	443/453 (98%)	425 (96%)	18 (4%)	37	23

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	ASP
1	A	161	PHE
1	A	179	HIS
1	A	181	PRO
1	A	196	SER
1	A	237	ASP
1	A	242	TYR
1	A	273	LEU
1	A	351	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	360	SER
1	A	369	ARG
1	A	447	LEU
1	A	453	HIS
1	A	476	LEU
1	A	488	GLU
1	A	512	ARG
1	A	522	HIS
1	A	526	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	ASN
1	A	152	GLN
1	A	315	ASN
1	A	406	ASN
1	A	526	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	601	-	48,58,58	1.68	9 (18%)	54,89,89	2.48	10 (18%)
3	N01	A	602	-	42,52,52	2.32	11 (26%)	55,80,80	2.18	12 (21%)
4	PGE	A	603	-	9,9,9	0.25	0	8,8,8	1.72	2 (25%)
6	ACT	A	605	-	1,3,3	1.92	0	0,3,3	0.00	-
7	GOL	A	606	-	5,5,5	0.37	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	0/30/50/50	0/6/6/6
3	N01	A	602	-	-	0/27/67/67	0/5/5/5
4	PGE	A	603	-	-	0/7/7/7	0/0/0/0
6	ACT	A	605	-	-	0/0/0/0	0/0/0/0
7	GOL	A	606	-	-	0/4/4/4	0/0/0/0

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	N01	CAA-C7N	-7.77	1.25	1.49
2	A	601	FAD	O2B-C2B	-2.17	1.37	1.43
3	A	602	N01	C4A-N3A	-2.04	1.32	1.35
2	A	601	FAD	C5'-C4'	2.07	1.54	1.51
3	A	602	N01	C5N-C4N	2.16	1.43	1.38
3	A	602	N01	C3B-C2B	2.23	1.58	1.53
3	A	602	N01	C2A-N3A	2.23	1.36	1.32
3	A	602	N01	C6A-N6A	2.35	1.42	1.34
2	A	601	FAD	C5X-N5	2.63	1.39	1.35
2	A	601	FAD	C10-N1	2.95	1.40	1.35
3	A	602	N01	O4B-C1B	3.22	1.45	1.41
2	A	601	FAD	C2A-N3A	3.24	1.37	1.32
2	A	601	FAD	C4-N3	3.38	1.39	1.33
2	A	601	FAD	C4X-N5	3.43	1.38	1.33
2	A	601	FAD	C2A-N1A	3.53	1.40	1.33
3	A	602	N01	P2B-O2B	3.65	1.68	1.62
2	A	601	FAD	C9A-N10	3.91	1.44	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	N01	O7N-C7N	4.31	1.37	1.22
3	A	602	N01	P2B-O2X	5.28	1.61	1.51
3	A	602	N01	P2B-O3X	6.67	1.64	1.51

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	N3A-C2A-N1A	-11.76	119.89	128.89
3	A	602	N01	N3A-C2A-N1A	-8.61	122.30	128.89
2	A	601	FAD	C4X-C10-N10	-5.46	117.30	120.52
2	A	601	FAD	C4-C4X-C10	-5.31	116.54	119.94
3	A	602	N01	C3B-C2B-C1B	-4.81	93.43	102.73
4	A	603	PGE	C5-O3-C4	-3.33	99.00	113.31
3	A	602	N01	O2X-P2B-O3X	-3.28	105.14	112.76
2	A	601	FAD	O2'-C2'-C1'	-3.13	102.25	109.94
3	A	602	N01	O5B-C5B-C4B	-3.11	97.64	109.12
2	A	601	FAD	C7M-C7-C6	-3.02	112.08	120.28
2	A	601	FAD	O3B-C3B-C4B	-2.83	102.57	111.05
3	A	602	N01	O2X-P2B-O2B	-2.39	98.21	105.83
3	A	602	N01	O1A-PA-O3	-2.38	94.30	105.09
3	A	602	N01	C4B-O4B-C1B	-2.34	107.15	109.72
4	A	603	PGE	O4-C6-C5	-2.19	98.53	112.03
3	A	602	N01	O3B-C3B-C2B	-2.07	105.19	111.16
2	A	601	FAD	C1B-N9A-C4A	-2.04	123.87	126.94
2	A	601	FAD	C8M-C8-C7	2.61	126.45	120.73
2	A	601	FAD	C7M-C7-C8	3.01	127.34	120.73
3	A	602	N01	O3-PA-O5B	3.57	112.40	102.94
3	A	602	N01	O4D-C1D-N1N	3.91	112.42	108.13
3	A	602	N01	N6A-C6A-N1A	4.39	128.62	119.20
3	A	602	N01	O2X-P2B-O1X	6.37	127.00	112.85
2	A	601	FAD	C4-N3-C2	6.40	120.78	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	FAD	4	0
3	A	602	N01	3	0
4	A	603	PGE	3	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	605	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	531/542 (97%)	0.04	10 (1%) 70 72	24, 41, 61, 73	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	468	GLY	3.2
1	A	142	ARG	2.6
1	A	467	ASN	2.4
1	A	442	ALA	2.3
1	A	463	TYR	2.3
1	A	376	ARG	2.2
1	A	542	THR	2.2
1	A	320	PRO	2.2
1	A	469	LEU	2.1
1	A	448	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PGE	A	603	10/10	0.89	0.12	1.91	37,48,59,59	0
3	N01	A	602	48/48	0.97	0.14	0.64	23,32,46,49	0
2	FAD	A	601	53/53	0.97	0.12	0.29	26,32,39,41	0
5	CL	A	604	1/1	0.99	0.03	-	41,41,41,41	0
7	GOL	A	606	6/6	0.88	0.14	-	59,62,67,70	0
6	ACT	A	605	4/4	0.97	0.11	-	50,53,56,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.