



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 07:28 PM GMT

PDB ID : 4OYY
Title : Humicola insolens cutinase
Authors : Dauter, Z.D.; Brzozowski, A.M.; Turkenburg, J.P.; Wilson, K.S.
Deposited on : 2014-02-13
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	F	194	%	87%	11%	..
1	G	194		90%	8%	..
1	H	194		88%	9%	..
1	I	194	%	90%	8%	..
1	J	194		88%	8%	..
1	K	194		88%	9%	..
1	L	194	%	91%	7%	..

2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 16975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cutinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	B	191	Total	C	N	O	S	0	0	0
			1397	872	254	266	5			
1	C	193	Total	C	N	O	S	0	0	0
			1419	885	260	269	5			
1	D	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	E	191	Total	C	N	O	S	0	0	0
			1396	871	254	266	5			
1	F	193	Total	C	N	O	S	0	0	0
			1419	885	260	269	5			
1	G	193	Total	C	N	O	S	0	0	0
			1416	884	260	267	5			
1	H	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	I	193	Total	C	N	O	S	0	0	0
			1410	881	256	268	5			
1	J	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	K	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	L	192	Total	C	N	O	S	0	0	0
			1410	880	258	267	5			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	10	Total	O	0	0
			10	10		
2	B	4	Total	O	0	0
			4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	7	Total O 7 7	0	0
2	D	8	Total O 8 8	0	0
2	E	3	Total O 3 3	0	0
2	F	8	Total O 8 8	0	0
2	G	15	Total O 15 15	0	0
2	H	8	Total O 8 8	0	0
2	I	8	Total O 8 8	0	0
2	J	10	Total O 10 10	0	0
2	K	9	Total O 9 9	0	0
2	L	8	Total O 8 8	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cutinase

Chain A:  91% 7% ..



- Molecule 1: Cutinase

Chain B:  88% 10% ..



- Molecule 1: Cutinase

Chain C:  87% 11% ..



- Molecule 1: Cutinase

Chain D:  91% 7% ..




- Molecule 1: Cutinase

Chain E:  87% 9% ...



- Molecule 1: Cutinase

Chain F:  87% 11% ..





- Molecule 1: Cutinase

Chain G: 90% 8% ..



- Molecule 1: Cutinase

Chain H: 88% 9% ..



- Molecule 1: Cutinase

Chain I: 90% 8% ..



- Molecule 1: Cutinase

Chain J: 88% 8% ..



- Molecule 1: Cutinase

Chain K: 88% 9% ..



- Molecule 1: Cutinase

Chain L: 91% 7% ..



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	125.55 Å 127.00 Å 134.50 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.76 – 3.00 29.76 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.76-3.00) 99.5 (29.76-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	4.76 (at 3.00 Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R , R_{free}	0.174 , 0.196 0.181 , 0.201	Depositor DCC
R_{free} test set	1338 reflections (3.18%)	DCC
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.332	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 35.5	EDS
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	10 of 43457 reflections (0.023%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16975	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.72 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.2586e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/1426	0.77	2/1936 (0.1%)
1	B	0.60	0/1421	0.79	3/1930 (0.2%)
1	C	0.59	0/1443	0.81	3/1959 (0.2%)
1	D	0.59	0/1426	0.74	1/1936 (0.1%)
1	E	0.58	0/1420	0.80	2/1929 (0.1%)
1	F	0.58	0/1443	0.81	3/1959 (0.2%)
1	G	0.61	0/1440	0.81	3/1955 (0.2%)
1	H	0.59	0/1426	0.82	7/1936 (0.4%)
1	I	0.56	0/1434	0.75	1/1948 (0.1%)
1	J	0.59	0/1426	0.79	3/1936 (0.2%)
1	K	0.59	0/1426	0.76	3/1936 (0.2%)
1	L	0.62	0/1434	0.81	1/1947 (0.1%)
All	All	0.59	0/17165	0.79	32/23307 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

There are no bond length outliers.

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	34	MET	CG-SD-CE	-10.76	82.98	100.20
1	E	4	ALA	N-CA-CB	9.26	123.07	110.10
1	J	51	ARG	NE-CZ-NH1	7.87	124.23	120.30
1	A	51	ARG	NE-CZ-NH1	7.81	124.21	120.30
1	G	192	ILE	CA-CB-CG1	7.56	125.37	111.00
1	G	51	ARG	NE-CZ-NH1	7.55	124.08	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	186	ARG	NE-CZ-NH2	-7.37	116.61	120.30
1	H	167	LEU	CB-CG-CD1	6.70	122.39	111.00
1	H	186	ARG	CD-NE-CZ	6.44	132.62	123.60
1	C	51	ARG	NE-CZ-NH1	6.43	123.52	120.30
1	F	51	ARG	NE-CZ-NH1	6.31	123.46	120.30
1	C	141	ARG	NE-CZ-NH1	6.21	123.41	120.30
1	H	193	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	J	73	ARG	N-CA-CB	-6.10	99.61	110.60
1	A	181	ARG	NE-CZ-NH1	6.01	123.30	120.30
1	H	181	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	I	51	ARG	NE-CZ-NH1	5.90	123.25	120.30
1	L	10	GLU	CG-CD-OE1	-5.69	106.92	118.30
1	H	167	LEU	CB-CG-CD2	-5.55	101.57	111.00
1	G	181	ARG	NE-CZ-NH1	5.55	123.07	120.30
1	K	193	ARG	NE-CZ-NH1	5.51	123.06	120.30
1	H	186	ARG	NE-CZ-NH1	5.51	123.05	120.30
1	K	181	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	E	181	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	J	181	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	B	51	ARG	NE-CZ-NH1	5.30	122.95	120.30
1	F	47	GLU	N-CA-CB	5.29	120.13	110.60
1	F	141	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	B	193	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	D	186	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	C	191	ARG	NE-CZ-NH2	-5.07	117.76	120.30
1	K	181	ARG	CB-CG-CD	5.02	124.65	111.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	1	GLN	Peptide
1	C	3	GLY	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1402	0	1394	5	0
1	B	1397	0	1385	6	0
1	C	1419	0	1416	8	0
1	D	1402	0	1394	6	0
1	E	1396	0	1383	13	0
1	F	1419	0	1416	10	0
1	G	1416	0	1414	10	0
1	H	1402	0	1394	6	0
1	I	1410	0	1401	8	0
1	J	1402	0	1394	9	0
1	K	1402	0	1394	10	0
1	L	1410	0	1405	7	0
2	A	10	0	0	0	0
2	B	4	0	0	1	0
2	C	7	0	0	0	0
2	D	8	0	0	0	0
2	E	3	0	0	0	0
2	F	8	0	0	0	0
2	G	15	0	0	0	0
2	H	8	0	0	1	0
2	I	8	0	0	0	0
2	J	10	0	0	0	0
2	K	9	0	0	1	0
2	L	8	0	0	1	0
All	All	16975	0	16790	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:68:THR:HG23	1:E:106:GLN:OE1	1.63	0.98
1:C:94:CYS:HB3	1:C:97:THR:HG23	1.48	0.96
1:G:1:GLN:OE1	1:G:1:GLN:N	2.03	0.90
1:E:68:THR:CG2	1:E:106:GLN:OE1	2.22	0.88
1:L:72:PRO:HG2	2:L:203:HOH:O	1.75	0.86
1:K:181:ARG:HG2	1:K:181:ARG:HH11	1.49	0.77
1:F:45:GLY:HA2	1:F:181:ARG:NH1	2.01	0.76
1:L:45:GLY:HA2	1:L:181:ARG:NH1	2.01	0.76
1:B:181:ARG:HD3	2:B:202:HOH:O	1.88	0.73
1:F:123:LYS:HE2	1:F:146:ASN:O	1.95	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:17:CYS:HB3	1:C:97:THR:HG21	1.77	0.67
1:C:94:CYS:HB3	1:C:97:THR:CG2	2.24	0.67
1:I:123:LYS:HE2	1:I:146:ASN:O	1.96	0.66
1:D:123:LYS:HE2	1:D:146:ASN:O	1.96	0.66
1:H:123:LYS:HE2	1:H:146:ASN:O	1.96	0.66
1:L:123:LYS:HE2	1:L:146:ASN:O	1.96	0.66
1:K:123:LYS:HE2	1:K:146:ASN:O	1.96	0.66
1:H:193:ARG:NH1	2:H:205:HOH:O	2.25	0.65
1:C:123:LYS:HE2	1:C:146:ASN:O	1.97	0.65
1:G:123:LYS:HE2	1:G:146:ASN:O	1.96	0.65
1:J:123:LYS:HE2	1:J:146:ASN:O	1.96	0.65
1:E:123:LYS:HE2	1:E:146:ASN:O	1.97	0.65
1:B:123:LYS:HE2	1:B:146:ASN:O	1.97	0.64
1:A:123:LYS:HE2	1:A:146:ASN:O	1.97	0.64
1:C:138:LEU:HD11	1:D:5:ILE:HG21	1.81	0.62
1:E:51:ARG:NH2	1:I:10:GLU:OE1	2.35	0.60
1:F:138:LEU:HD12	1:G:5:ILE:HG21	1.83	0.60
1:E:13:SER:OG	1:E:15:ASN:OD1	2.20	0.59
1:D:13:SER:OG	1:D:15:ASN:OD1	2.20	0.58
1:E:119:SER:O	1:E:122:VAL:HG23	2.02	0.58
1:F:13:SER:OG	1:F:15:ASN:OD1	2.21	0.58
1:J:13:SER:OG	1:J:15:ASN:OD1	2.20	0.58
1:K:193:ARG:NH1	2:K:204:HOH:O	2.35	0.58
1:H:3:GLY:HA3	1:H:40:PRO:HG3	1.86	0.57
1:K:3:GLY:HA3	1:K:40:PRO:HG3	1.88	0.56
1:B:3:GLY:HA3	1:B:40:PRO:HG3	1.87	0.56
1:L:45:GLY:HA2	1:L:181:ARG:HH11	1.70	0.56
1:J:3:GLY:HA3	1:J:40:PRO:HG3	1.89	0.54
1:F:45:GLY:HA2	1:F:181:ARG:HH11	1.70	0.54
1:E:51:ARG:HH22	1:I:10:GLU:HB3	1.73	0.54
1:D:3:GLY:HA3	1:D:40:PRO:HG3	1.89	0.54
1:A:3:GLY:HA3	1:A:40:PRO:HG3	1.89	0.54
1:G:100:VAL:HG13	1:G:192:ILE:HD11	1.90	0.53
1:B:162:VAL:HA	1:B:166:THR:O	2.09	0.52
1:G:100:VAL:HG13	1:G:192:ILE:CD1	2.41	0.51
1:G:162:VAL:HA	1:G:166:THR:O	2.11	0.51
1:I:162:VAL:HA	1:I:166:THR:O	2.11	0.51
1:D:162:VAL:HA	1:D:166:THR:O	2.12	0.50
1:F:162:VAL:HA	1:F:166:THR:O	2.12	0.50
1:K:162:VAL:HA	1:K:166:THR:O	2.10	0.50
1:C:162:VAL:HA	1:C:166:THR:O	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:162:VAL:HA	1:H:166:THR:O	2.11	0.50
1:A:162:VAL:HA	1:A:166:THR:O	2.13	0.49
1:J:162:VAL:HA	1:J:166:THR:O	2.12	0.49
1:E:162:VAL:HA	1:E:166:THR:O	2.11	0.49
1:L:162:VAL:HA	1:L:166:THR:O	2.12	0.49
1:F:43:ALA:HB1	1:F:55:ILE:HD13	1.94	0.49
1:G:43:ALA:HB1	1:G:55:ILE:HD13	1.96	0.48
1:B:43:ALA:HB1	1:B:55:ILE:HD13	1.95	0.48
1:C:43:ALA:HB1	1:C:55:ILE:HD13	1.95	0.47
1:I:43:ALA:HB1	1:I:55:ILE:HD13	1.95	0.47
1:D:43:ALA:HB1	1:D:55:ILE:HD13	1.96	0.47
1:H:43:ALA:HB1	1:H:55:ILE:HD13	1.96	0.47
1:J:43:ALA:HB1	1:J:55:ILE:HD13	1.96	0.47
1:E:43:ALA:HB1	1:E:55:ILE:HD13	1.97	0.47
1:F:141:ARG:NH1	1:G:6:GLU:OE2	2.48	0.47
1:K:43:ALA:HB1	1:K:55:ILE:HD13	1.97	0.46
1:J:28:SER:O	1:J:29:THR:HB	2.15	0.46
1:K:181:ARG:NH1	1:K:181:ARG:HG2	2.26	0.46
1:A:43:ALA:HB1	1:A:55:ILE:HD13	1.98	0.46
1:E:51:ARG:HA	1:E:51:ARG:HD2	1.87	0.44
1:G:100:VAL:CG1	1:G:192:ILE:HD11	2.47	0.44
1:L:191:ARG:HA	1:L:191:ARG:HD3	1.70	0.44
1:J:29:THR:HG23	1:J:29:THR:O	2.18	0.44
1:G:48:SER:HB2	1:H:96:ASN:HB2	1.99	0.43
1:E:3:GLY:HA3	1:E:4:ALA:CB	2.48	0.43
1:K:51:ARG:HA	1:K:51:ARG:HD3	1.81	0.43
1:E:3:GLY:HA3	1:E:4:ALA:HB2	2.01	0.42
1:K:181:ARG:HH11	1:K:181:ARG:CG	2.27	0.42
1:I:40:PRO:O	1:I:44:ASN:HB2	2.19	0.42
1:L:2:LEU:O	1:L:2:LEU:HG	2.20	0.42
1:J:29:THR:CG2	1:J:29:THR:O	2.68	0.41
1:E:65:ALA:O	1:E:68:THR:HG22	2.20	0.41
1:I:51:ARG:HG3	1:I:52:ASN:N	2.36	0.41
1:A:40:PRO:O	1:A:44:ASN:HB2	2.21	0.41
1:C:40:PRO:O	1:C:44:ASN:HB2	2.21	0.41
1:I:51:ARG:HG3	1:I:52:ASN:H	1.85	0.41
1:B:72:PRO:O	1:B:73:ARG:HB2	2.21	0.41
1:K:72:PRO:O	1:K:73:ARG:HB2	2.20	0.41
1:F:72:PRO:O	1:F:73:ARG:HB2	2.21	0.41
1:J:18:PRO:HG3	1:J:54:TRP:CG	2.57	0.40
1:F:40:PRO:O	1:F:44:ASN:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	189/194 (97%)	181 (96%)	7 (4%)	1 (0%)	34 76
1	B	189/194 (97%)	180 (95%)	8 (4%)	1 (0%)	34 76
1	C	191/194 (98%)	181 (95%)	9 (5%)	1 (0%)	34 76
1	D	189/194 (97%)	181 (96%)	7 (4%)	1 (0%)	34 76
1	E	189/194 (97%)	180 (95%)	7 (4%)	2 (1%)	17 58
1	F	191/194 (98%)	182 (95%)	8 (4%)	1 (0%)	34 76
1	G	191/194 (98%)	183 (96%)	7 (4%)	1 (0%)	34 76
1	H	189/194 (97%)	181 (96%)	7 (4%)	1 (0%)	34 76
1	I	191/194 (98%)	182 (95%)	8 (4%)	1 (0%)	34 76
1	J	189/194 (97%)	181 (96%)	7 (4%)	1 (0%)	34 76
1	K	189/194 (97%)	180 (95%)	8 (4%)	1 (0%)	34 76
1	L	190/194 (98%)	183 (96%)	6 (3%)	1 (0%)	34 76
All	All	2277/2328 (98%)	2175 (96%)	89 (4%)	13 (1%)	30 72

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	4	ALA
1	E	120	GLY
1	F	120	GLY
1	I	120	GLY
1	J	120	GLY
1	L	120	GLY
1	B	120	GLY
1	D	120	GLY
1	G	120	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	120	GLY
1	K	120	GLY
1	A	120	GLY
1	C	120	GLY

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	141/143 (99%)	136 (96%)	5 (4%)	43 80
1	B	140/143 (98%)	133 (95%)	7 (5%)	30 70
1	C	143/143 (100%)	134 (94%)	9 (6%)	22 60
1	D	141/143 (99%)	137 (97%)	4 (3%)	51 84
1	E	140/143 (98%)	133 (95%)	7 (5%)	30 70
1	F	143/143 (100%)	136 (95%)	7 (5%)	31 71
1	G	142/143 (99%)	136 (96%)	6 (4%)	36 76
1	H	141/143 (99%)	133 (94%)	8 (6%)	25 64
1	I	141/143 (99%)	134 (95%)	7 (5%)	30 70
1	J	141/143 (99%)	133 (94%)	8 (6%)	25 64
1	K	141/143 (99%)	134 (95%)	7 (5%)	30 70
1	L	142/143 (99%)	137 (96%)	5 (4%)	43 80
All	All	1696/1716 (99%)	1616 (95%)	80 (5%)	32 72

All (80) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	SER
1	A	28	SER
1	A	51	ARG
1	A	123	LYS
1	A	156	CYS
1	B	13	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	28	SER
1	B	119	SER
1	B	123	LYS
1	B	156	CYS
1	B	158	VAL
1	B	167	LEU
1	C	2	LEU
1	C	13	SER
1	C	28	SER
1	C	51	ARG
1	C	97	THR
1	C	123	LYS
1	C	156	CYS
1	C	179	GLU
1	C	193	ARG
1	D	13	SER
1	D	28	SER
1	D	123	LYS
1	D	156	CYS
1	E	13	SER
1	E	28	SER
1	E	122	VAL
1	E	123	LYS
1	E	126	VAL
1	E	156	CYS
1	E	179	GLU
1	F	2	LEU
1	F	13	SER
1	F	28	SER
1	F	123	LYS
1	F	149	ARG
1	F	156	CYS
1	F	193	ARG
1	G	13	SER
1	G	28	SER
1	G	51	ARG
1	G	123	LYS
1	G	156	CYS
1	G	179	GLU
1	H	13	SER
1	H	28	SER
1	H	119	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	123	LYS
1	H	141	ARG
1	H	156	CYS
1	H	158	VAL
1	H	167	LEU
1	I	13	SER
1	I	28	SER
1	I	123	LYS
1	I	129	VAL
1	I	156	CYS
1	I	158	VAL
1	I	193	ARG
1	J	13	SER
1	J	28	SER
1	J	29	THR
1	J	51	ARG
1	J	123	LYS
1	J	141	ARG
1	J	156	CYS
1	J	179	GLU
1	K	13	SER
1	K	28	SER
1	K	123	LYS
1	K	129	VAL
1	K	141	ARG
1	K	156	CYS
1	K	158	VAL
1	L	13	SER
1	L	28	SER
1	L	123	LYS
1	L	156	CYS
1	L	179	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	44	ASN
1	L	52	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	191/194 (98%)	-0.69	0	100	100	0
1	B	191/194 (98%)	-0.52	0	100	100	0
1	C	193/194 (99%)	-0.61	0	100	100	0
1	D	191/194 (98%)	-0.61	0	100	100	0
1	E	191/194 (98%)	-0.42	0	100	100	0
1	F	193/194 (99%)	-0.61	1 (0%)	91	76	0
1	G	193/194 (99%)	-0.72	0	100	100	0
1	H	191/194 (98%)	-0.65	0	100	100	0
1	I	193/194 (99%)	-0.55	1 (0%)	91	76	0
1	J	191/194 (98%)	-0.61	0	100	100	0
1	K	191/194 (98%)	-0.56	0	100	100	0
1	L	192/194 (98%)	-0.52	1 (0%)	91	76	0
All	All	2301/2328 (98%)	-0.59	3 (0%)	95	90	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	1	GLN	2.7
1	L	15	ASN	2.2
1	F	15	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.