



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:32 AM GMT

PDB ID : 3P4D
Title : Alternatingly modified 2'Fluoro RNA octamer f/rC4G4
Authors : Pallan, P.S.; Greene, E.M.; Jicman, P.A.; Pandey, R.K.; Manoharan, M.;
Rozners, E.; Egli, M.
Deposited on : 2010-10-06
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)
RNA backbone	2183	1033 (2.70-1.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	8	<div> <div style="width: 75%; background-color: green;"></div> <div style="width: 25%; background-color: yellow;"></div> </div> <div>75% 25%</div>
1	B	8	<div> <div style="width: 75%; background-color: green;"></div> <div style="width: 25%; background-color: yellow;"></div> </div> <div>75% 25%</div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 366 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*(CFZ)P*CP*(CFZ)P*CP*(GF2)P*GP*(GF2)P*G)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	8	Total	C	F	N	O	P	0	0	0
			169	76	4	32	50	7			
1	B	8	Total	C	F	N	O	P	0	0	0
			169	76	4	32	50	7			


- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	14	Total	O	0	0
			14	14		
2	B	14	Total	O	0	0
			14	14		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-R(*(CFZ)P*CP*(CFZ)P*CP*(GF2)P*GP*(GF2)P*G)-3'

Chain A:  75% 25%



- Molecule 1: 5'-R(*(CFZ)P*CP*(CFZ)P*CP*(GF2)P*GP*(GF2)P*G)-3'

Chain B:  75% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	41.63 Å 41.63 Å 126.49 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	21.08 – 1.85 21.08 – 1.85	Depositor EDS
% Data completeness (in resolution range)	100.0 (21.08-1.85) 99.1 (21.08-1.85)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.99 (at 1.85 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.210 , 0.271 0.218 , 0.279	Depositor DCC
R_{free} test set	279 reflections (7.88%)	DCC
Wilson B-factor (Å ²)	30.1	Xtriage
Anisotropy	0.334	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 42.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 3824 reflections (0.026%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	366	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GF2, CFZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/92	1.50	2/134 (1.5%)
1	B	1.01	0/92	1.38	0/134
All	All	0.92	0/184	1.44	2/268 (0.7%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	106	G	O4'-C1'-N9	5.74	112.80	108.20
1	A	102	C	O4'-C4'-C3'	-5.03	98.97	104.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	169	0	85	0	0
1	B	169	0	85	5	0
2	A	14	0	0	0	0
2	B	14	0	0	0	0
All	All	366	0	170	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 10.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:205:GF2:H5'	1:B:205:GF2:H8	1.24	1.14
1:B:205:GF2:C5'	1:B:205:GF2:H8	1.97	0.94
1:B:205:GF2:H5'	1:B:205:GF2:C8	2.14	0.69
1:B:205:GF2:C8	1:B:205:GF2:C5'	2.74	0.66
1:B:204:C:H2'	1:B:205:GF2:H5'	1.98	0.45

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	3/8 (37%)	0	0
1	B	3/8 (37%)	0	0
All	All	6/16 (37%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CFZ	A	101	1	14,18,22	1.00	2 (14%)	18,26,33	1.20	2 (11%)
1	CFZ	A	103	1	13,21,22	0.89	0	18,30,33	1.20	3 (16%)
1	GF2	A	105	1	17,25,26	1.55	2 (11%)	20,37,40	2.85	6 (30%)
1	GF2	A	107	1	17,25,26	1.61	2 (11%)	20,37,40	2.78	4 (20%)
1	CFZ	B	201	1	14,18,22	0.81	0	18,26,33	1.13	2 (11%)
1	CFZ	B	203	1	13,21,22	0.80	1 (7%)	18,30,33	1.01	1 (5%)
1	GF2	B	205	1	17,25,26	1.45	2 (11%)	20,37,40	3.19	9 (45%)
1	GF2	B	207	1	17,25,26	1.47	3 (17%)	20,37,40	2.36	6 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CFZ	A	101	1	-	0/2/22/26	0/2/2/2
1	CFZ	A	103	1	-	0/3/25/26	0/2/2/2
1	GF2	A	105	1	-	0/3/25/26	0/3/3/3
1	GF2	A	107	1	-	0/3/25/26	0/3/3/3
1	CFZ	B	201	1	-	0/2/22/26	0/2/2/2
1	CFZ	B	203	1	-	0/3/25/26	0/2/2/2
1	GF2	B	205	1	-	0/3/25/26	0/3/3/3
1	GF2	B	207	1	-	0/3/25/26	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	101	CFZ	C6-C5	-2.15	1.33	1.38
1	B	203	CFZ	C6-C5	-2.10	1.33	1.38
1	A	101	CFZ	C4-N3	-2.03	1.31	1.35
1	B	207	GF2	C6-N1	2.02	1.36	1.33
1	B	205	GF2	C6-C5	2.51	1.46	1.41
1	B	207	GF2	C6-C5	3.17	1.47	1.41
1	A	105	GF2	C6-C5	3.46	1.48	1.41
1	B	207	GF2	O6-C6	3.62	1.33	1.24
1	A	107	GF2	C6-C5	4.00	1.49	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	205	GF2	O6-C6	4.21	1.34	1.24
1	A	105	GF2	O6-C6	4.56	1.35	1.24
1	A	107	GF2	O6-C6	4.65	1.35	1.24

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	GF2	C5-C6-N1	-8.72	111.67	123.59
1	B	205	GF2	C5-C6-N1	-8.39	112.12	123.59
1	A	105	GF2	C5-C6-N1	-8.01	112.63	123.59
1	B	207	GF2	C5-C6-N1	-6.23	115.07	123.59
1	B	205	GF2	C5'-C4'-C3'	-3.91	99.70	115.21
1	B	207	GF2	C6-C5-C4	-3.83	116.31	120.90
1	B	205	GF2	O4'-C4'-C5'	-3.54	96.65	109.32
1	B	205	GF2	C6-C5-C4	-3.00	117.31	120.90
1	A	105	GF2	C6-C5-C4	-2.97	117.35	120.90
1	A	107	GF2	C6-C5-C4	-2.91	117.42	120.90
1	A	105	GF2	N3-C2-N1	-2.73	123.28	127.44
1	A	101	CFZ	C4'-O4'-C1'	-2.66	106.80	109.72
1	A	107	GF2	O4'-C4'-C3'	-2.56	99.99	105.15
1	B	205	GF2	O5'-C5'-C4'	-2.45	100.14	109.12
1	B	207	GF2	N3-C2-N1	-2.33	123.89	127.44
1	B	205	GF2	N3-C2-N1	-2.32	123.91	127.44
1	A	103	CFZ	C5-C4-N4	-2.29	117.79	121.31
1	B	207	GF2	C2'-C1'-N9	-2.04	110.12	113.76
1	A	103	CFZ	F2'-C2'-C3'	2.10	114.16	109.07
1	B	201	CFZ	O4'-C1'-N1	2.13	112.58	108.08
1	B	203	CFZ	C2-N3-C4	2.15	118.65	115.61
1	B	205	GF2	O3'-C3'-C2'	2.21	119.35	111.68
1	A	103	CFZ	C2-N3-C4	2.26	118.79	115.61
1	A	101	CFZ	C2-N3-C4	2.32	118.88	115.61
1	A	105	GF2	F-C2'-C3'	2.47	115.06	109.07
1	A	105	GF2	C4-C5-N7	3.01	112.25	109.48
1	B	207	GF2	C4-C5-N7	3.19	112.41	109.48
1	B	205	GF2	C4-C5-N7	3.29	112.51	109.48
1	B	201	CFZ	C2-N3-C4	3.33	120.30	115.61
1	B	207	GF2	C6-N1-C2	5.43	123.47	115.94
1	A	107	GF2	C6-N1-C2	7.12	125.82	115.94
1	A	105	GF2	C6-N1-C2	7.35	126.14	115.94
1	B	205	GF2	C6-N1-C2	7.50	126.35	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	205	GF2	5	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	4/8 (50%)	-0.47	0 100 100	29, 31, 35, 46	0
1	B	4/8 (50%)	-0.59	0 100 100	22, 27, 27, 44	0
All	All	8/16 (50%)	-0.53	0 100 100	22, 29, 44, 46	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	GF2	B	207	23/24	0.96	0.07	-	22,24,27,27	0
1	CFZ	B	201	17/21	0.80	0.16	-	46,55,58,60	0
1	GF2	B	205	23/24	0.95	0.12	-	19,24,27,34	0
1	CFZ	A	103	20/21	0.98	0.07	-	21,30,33,34	0
1	GF2	A	107	23/24	0.94	0.10	-	34,39,46,48	0
1	CFZ	A	101	17/21	0.92	0.10	-	29,33,40,41	0
1	GF2	A	105	23/24	0.97	0.07	-	25,29,36,39	0
1	CFZ	B	203	20/21	0.93	0.08	-	28,33,43,44	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.